

PRELIMINARY ASSESSMENT

AMERICAN FORK CANYON

PACIFIC MINE
MARY ELLEN GULCH MINE
LOWER BOG MINE

UINTA NATIONAL FOREST

PLEASANT GROVE RANGER DISTRICT

June 1994

PRELIMINARY ASSESMENT

UPPER AMERICAN FORK CANYON

UINTA NATIONAL FOREST PLEASANT GROVE RANGER DISTRICT

PACIFIC MINE, LOWER BOG MINE, AND MARY ELLEN GULCH MINE

PREPARED BY:

REVIEWED AND RECOMMENDED BY:

DISTRICT RANGER

RESOURCE ASSISTANT

EXECUTIVE SUMMARY

PACIFIC, LOWER BOG, AND MARY ELLEN MINES AMERICAN FORK CANYON, UTAH

The Pacific, Lower Bog, and Mary Ellen mines are located on National Forest System lands on the Uinta National Forest. Each mine has associated tailings piles with ground water running out of the mine adits. This water has been tested periodically, and is known to contain elevated levels of copper, zinc, and cadmium.

The area near the Pacific mine is used by recreationists. OHV (Off Road Vehicle) use occurs on the tailings pile of the Pacific mine. The Lower Bog and Mary Ellen mines are less accessible to publics; however water from these adits still enter the North Fork of American Fork creek.

The Uinta National Forest recommends mitigation and reclamation to varying degrees at each site. This Preliminary Assessment makes no effort to recommend specific techniques. Rather, the P.A. is written to give the reader an overview of the situation at each site along with a brief history, ownership, and condition of sites.

TABLE OF CONTENTS

PART I

General Site Information

Site Name and Location Type of Facility Type of Ownership Site Status Years of Operation Owner/Operator Information Environmental Setting Site Size

Source and Waste Characteristics

Source Types and Locations
Hazardous Substances Present
Ground Water Use and Characteristics
Surface Water use and Characteristics
Surface Water Body Types Within 15 Downstream Miles
Fisheries Within 15 Downstream Miles
Sensitive Environments and Wetlands Within 15 Downstream Miles
Acreage of Wetlands within Four Miles
Soil Exposure Characteristics
Air Pathway Caracteristics

PART II

Preliminary assessment Scoresheet

PART III

Correspondence

PART IV

Sample Results From Water, Soil, and Tailings at the Three Sites

Appendix A

"American Fork Hydrology and Water Quality Study" Lidstone and Anderson, Inc. February 3, 1993

Appendix B


"Aquatic Ecosystem Inventory", Macroinvertebrate Analysis


PART I


General Site Information and Source and Waste Characteristics




Photo circa 1910 pacific mine









GENERAL SITE INFORMATION

CERCLIS ID NUMBER:

UTD 988074951

SITE NAME AND LOCATION:

The site has been identified and will be referred to as the American Fork Mining District Site which is composed of three separate locations which are in close proximity to each other. These sites include the: Mary Ellen Gulch mine and tailings (MEG), Lower Bog mine and tailings (LB), and Pacific mine and tailings (PM). General location is in Utah County, in the Upper American Fork Canyon area. The MEG mine is located in Township 3 South, Range 3 East, NW1/4 of SE1/4 Section 20. The Lower Bog Mine is in Township 3 South, Range 3 East, SW1/4 of SE1/4 Section 16. The Pacific Mine and Tailings are located at Township 3 South, Range 3 East, NW1/4 of SE1/4 Section 22. All legal descriptions are Salt Lake Based Meridian (SLBM).

Ground water is present in all three mines. The water is exposed to mineralized rock, spent ore, and or tailings changing the chemical composition of the water (Lidstone & Anderson, Inc 1993). In the case of the three mines, the water runs out of the adit across tailings piles and into the American Fork River. In addition to containing trace elements picked up in the mine shafts, except at the Lower Bog Mine, the water picks up more contaminants as it passes through the tailings piles. Precipitation events also contribute to the pollution of the American Fork River through surface run-off from the tailings. In both the Pacific and Lower Bog situations, tailings piles at both sites are within 10 feet of the North Fork of American Fork river. This close proximity to surface water allows a high potential for contamination to occur to the river during and after most precipitation events.

The area surrounding the three sites is used throughout much of the year by outdoor enthusiests. Recreational opportunities exist throughout the area including camping, fishing, hunting, off road vehicle use, and exploring. The ability for people get close to and travel virtually unrestricted through old mining operations appeals to many people. The area has a rich mining history that attracts people to it. Unfortunately, people who visit these sites are exposing themselves to more than just the appeal of the area.

Public access to the effluent and tailing piles is generally unrestricted particularly at the Pacific mine. Efforts were made to fence the area but were unsuccessful in restricting all publics from being exposed to the area. The tailings pile at the Pacific Mine is used by Off Highway Vehicles (OHV) as a hill climb and OHV play area.

The Lower Bog mine is less accessible, requiring a short hike or four wheel drive to get close enough to make the 200 yard hike to the foot of the tailings pile. The Mary Ellen Gulch mine is on private land and vehicle access requires travel with high clearance vehicles.

Exposure to the sites has not been directly linked to any health problems however that possibility exists.

TYPE OF FACILITY:

The three sites are facilities associated with early 20th century hard rock mining claims. Silver, Iron ore, and gold were all mined at these sites (Keech). Along with the mining activities, milling also occurred on site, leaving tailing piles at the Pacific and Lower Bog mines (See Attached Photos). Ground water is flowing out of each of the three mine adits at varying flow rates. The ground water is exposed to elevated levels of Zinc, Cadmium, Copper, and Lead (See Appendix A). In addition to the contamination that occurs within the adit, in the case of the Pacific and Mary Ellen mines, the same effluence flows over mine tailings with similar elevated elements.

TYPE OF OWNERSHIP:

MARY ELLEN GULCH MINE: (Survey Number L57, Plat Index Number 392) Sold by Mann Enterprises to William D. Schnack on 8/20/1987. This mine is privately owned and currently not in operation. The water that flows out of the mine adit flows across mine tailings directly into the Mary Ellen Gulch tributary of the American Fork River. Shortly after the adit water enters the Mary Ellen Gulch tributary, (within 300 feet) it enters onto National Forest System lands.

PACIFIC MINE: (Survey Number 5361, Plat Index Number 491 originally known as the Blue Rock #2 claim) the Mine is owned by the Euro-Nevada Mining Corporation, Inc. 6121 Lakeside Drive, Suite 240, Reno, Nevada 89511, (702) 825-8890. The majority of the tailings pile and settling pond exist on National Forest System land.

LOWER BOG MINE: (Survey Number 5422, Plat Index Number 451) Originally patent 5/24/1910. Last owner Lorraine B. Jack et al who sold the land to United States of America on 10/14/1966 and is now National Forest System lands.

SITE STATUS:

MARY ELLEN GULCH MINE: The Mary Ellen mine is currently inactive however, the Globe mine which is adjacent (upstream) to the Mary Ellen Mine is active.

PACIFIC MINE: The Pacific mine is currently inactive.

LOWER BOG MINE: The Lower Bog mine is currently inactive.


YEARS OF OPERATION:

Each of the mines have been reviewed by Uinta National Forest Archeologist for cultural and historical significance and are all eligible for National Historic Register status.

MARY ELLEN GULCH: The Mary Ellen gulch mine was __cated in 1870. A patent was filed for operation in 1876. Activity occured periodocially through 1959.

PACIFIC MINE: Formally known as the Blue Rock #2 was located in 1903. At this time, there was evidence of three tunnels prior to location. Activity at this


Lower Bay Mine American fork Canyon Vinta National Estest Pleasant Grove Ranger Pistrict November 04, 1993 1040 hours Tim Garcia & Paul Skabland



THE

(Approximate scale)

* 0.2 GPM DISCHARGE WH 44.5 GPM DISCHARGE




120,000 fe 2



PACIFIC MINE 6-94 Adit discharge running through tailings.



PACIFIC MINE 6-94 Beaver pond (foreground) Pacific tailings



LOWER BOG MINE 10/93

ADIT discharge (right) entering North Fork American Fork stream. Tailings pile (left)

LOWER BOG MINE 10/93

"Yellow Boy" discolored sediment entering North Fork American Fork River



mine was at it's height between 1910 and the late 1940's. There has been a resurgence of interest in making further explorations of this mine in the last decade by it's current owner; however no significant work has been done since the 1940's.

LOWER BOG: The Bog mine was located in 1895 by Ed Hines. Initial surveys were conducted in 1905 with actual work begining in 1914. Active mining occured through the 1940's and finally operations shut down in the late 1940's. Some prospecting occured later in the 1970's however the majority of activity occured between 1914 and the late 1940's

OWNER/OPERATOR INFORMATION:

MARY ELLEN GULCH: William D. Schnack c/o Associated Title Co., P.O. Box 478, Salt Lake City , UT,84110-0478, Attn: Lyle Swenson

PACIFIC MINE: Euro-Nevada Mining Corporation, Inc. 6121 Lakeside Drive, Suite 240, Reno, Nevada 89511, (702) 825-8890 owns the mine and some tailings however, the majority of the tailings pile and settling pond exist on National Forest System land.

LOWER BOG: United States of America, National Forest System Lands.

ENVIRONMENTAL SETTING:

MARY ELLEN GULCH: The Mary Ellen Gulch mine is located at approximately 9,000 feet above sea level. Dominant vegetation types include upper elevation riparian, alpine spruce/fir type and high elevation mountain brush.

PACIFIC MINE: The Pacific Mine is located in the bottom of the North Fork of the American Fork Canyon at approximately 7800 feet AMSL. Vegetation consists of mixed conifer stands to the west and riparian vegetation skirting the east perimeter of the tailings pile and settling pond. The North Fork of the A.F. river runs within 10 feet of the tailings around the east side of the mine tailings area.

LOWER BOG: The Lower Bog is located along a stream corridor consisting of associated high elevation riparian vegetation types. The adit is in a high elevation mountain brush zone.

APPROXIMATE SIZE OF SITE:

MARY ELLEN GULCH

PACIFIC MINE: Operations at the Pacific mine cover an area of approximately 120,000 square feet. The majority of this area is v d as a tailings and settling pond. The average depth of the tailings around the area is estimated at approximately five feet. The total volume of the tailings has been estimated at 600,000 cubic feet of tailings containing elevated levels of zinc, cadmium, lead, and copper. There are remains of buildings associated with the Pacific mine operation however; no intact structures are present.

LOWER BOG: Groundwater discharge and tailings pile make up the facility at the Lower Bog mine. The area associated with the mine involves about 6900 square feet. The average depth of the tailings is approximately 10 feet, with total volume being approximately 69,000 cubic feet. There are no facilities associated with the Lower Bog mine.

SOURCE AND WASTE CHARACTERISTICS:

SOURCE TYPES AND LOCATIONS:

MARY ELLEN GULCH: Groundwater discharge is the primary source of contamination in the Mary Ellen Gulch location. Groundwater surfacing from the adit contains elevated levels of zinc, iron, copper, lead, and cadmium. The Mary Ellen Gulch Mine is located along a south east flowing tributary drainage to the North Fork of the American Fork River at an elevation of 9,100 feet. The site has several portals, tailings and waste rock piles. The North portal has a pH of 5.95, while the south portal has a 7.2 pH. The North Portal discharges 70 GPM (Gallon Per Minute) with the south portal discharging only 2.5 GPM (Lidstone & Anderson 1993).

PACIFIC MINE: There are two major waste characteristics involved at the Pacific mine site. The first is the extensive tailings pile and settling pond associated with past mining activities. Dust transported by wind and precipitation run-off are both causes for the spread of these tailings from the site. Tailings and the settling pond are both within a distance of 10 to 50 feet from the American Fork river. The second Source of pollution is ground water discharge from the Pacific mine adit itself. 144 GPM discharge with a pH of 6.5 was measured from the Pacific mine portal with elevated levels of lead, zinc, copper, and cadmium (Lidstone & Anderson 1993).

LOWER BOG: The Lower Bog mine has an elevation of about 8500 feet. The site consists of a single bedrock opening, tailings dump, and miscellaneous spoil piles. Discharge from the adit is approximately 44 GPM with "yellow boy" or hydrous iron oxide deposits around the area of discharge. pH levels were measured at 5.1 with total disolved solids at 80 parts per million (PPM). 1992 samples indicate elevated levels of iron, cadmium, zinc, copper, and lead. Discharge from the mine adit flow boths around both sides of the tailings located below the mine opening (Lidstone and Anderson 1993).

HAZARDOUS SUBSTANCES PRESENT:

The following elements identified exceed aquatic standards:

MARY ELLEN GULCH: Elevated levels of zinc and iron are present (Mangum, 1988).

PACIFIC MINE: Elevated levels of lead, cadmium, zinc, and copper are present (Mangum, 1988).

LOWER BOG: Elevated levels of lead, cadmium, zinc, copper, and iron are present (Mangum, 1988).

Testing of each site has occured on several occasions. Results of these tests can be seen in section IV of this text.

GROUND WATER USE AND CHARACTERISTICS

General Narrative:

Sources of contamination are poorly contained. The tailings are not enclosed allowing infiltration to freely occur. Ground water that is discharged from the mine adit is being discharged already contaminated (Mangum, 1988). The source is less likely to contaminate ground water than it is to contaminate near by surface water. The waste quantities at any one of the three would not be considered particularly large; however the waste at all three sites combined would be considered large.

Annual precipitation for all three areas is approximately 40 inches annually. Much of the precipitation comes in the form of snow between the months of November and April. Infiltration rates at all three areas would not be considered exceptionally high; but rather should be considered average with none of the areas having evidence of karst terrain.

PRIVATE WELLS WITHIN 4 MILES: There are no known private wells within four miles of any of the three mines sites identified. The areas downstream from the Pacific mine particularly is a popular site for camping and fishing. Campers, upon occasion may still drink directly form the American Fork River directly below the Pacific mine tailings.

SURFACE WATER USE AND CHARACTERISTICS

DISTANCE TO NEAREST SURFACE WATER:

MARY ELLEN GULCH: The closest surface water to the Mary Ellen adit is within 30 feet. The effluent from the adit flows down across mine waste and directly into the Mary Ellen Gulch tributary of the American Fork River.

PACIFIC MINE: The tailings pile and settling pond is within 10 feet of the American Fork River. During precipitation events, run off will flow directly across the tailings and into the river. The effluent from the Pacific mine adit flows into a wetland area created by beaver activity. This beaver pond captures some of the contaminants preventing a strong solution from entering the American Fork stream channel (Lidstone & Anderson, 1993). However there is evidence that some elements enter the stream.

LOWER BOG MINE: Tailings from the Lower Bog mine are within 3 feet of the main channel of the American Fork River. In addition to the exposure of surface water, adit discharge runs over and around the tailings. Either adit discharge or springflow flows beneath the tailings pile and enters the stream from beneath the mine tailings.

SURFACE WATER BODY TYPES WITHIN 15 DOWNSTREAM MILES

Tibble Fork Reservoir is approximately 7 downstream miles from the lowest site (Mary Ellen Gulch). It is used as a flood control structure. Water collected there is also used for agricultural irrigation in the Utah County area. No evidence has been collected indicating the contamination of Tibble Fork

Reservoir as a result of these sites. Evidence in fact shows little effects of the contaminants less than a mile down stream from the lowest source.

FISHERIES WITHIN 15 DOWNSTREAM MILES:

All three mines are located in the American Fork drainage. The American Fork river, including Tibble Fork Reservior is a put and take fishery managed primarily for rainbow trout. Secondary management is for brown and cutthroat trout. The Utah Division of Wildlife Resources (DWR) stocks approximately 35,500 fish a year in the stream reach from Mary Ellen Gulch to the mouth of American Fork Canyon, which is a distance of approximately 11.6 stream miles.

No studies have monitored fish downstream of the mines for contaminants. It is not known if, or at what levels fish retain contaminants from the mines. Many of the planted fish do not overwinter and spawn. A small, but important native cutthroat trout population does overwinter and spawn in this drainage. The majority of fish caught in the American Fork river have been in the drainage less than one year. Fisherpersons commonly keep and eat the fish they catch.

Quantifying the actual number of recreation fishing hours on the American Fork river is difficult, but the DWR manages the American Fork river as a "heavy use" area and has a goal of 500 angler-hours/acre/year.

Numerous log structures designed to enhance fish habitat have been installed along the upper reaches of the American Fork River. Rainbow trout congregate in the pools below these structures and encourage fishing below the discharge of the three mines. Tibble Fork Reservoir was built as a sediment trap and traps sediment associated with the discharge from the sites. Dissolved pollutants may travel below the reservoir.

SENSITIVE ENVIRONMENTS AND WETLANDS WITHIN 15 DOWNSTREAM MILES:

SOIL EXPOSURE CHARACTERISTICS:

General Narrative

Soil effects are localized and restricted to immediately around each of the three sites. Little evidence has been gathered indicating effects to the soil resources.

AIR PATHWAY CHARACTERISTICS:

General Narrative

Effect of the air pathway is localized at all three sites. Localized wind at each site has the potential to transport contaminated tailing dust within a close proximity of each site. The threat of air pathway contamination is not fully known. Dust from these areas has been witnessed by individuals and seems to be the only threat to the air pathway.

LOCATIONS OF SENSITIVE ENVIRONMENTS WIHTIN 4 MILES:

ACREAGE OF WETLANDS WITHIN 4 MILES:

Wetlands

The entire watershed within a radius of 1/4 and 1/2 miles of the Pacific mine drains into the North Fork of American Fork Creek. A wetland approximately 2 acres in size is associated with a beaver pond in the stream. The beaver pond is within 1/4 mile of the Pacific Mine. Approximately 4 acres of sensitive environments (riparian areas) exist along the stream channel. Two acres in the 1/4 mile radius and 2 acres within the 1/2 mile radius. No other wetlands or sensitive environments occur within 1/2 mile of the Pacific Mine.

	ONSITE	1/4 mi.	1/2 mi.
Wetlands	0.1 acres	2 acres	0 acres
Sensitive Env.	0.2 acres	2 acres	2 acres
Total	2.1 acres	4 acres	2 acres

PART II Preliminary Assessment Scoresheet

APPENDIX A

OMB Approval Number: 2050-0095 Approved for Use Through: 1/92

PA Scoresheets

Investigator: Timothy Garcia

Agency/Organization: USDA Forest Service

Street Address: 88 West 100 North

City/State/Zip: Provo, UT 84601

Date: January 18, 1994

GENERAL INFORMATION

Site Description and Operational History:

REFER TO PART I IN GENERAL SITE INFORMATION

Probable Substances of Concern: (Previous investigations, analytical data)

REFER TO PART IV AND APPENDICIES A THROUGH-D

GENERAL INFORMATION (continued)

Site Sketch: (Show all pertinent features, indicate sources and closest targets, indicate north) REFER TO PART I GENERAL SITE INFORMATION

SOURCE EVALUATION

Source Description:

Source Consists of tailings pile, two contaminated ground water discharge and spoil piles.

Source Description:

Source Description:

Source Description:

120,000 square feet total X average depth of tailings of 5 feet = 600,000 cubic feet of waste.

Source Name: MARY ELLEN GULCH
Source Description:

Tailing and waste rock piles; spoils dump, and contaminated ground water discharge

Source 03 | Source Name: LOWER BOG | Source Waste Quantity (WQ) Calculations: 6900 |

Source Description:

Tailings pile, miscellaneous spoils piles, and contaminated ground water discharge

Site WC:

PA TABLE 1: WASTE CHARACTERISTICS (WC) SCORES

PA Table 1s: WC Scores for Single Source Sites and Formulas for Multiple Source Sites

T		SINGLE	SOURCE SITES (assigned WC	scores)	MULTIPLE SOURCE SITES
ER	SOURCE TYPE	WC = 18	WC = 32	WC = 100	Formula for Assigning Source WQ Values
CORST-TOWNT	N/A	≤100 lb	>100 to 10,000 to	>10,000 њ	15 + 1
E ben Herstein ba b &	N/A	≤500,000 lb	> 500,000 to 50 million lb	>50 million lb	4b + 5,000
	Landfill	≤6.75 million h³ ≤250,000 yd³	> 6.75 million to 675 million ft ³ > 250,000 to 25 million yd ³	> 675 million ft ³ > 25 million yd ³	$tt^3 + 67,500$ $ya^3 + 2,500$
	Surface impoundment	≤8,750 ft ³ ≤250 yd ³	>6,750 to 675,000 ft ³ >250 to 25,000 yd ³	> 875,000 ft ³ > 25,000 yd ³	$tt^3 + 67.5$ $yd^3 + 2.5$
v	Drums	≤1,000 drums	>1,000 to 100,000 drume	>100,000 drume	drums + 10
0	Tanks and non- drum containers	≤ 50,000 gallens	>50,000 to 5 million gallons	>5 million gallons	gallons + 500
M	Contaminated soil	≤6.75 million ft ³ ≤250,000 yd ³	>6.75 million to 675 million ft ³ >250,000 to 25 million yd ³	> 675 million ft ³ > 25 million yd ³	$tt^3 + 67,500$ $yd^3 + 2,500$
	Pile	≤6,750 ft ³ ≤250 yd³	>8,750 to 875,000 ft ² > 250 to 25,000 yd ²	>675,000 ft ³ >25,000 yd ³	ft ³ + 67.5 yd ³ + 2.5
	Other	≤8,750 ft ³ ≤250 yd ³	>6,750 to 675,000 ft ² >250 to 25,000 yd ⁵	> 675,000 ft ³ > 25,000 yd ³	$ft^3 + 67.5$ $ya^3 + 2.5$
	Landfill	≤340,000 ft ³ ≤7.8 acres	>340,000 to 34 million ft ² >7.8 to 780 scree	>34 million ft ² >780 scree	ft ² + 3,400 acres + 0.078
	Surface impoundment	≤1,300 ft ³ ≤0.029 scree	>1,300 to 130,000 ft ² >0.029 to 2.9 acree	>130,000 ft ² >2.9 scree	ft ² + 13 acres + 0.00029
RE	Contaminated soil	≤3.4 million ft ³ ≤78 scree	>3.4 million to 340 million ft ³ >78 to 7,800 ecree	>340 million ft ³ >7,800 scree	ft ² + 34,000 acres + 0.78
A	Pile*	≤1,300 ft ³ ≤0,029 scree	>1,300 to 130,000 ft ² >0,025 to 2.5 acres	>130,000 ft ² >2.5 sores	ft ² + 13 acres + 0.00029
	Land treatment	≤27,000 h² ≤0.62 acres	>27,000 to 2.7 million ft ² >0.62 to 62 acres	>2.7 million ft ² >62 scree	$tt^2 + 270$ acres + 0.0062

1 ton = 2,000 lb = 1 yds = 4 drums = 200 gallons

* Use area of land ourface under pile, not surface area of pile.

PA Table 1b: WC Scores for Multiple Source Sites

WQ Total	WC Soore
>0 to 100	18
> 100 to 10,000	32
>10,000	100

GROUND WATER PATHWAY GROUND WATER USE DESCRIPTION

See	Ground water	section	in	narrative	

Calculations for Drinking Water Populations Served by Ground Water:

11

100

SUSPECTED RELEASE	PRIMARY TARGETS
Y N U s o n _ s k	Y N U o o n s k S X Is any drinking water well nearby? Has any nearby drinking water well been closed? Has any nearby drinking water user reported foul-testing or foul-smelling water? Does any nearby well have a large drawdown or high production rate? XX Is any drinking water well located between the site and other wells that are suspected to be exposed to a hazardous substance?
conductive? Is drinking water drawn from a shallow aquifer? Are suspected contaminants highly mobile in ground water? Does analytical or circumstantial evidence suggest ground water contamination? Other criterie? SUSPECTED RELEASE?	□ ☑ □ Does analytical or circumstantial evidence suggest contamination at a drinking water well? □ ☑ □ Does any drinking water well warrant sampling? □ □ Other criteria? □ □ PRIMARY TARGET(S) IDENTIFIED?
Summerize the rationale for Suspected Release (attach an additional page if necessary):	Summarize the rationale for Primary Targets (attach-an additional page if necessary):

GROUND WATER PATHWAY SCORESHEET

	Pathway Characteristics			
	Do you suspect a release [see Ground Water Pathway Criteria List, page 7)? Is the site located in karst terrain? Depth to aquifer:	Yes Yes	- April	
	Distance to the nearest drinking water well:		ft	
		A	в.	
LIKELIH	OOD OF RELEASE	Suspected	Ne Suspected Release	
		Jeed	Anna San	References
	ECTED RELEASE: If you suspect a release to ground water (see page 7), in a score of 550. Use only column A for this pathway.		1906 m Jacq	
the si	USPECTED RELEASE: If you do not suspect a release to ground water, and te is in karst terrain or the depth to aquifer is 70 feet or less, assign a score 0; otherwise, assign a score of 340. Use only column B for this pathway.		340	
	LR -		340	
TARGET	s			
drinkir	ARY TARGET POPULATION: Determine the number of people served by ng water wells that you suspect have been exposed to a hazardous since from the site (see Ground Water Pathway Criteria List, page 7).	0		
drinkir	NDARY TARGET POPULATION: Determine the number of people served by ng water wells that you do NOT suspect have been exposed to a hazardous ince from the site, and assign the total population score from PA Table 2.		-500	
	Are any wells part of a blended system? YesNdXX	0	0	
water,	EST WELL: If you have identified a primary target population for ground assign a score of 50; otherwise, assign the Nearest Well score from ble 2. If no drinking water wells exist within 4 miles, assign a score of zero.	0	0	
		136 6 4 6	120, 5, 41 11	TOP:
or if y	HEAD PROTECTION AREA (WHPA): If any source lies within or above a WHPA, ou have identified any primary target well within a WHPA, assign a score of 20; 5 if neither condition holds but a WHPA is present within 4 miles; otherwise	0	0	
assign	zero.	n = a	is-a	
7. RESO	JRCES		5	
	т-		5	
WASTE	CHARACTERISTICS			
ch	you have identified any primary target for ground water, assign the waste aracteristics score calculated on page 4, or a score of 32, whichever is IEATER; do not evaluate part 8 of this factor.	(100 to 20)		
	you have NOT identified any primary target for ground water, assign the isse characteristics score calculated on page 4.	(104,35, or 106	100	
	wc =		445	
GROUND	WATER PATHWAY SCORE: LR x T x WC 82,500	(subject to a m	6	

PA TABLE 2: VALUES FOR SECONDARY GROUND WATER TARGET POPULATIONS

PA Table 2a: Non-Karst Aquifers

		Nearest			Popu	lation Se	rved by W	ells Withi	n Distance	Categor	Y		En elinis e
Distance from Site	Population	Well (choose highest)	1 to 10	11, te 30	31 te :	101 10 300	301 to 1,000	1,001 to 3,000	3,001 te 10,000	10,001 te 20,000	30,001 te 100,000	Greeter then 100,000	Population Value
0 to X mile	0	20	1	2	5	16	52	163	521	1,633	5,214	16,325	
> % to % mile	0	18	1	1	3	10	32	101	323	1,012	3,233	10,121	
> 1/4 to 1 mile	0 0	9	1	1	2	5	17	52	167	522	1,668	5,224	
>1 to 2 m a		5	1	1	1	3	9	29	94	294	939	2,938	
> 2 to 3 miles		3	1	1	1 .	2	. 7	21	68	212	678	2,122	
>3 to 4 miles	0	2	1	i	1	1	4	13	42	131	417	1,306	1
	Nearest Well =	2										Score =	1

PA Table 2b: Karst Aquifers

		Nearest	anya — 11		Рори	lation Sei	ved by W	ells With	n Distanc	e Categor	γ		7
Distance from Site	Population	(use 20 for karst)	1 to 10	11 to 30	31 to 100	101 to 300	301 te 1,000	1,001 to 3,000	2,001 to 10,000	10,001 to 30,000	30,001 te 100,000	Orester then 100,000	Population Value
O to ¼ mile	0	20	1	2	5	16	52	163	521	1,633	5,214	16,325	
> % to % mile	0	20	1	1	3	10	32	101	323	1,012	3,233	10,121	
> % to 1 mile	0	20	1	1	3	. 8	26	82	261	816	2,607	8,162	
> 1 to 2 miles	. 0	20	1	1	3	8	26	82	261	816	2,607	8,162	
> 2 to 3 miles	0	20	1	1	3	8	26	82	261	816	2,607	8,162	
> 3 to 4 miles	0	20	1	1	3	8	26	82	261	816	2,607	8,162	
	Nearest Well =										5	Score =	

SURFACE WATER PATHWAY MIGRATION ROUTE SKETCH

Suface Water Migration Route Sketch: (include runoff route, probable point of entry, 15-mile target distance limit, intakes, fisheries, and sensitive environments)

			SUSPECTED RELEASE				PRIMARY TARGETS
DB D	0000000000	000000000000000000000000000000000000000	THE PARTY OF THE P	Y	200 80 0 0 0 0	U e k 0	Drinking water intake Fishery Does analytical or circumstantial evidence suggest surface water contamination at or downstream of a target? Drinking water intake Fishery Does analytical or circumstantial evidence suggest surface water contamination at or downstream of a target? Does any target warrent sampling? If yes: Prinking water intake Fishery Sensitive environment Other criteria? PRIMARY INTAKE(S) IDENTIFIED?
20			Is ground water discharge to surface water likely? Does analytical or circumstantial evidence suggest surface water contamination? Other criteria?		0		PRIMARY SENSITIVE ENVIRONMENT(S) IDENTIFIED?
Su			s the rationals for Suspected Release (attach an page if necessary):	2.002			the rationale for Primery Targets (attach an page if necessary):

SURFACE WATER PATHWAY LIKELIHOOD OF RELEASE AND DRINKING WATER THREAT SCORESHEET

Pathway Characteristics

Do you suspect a release (see Surface Water Pathway Criteria List, page 11)? Distance to surface water: Flood frequency:

SUSPECTED RELEASE: If you suspect a release to surface water (see page 11), assign a score of \$50. Use only column A for this pathway. NO SUSPECTED RELEASE: If you do not suspect a release to surface water, use the table below to assign a score based on distance to surface water and flood frequency. Use only column B for this pathway. Distance to surface water > 2,500 feet	KELIHOOD OF RELEASE		Suspected Release	Ne Suspected Release	Refere
NO SUSPECTED RELEASE: If you do not suspect a release to surface water, use the table below to assign a score based on distance to surface water and flood frequency. Use only column 8 for this pathway. Distance to surface water > 2.500 feet			550		
Distance to surface water > 2.500 feet, and Site in annual or 10-year floodplain Site in 100-year floodplain Site in 500-year floodplain Site in 500-year floodplain Site outside 500-year floodplain Site outside 500-year floodplain LR = 550 LR = 550 Indicators in 100 year floodplain Indicato	water, use the table below to assign a score based on	distance to surface		, 508. 408. 200 ± 1001	
Site in 100-year floodplain 4-08 Site in 100-year floodplain 300 Site in 100-year floodplain 300 Site outside 500-year floodplain 700 LR = 550 RENKING WATER THREAT TARGETS Record the water body type, flow (if applicable), and number of people served by each drinking water intake within the target distance limit. If there is no drinking water intake within the target distance limit, If there is no drinking water intake within the target distance limit, If there is no drinking water intake within the target distance limit, factors 4, 5, and 6 each receive zero scores.	The state of the s	500			
Site in 100-year floodplain 300 Site outside 500-year floodplain 100 LR = 550 RINKING WATER THREAT TARGETS Record the water body type, flow (if applicable), and number of people served by each drinking water intake within the target distance limit. If there is no drinking water intake within the target distance limit, factors 4, 5, and 6 ach receive zero scores. PRIMARY TARGET POPULATION: If you suspect any drinking water intake listed above has been exposed to a hazardous substance from the site (see Surface Water Pathway Criteria List, page 11), list the intake name(s) and calculate the factor score based on the total population served. O people x 10 = 0 SECONDARY TARGET POPULATION: Determine the number of people served by drinking water intakes that you do NOT suspect have been exposed to a hazardous substance from the site, and assign the total population score from PA Table 3. Are any intakes part of a blended system? Yes No If yes, attach a page to show apportionment calculations. NEAREST INTAKE: If you have identified a primary target population for the drinking water threat (factor 4), assign a score of 50; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of 50; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of 50; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of 50; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of 50; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of 50; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit assign a score of 50		500	The second second		
Site in 500-year floodplain 100 LR = 550 LR = 550 Item. watter body type, flow (if applicable), and number of people served by each drinking water intake within the target distance limit. If there is no drinking water intake within the target distance limit, actors 4, 5, and 6 each receive zero scores. Mater Body Type Row People Served cach receive zero scores. Describe Name Water Body Type Row People Served cach receive zero scores. Describe Name PRIMARY TARGET POPULATION: If you suspect any drinking water intake listed above has been exposed to a hazardous substance from the site see Surface Water Pathway Criteria List, page 11), list the intake name(s) and calculate the factor score based on the total population served. Decople x 10 = 0 SECONDARY TARGET POPULATION: Determine the number of people served by drinking water intakes that you do NOT suspect have been exposed to a hazardous substance from the site, and assign the total population score from PA Table 3. Are any intakes part of a blended system? Yes No O People x 10 type, attach a page to show apportionment calculations. NEAREST INTAKE: If you have identified a primary target population for the drinking water threat flactor 4; assign a score of 50; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of 50; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of 20; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of 20; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of 20; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of 20; otherwise, assign the Nearest Intake Name Page 10;			100	4 7	
Site outside 500-year floodplain LR = 550 Index out 100 100		ATT DESCRIPTION ASSESSMENT OF THE PARTY OF T			
Record the water body type, flow (if applicable), and number of people served by each drinking water intake within the target distance limit. If there is no drinking water intake within the target distance limit, actors 4, 5, and 6 each receive zero scores. Intake Name					
REINKING WATER THREAT TARGETS Record the water body type, flow (if applicable), and number of people served by each drinking water intake within the target distance limit. If there is no drinking water intake within the target distance limit, factors 4, 5, and 6 each receive zero scores. Intake Name			E CO	\$400, 4000, 3000 or 1,000	
RENKING WATER THREAT TARGETS Record the water body type, flow (if applicable), and number of people served by each drinking water intake within the target distance limit. If there is no drinking water intake within the target distance limit, factors 4, 5, and 6 each receive zero scores. Intake Name Water Body Type Flow People Served		LR	550		
Record the water body type, flow (if applicable), and number of people served by each drinking water intake within the target distance limit. If there is no drinking water intake within the target distance limit, factors 4, 5, and 6 each receive zero scores. Intake Name					
Record the water body type, flow (if applicable), and number of people served by each drinking water intake within the target distance limit. If there is no drinking water intake within the target distance limit, factors 4, 5, and 6 each receive zero scores. Intake Name	RINKING WATER THREAT TARGETS '				
above has been exposed to a hazardous substance from the site (see Surface Water Pathway Criteria List, page 11), list the intake name(s) and calculate the factor score based on the total population served. O	each receive zero scores.	Row People Servedctscts			
SECONDARY TARGET POPULATION: Determine the number of people served by drinking water intakes that you do NOT suspect have been exposed to a hazardous substance from the site, and assign the total population score from PA Table 3. Are any intakes part of a blended system? Yes No 19 yes, attach a page to show apportionment calculations. NEAREST INTAKE: If you have identified a primary target population for the drinking water threat (factor 4;, assign a score of 50; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of zero. RESOURCES 5	PRIMARY TARGET POPULATION: If you suspect any				
drinking water intakes that you do NOT suspect have been exposed to a hazardous substance from the site, and assign the total population score from PA Table 3. Are any intakes part of a blended system? Yes No 0 If yes, attach a page to show apportionment calculations. NEAREST INTAKE: If you have identified a primary target population for the drinking water threat (factor 4; assign a score of 50; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of zero.	Pathway Criteria List, page 11), list the intake name(s)	n the site (see Surface Water			
If yes, attach a page to show apportionment calculations. NEAREST INTAKE: If you have identified a primary target population for the drinking water threat (factor 4; assign a score of 50; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of zero.	Pathway Criteria List, page 11), list the intake name(s)	n the site (see Surface Water and calculate the factor	- 0		
NEAREST INTAKE: If you have identified a primary target population for the drinking water threat (factor 4), assign a score of 50; otherwise, assign the Nearest Intake score from PA Table 3. If no drinking water intake exists within the target distance limit, assign a score of zero. RESOURCES 5	Pathway Criteria List, page 11), list the intake name(s) score based on the total population served. SECONDARY TARGET POPULATION: Determine the drinking water intakes that you do NOT suspect have be	n the site (see Surface Water and calculate the factor O people x 10 number of people served by seen exposed to a hazardous	- 0		
RESOURCES 5	Pathway Criteria List, page 11), list the intake name(s) score based on the total population served. SECONDARY TARGET POPULATION: Determine the drinking water intakes that you do NOT suspect have be substance from the site, and assign the total population. Are any intakes part of a blended system? You	D people x 10 number of people served by seon exposed to a hazardous a score from PA Table 3.	0		
22	Pathway Criteria List, page 11), list the intake name(s) score based on the total population served. SECONDARY TARGET POPULATION: Determine the drinking water intakes that you do NOT suspect have be substance from the site, and assign the total population. Are any intakes part of a blended system? You fixe, attach a page to show apportionment can be needed. NEAREST INTAKE: If you have identified a primary tark drinking water threat (factor 4), assign a score of 50; on Nearest Intake score from PA Table 3. If no drinking water threat score from PA Table 3. If no drinking water threat score from PA Table 3. If no drinking water threat factor 4 is assign a score of 50; on the score from PA Table 3. If no drinking water threat factor 4 is assign a score of 50; on the score from PA Table 3. If no drinking water threat factor 4 is assign a score of 50; on the score factor 4 is assign a score of 50; on the score factor 4 is assign a score of 50; or 50	D people x 10 number of people served by seen exposed to a hazardous a score from PA Table 3. The second people served by seen exposed to a hazardous a score from PA Table 3. The second people served by seen exposed to a hazardous a score from PA Table 3. The second people served by seen exposed to a hazardous a score from PA Table 3.	0 (64.34.16.2.1., # 38	- (M. G.L.), = 0	
т - 55	Pathway Criteria List, page 11), list the intake name(s) score based on the total population served. SECONDARY TARGET POPULATION: Determine the drinking water intakes that you do NOT suspect have be substance from the site, and assign the total population. Are any intakes part of a blended system? You have identified a primary tardrinking water threat (factor 4), assign a score of 50; of Nearest Intake score from PA Table 3. If no drinking water target distance limit, assign a score of zero.	D people x 10 number of people served by seen exposed to a hazardous a score from PA Table 3. The second people served by seen exposed to a hazardous a score from PA Table 3. The second people served by seen exposed to a hazardous a score from PA Table 3. The second people served by seen exposed to a hazardous a score from PA Table 3.	0 (MARIELL - 00		-
	Pathway Criteria List, page 11), list the intake name(s) score based on the total population served. SECONDARY TARGET POPULATION: Determine the drinking water intakes that you do NOT suspect have be substance from the site, and assign the total population. Are any intakes part of a blended system? You have identified a primary tardrinking water threat (factor 4), assign a score of 50; of Nearest Intake score from PA Table 3. If no drinking water target distance limit, assign a score of zero.	D people x 10 number of people served by seen exposed to a hazardous a score from PA Table 3. The second people served by seen exposed to a hazardous a score from PA Table 3. The second people served by seen exposed to a hazardous a score from PA Table 3. The second people served by seen exposed to a hazardous a score from PA Table 3.	0 (MARIELL - 00		

PA TABLE 3: VALUES FOR SECONDARY SURFACE WATER TARGET POPULATIONS

	200	Nearest			P	opulation	Served by	Intakes	Within Flo	w Catego	ry			
Surface Water Body Flow (see PA Table 4)	Population	Intake (choose highest)	1 to 30	31 10 100	101 10 300	1,000	1,001 to 3,000	3,001 to 10,000	10,001 to 30,000	30,001 te 100,000	100,001 te 300,000	300,001 te 1,000,000	Olwater than 1,000,000	Population Value
<10 cfs	-	20	2	5	16	52	163	521	1,633	5,214	16,325	52,138	163,246	
10 to 100 cfs	0	2	1	1	2	5	16	52	163	521	1,633	5,214	16,325	1
> 100 to 1,000 cfs		1	0	0	1	1	2	6	16	52	163	521	1,633	
> 1,000 to 10,000 afs		0	0	0	0	. 0	1	1	2	5	18	52	163	
>10,000 cfs or Great Lakes	-	0	0	0	0	0	. 0	0	1	1	2	5	16	
3-mile Mixing Zone		10	1	3	8	26	82	261	816	2,607	8,162	26,068	81,663	
Neare	st Intake =	2										s	core =	. 1

PA TABLE 4: SURFACE WATER TYPE / FLOW CHARACTERISTICS WITH DILUTION WEIGHTS FOR SECONDARY SURFACE WATER SENSITIVE ENVIRONMENTS

Type of Si	wlace Wa	ter Body	Dilution
Water Body Type	OR	Flow	Weight
minimal stream		< 10 cfs	1
small to moderate stream		10 to 100 cfs	0.1
moderate to large stream		> 100 to 1,000 cfs	N/A
large stream to river		> 1,000 to 10,000 cfs	N/A
large river		> 10,000 cfs	N/A
3-mile mixing zone of quiet flowing streams or rivers		10 ofs or greater	N/A
coastal tidal water (herbors, sounds, bays, stc.), ocean, or Great Lakes	111	N/A	N/A

SURFACE WATER PATHWAY (continued) HUMAN FOOD CHAIN THREAT SCORESHEET

HUMAN FOOD CHAIN THREAT TARGETS 8. Record the water body type and flow (if applicable) for each fishery within the target distance limit. If there is no fishery within the target distance limit, assign a Targets score of 0 at the bottom of the page. Fishery Name	HUMAN FOOD CHAIN THREAT TARGETS 8. Record the water body type and flow (if applicable) for each fishery within the target distance limit, assign a Targets score of 0 at the bottom of the page. Fishery Name	Heare Reference
8. Record the water body type and flow (if applicable) for each fishery within the target distance limit. It there is no fishery within the target distance limit, assign a Targets score of 0 at the bottom of the page. Fishery Name	8. Record the water body type and flow (if applicable) for each fishery within the target distance limit. If there is no fishery within the target distance limit, assign a Targets score of 0 at the bottom of the page. Fishery Name	8.300 w 100)
the target distance limit. If there is no fishery within the target distance limit, assign a Targets score of 0 at the bottom of the page. Fishery Name	the target distance limit. If there is no fishery within the target distance limit, assign a Targets score of 0 at the bottom of the page. Fishery Name	
NORTH FORK OF AF RIVER RIVER # 6.7 cfs	North Fork of AF River River * 6.7 cfs	
Cfs	Cfs c	
PRIMARY FISHERIES: If you suspect any fishery listed above has been exposed to a hazardous substance from the site (see Surface Water Criteria List, page 11), assign a score of 300 and do not evaluate Factor 10. List the primary fisheries: AF River SECONDARY FISHERIES If you suspect a release to surface water and have identified a secondary fishery but no primary fishery, assign a score of 210. If you do not suspect a release, assign a Secondary Fisheries score from the table below using the lowest flow at any fishery within the target distance limit. Lowert Flow Secondary Fisheries Score (< 10 cfs 210 10 to 100 cfs 30) 100 cfs, coastal 11dal waters, oceans, 12	PRIMARY FISHERIES: If you suspect any fishery listed above has been exposed to a hazardous substance from the site (see Surface Water Criteria List, page 11), assign a score of 300 and do not evaluate Factor 10. List the primary fisheries: AF River SECONDARY FISHERIES If you suspect a release to surface water and have identified a secondary lishery but no primary fishery, assign a score of 210. If you do not suspect a release, assign a Secondary Fisheries score from the table	
PRIMARY FISHERIES: If you suspect any fishery listed above has been exposed to a hazardous substance from the site (see Surface Water Criteria List, page 11), assign a score of 300 and do not evaluate Factor 10. List the primary fisheries: AF River SECONDARY FISHERIES If you suspect a release to surface water and have identified a secondary lishery but no primary fishery, assign a score of 210. If you do not suspect a release, assign a Secondary Fisheries score from the table below using the lowest flow at any fishery within the target distance limit. Lower flow Secondary Fisheries Score (10 cfs 210 10 to 100 cfs 30 10 to 100 cfs 30 10 to 100 cfs, coastal tidal waters, oceans, 12	PRIMARY FISHERIES: If you suspect any fishery listed above has been exposed to a hazardous substance from the site (see Surface Water Criteria List, page 11), assign a score of 300 and do not evaluate Factor 10. List the primary fisheries: AF River SECONDARY FISHERIES If you suspect a release to surface water and have identified a secondary lishery but no primary fishery, assign a score of 210. If you do not suspect a release, assign a Secondary Fisheries score from the table	
PRIMARY FISHERIES: If you suspect any fishery listed above has been exposed to a hazardous substance from the site (see Surface Water Criteria List, page 11), assign a score of 300 and do not evaluate Factor 10. List the primary fisheries: AF River SECONDARY FISHERIES If you suspect a release to surface water and have identified a secondary lishery but no primary fishery, assign a score of 210. If you do not suspect a release, assign a Secondary Fisheries score from the table below using the lowest flow at any fishery within the target distance limit. Lower flow Secondary Fisheries Score (10 cfs 210 10 to 100 cfs 30 10 to 100 cfs 30 10 to 100 cfs, coastal tidal waters, oceans, 12	PRIMARY FISHERIES: If you suspect any fishery listed above has been exposed to a hazardous substance from the site (see Surface Water Criteria List, page 11), assign a score of 300 and do not evaluate Factor 10. List the primary fisheries: AF River SECONDARY FISHERIES If you suspect a release to surface water and have identified a secondary lishery but no primary fishery, assign a score of 210. If you do not suspect a release, assign a Secondary Fisheries score from the table	
PRIMARY FISHERIES: If you suspect any fishery listed above has been exposed to a hazardous substance from the site (see Surface Water Criteria List, page 11), assign a score of 300 and do not evaluate Factor 10. List the primary fisheries: AF RIVER SECONDARY FISHERIES If you suspect a release to surface water and have identified a secondary lishery but no primary fishery, assign a score of 210. If you do not suspect a release, assign a Secondary Fisheries score from the table below using the lowest flow at any fishery within the target distance limit. Lower Flow Secondary Fisheries Score (10 cfs 210 10 to 100 cfs 30 100 cfs, coastal lidal waters, oceans, 12	PRIMARY FISHERIES: If you suspect any fishery listed above has been exposed to a hazardous substance from the site (see Surface Water Criteria List, page 11), assign a score of 300 and do not evaluate Factor 10. List the primary fisheries: AF River SECONDARY FISHERIES If you suspect a release to surface water and have identified a secondary lishery but no primary fishery, assign a score of 210. If you do not suspect a release, assign a Secondary Fisheries score from the table	
PRIMARY FISHERIES: If you suspect any fishery listed above has been exposed to a hazardous substance from the site (see Surface Water Criteria List, page 11), assign a score of 300 and do not evaluate Factor 10. List the primary fisheries: AF River 300 SECONDARY FISHERIES If you suspect a release to surface water and have identified a secondary lishery but no primary fishery, assign a score of 210. If you do not suspect a release, assign a Secondary Fisheries score from the table below using the lowest flow at any fishery within the target distance limit. Lewest Rew Secondary Reharines Score (10 to 100 cfs 210 10 to 100 cfs 30 10 to 100 cfs 30 10 to 100 cfs, coastal 10 tidal waters, oceans, 12	PRIMARY FISHERIES: If you suspect any fishery listed above has been exposed to a hazardous substance from the site (see Surface Water Criteria List, page 11), assign a score of 300 and do not evaluate Factor 10. List the primary fisheries: AF River 300 D. SECONDARY FISHERIES If you suspect a release to surface water and have identified a secondary fishery but no primary fishery, assign a score of 210. B. If you do not suspect a release, assign a Secondary Fisheries score from the table	Di-
to a hazardous substance from the site (see Surface Water Criteria List, page 11), assign a score of 300 and do not evaluate Factor 10. List the primary fisheries: AFRIVEY D. SECONDARY FISHERIES A. If you suspect a release to surface water and have identified a secondary lishery but no primary fishery, assign a score of 210. B. If you do not suspect a release, assign a Secondary Fisheries score from the table below using the lowest flow at any fishery within the target distance limit. Lewest Flow Secondary Fisheries Score < 10 cfs 210 10 to 100 cfs 30 > 100 cfs, coastal tidal waters, oceans, 12	to a hazardous substance from the site (see Surface Water Criteria List, page 11), assign a score of 300 and do not evaluate Factor 10. List the primary fisheries: AFRIVEY D. SECONDARY FISHERIES A. If you suspect a release to surface water and have identified a secondary fishery but no primary fishery, assign a score of 210. B. If you do not suspect a release, assign a Secondary Fisheries score from the table	
D. SECONDARY FISHERIES A. If you suspect a release to surface water and have identified a secondary fishery but no primary fishery, assign a score of 210. B. If you do not suspect a release, assign a Secondary Fisheries score from the table below using the lowest flow at any fishery within the target distance limit. Lower Flow Secondary Fisheries Score	b. SECONDARY FISHERIES If you suspect a release to surface water and have identified a secondary fishery but no primary fishery, assign a score of 210. If you do not suspect a release, assign a Secondary Fisheries score from the table	
but no primary fishery, assign a score of 210. 3. If you do not suspect a release, assign a Secondary Fisheries score from the table below using the lowest flow at any fishery within the target distance limit. Lowest Flow Secondary Fisheries Score	but no primary fishery, assign a score of 210. 8. If you do not suspect a release, assign a Secondary Fisheries score from the table	•
but no primary fishery, assign a score of 210. 3. If you do not suspect a release, assign a Secondary Fisheries score from the table below using the lowest flow at any fishery within the target distance limit. Lowest Flow Secondary Fisheries Score	but no primary fishery, assign a score of 210. 8. If you do not suspect a release, assign a Secondary Fisheries score from the table	
B. If you do not suspect a release, assign a Secondary Fisheries score from the table below using the lowest flow at any fishery within the target distance limit. Lowest Flow Secondary Fisheries Score	B. If you do not suspect a release, assign a Secondary Fisheries score from the table	
below using the lowest flow at any fishery within the target distance limit. Lewest Flow Secondary Fleherine Score		
Lowert Flow Secondary Fleheres Score	below using the lawaer flow at the figher within the target distance limit	30, - 13
< 10 cfs 210 10 to 100 cfs 30 > 100 cfs, coastal tidal waters, oceans, 12	- DEIOM DRING THE IDMEST HOM ST SHY HENCEY MITHIN THE FRANCE CHEEK CHEEK HINT.	
< 10 cfs 210 10 to 100 cfs 30 > 100 cfs, coastal tidal waters, oceans, 12	Lawrett Share Secondary Schoolst Score	. 1
10 to 100 cfs 30 > 100 cfs, coastal tidal waters, oceans, 12		
> 100 cfs, coastal tidal waters, oceans, 12		
tidal waters, oceans, 12		
	tidal waters, oceans, 12	
000,PM = 8		

SURFACE WATER PATHWAY (continued) ENVIRONMENTAL THREAT SCORESHEET

IKELIHOOD OF RE	LEMSE			Release	Release 100	References
iter Surface Water Lik	elihood of Release sco	re from page 12.	LR =	550	Indiana a las	
NVIRONMENTAL	HREAT TARGETS					
sensitive environme and 5). If there is	ent within the target d	applicable) for each surface wi listance limit (see PA Tables 4 ant within the target distance action of the page.				
Environment Name		Weter Body Type	Rew			
			cfs			
1			cfs			
	Contract Con		cfs			
	C. G. S. Creater		cfs	L. E. Sille		100
				199	were france	
	as been exposed to a	hazardous substance from the	site (see	1 1 1 R		1
ment listed above I Surface Water Criti	as been exposed to a	hazardous substance from the tign a score of 300 and do not	site (see	300		
ment listed above I Surface Water Critifactor 13. List the 3. SECONDARY SENS present, but none it Sensitive Environm A. For secondary s	as been exposed to a inia List, page 11], ass primary sensitive envi TIVE ENVIRONMENT a primary sensitive e- ents based on flow.	hazardous substance from the light a score of 300 and do not ironments: S: If sensitive environments a nvironment, evaluate Seconda on surface water bodies with	site (see evaluate evaluate evaluate	300		
ment listed above I Surface Water Critifactor 13. List the 3. SECONDARY SENS present, but none it Sensitive Environm A. For secondary s	as been exposed to a inia List, page 11], ass primary sensitive envi TIVE ENVIRONMENT a primary sensitive e- ents based on flow.	hazardous substance from the tign a score of 300 and do not fronments: S: If sensitive environments a nvironment, evaluate Seconda	site (see evaluate evaluate evaluate	300		
ment listed above I Surface Water Critifactor 13. List the 3. SECONDARY SENS present, but none is Sensitive Environm A. For secondary s 100 cfs or less, this factor:	as been exposed to a iria List, page 11], ass primary sensitive envi- TIVE ENVIRONMENT a primary sensitive e- ents based on flow. ensitive environments assign scores as follo Odwien Weight	hazardous substance from the light a score of 300 and do not fromments: S: If sensitive environments a nivironment, evaluate Seconda on surface water bodies with twis, and do not evaluate part if the light and t	site (see evaluate ev	300		
ment listed above I Surface Water Critifactor 13. List the Secondary Sens present, but none it Sensitive Environm A. For secondary s 100 cfs or less, this factor:	as been exposed to a iria List, page 11), ass primary sensitive envi- TITVE ENVIRONMENT a primary sensitive e- ents based on flow. ensitive environments assign scores as follo Danden Weight (PA Table 4)	hazardous substance from the tign a score of 300 and do not fronments: S: If sensitive environments a nivironment, evaluate Seconda on surface water bodies with ws, and do not evaluate part I	site (see evaluate ev	300		
ment listed above I Surface Water Criti factor 13. List the SECONDARY SENS present, but none it Sensitive Environm A. For secondary s 100 cfs or less, this factor: Rew cfs	as been exposed to a ria List, page 11), ass primary sensitive environment a primary sensitive ents based on flow. ensitive environments assign scores as follo **Distribution** **Distributi	hazardous substance from the light a score of 300 and do not fromments: S: If sensitive environments a nivironment, evaluate Seconda on surface water bodies with twis, and do not evaluate part if the light and t	site (see evaluate ev	300		
ment listed above I Surface Water Criti factor 13. List the S. SECONDARY SENS present, but none it Sensitive Environm A. For secondary s 100 cfs or less, this factor: Rew	as been exposed to a ria List, page 111, ass primary sensitive environment a primary sensitive ents based on flow. Consitive environments assign scores as follo Division Weight (PA Table 4) .01 x	hazardous substance from the light a score of 300 and do not renments: S: If sensitive environments a nivironment, evaluate Seconda on surface water bodies with wis, and do not evaluate part I	site (see evaluate ev	300		
ment listed above I Surface Water Critifactor 13. List the 3. SECONDARY SENS present, but none is Sensitive Environme A. For secondary s 100 cfs or less, this factor: ###################################	as been exposed to a ria List, page 11), ass primary sensitive environment a primary sensitive ents based on flow. ensitive environments assign scores as follo **Distribution** **Distributi	hazardous substance from the light a score of 300 and do not fromments: S: If sensitive environments a nivironment, evaluate Seconda on surface water bodies with twis, and do not evaluate part if the light Tablee 5 and 6) 400	site (see evaluate ev	300		
ment listed above I Surface Water Critifactor 13. List the 3. SECONDARY SENS present, but none is Sensitive Environm A. For secondary s 100 cfs or leas, this factor: ###################################	as been exposed to a ria List, page 11), ass primary sensitive environment a primary sensitive emits based on flow. ensitive environments assign scores as follo Disclere Weight (PA Table 4) .01 x x	hazardous substance from the light a score of 300 and do not fromments: S: If sensitive environments a nivironment, evaluate Seconda on surface water bodies with twis, and do not evaluate part if the light of the	site (see evaluate ev	300		

304

PA TABLE 5: SURFACE WATER AND AIR PATHWAY SENSITIVE ENVIRONMENTS VALUES

Sensitive Environment	Assigned Value
Critical habitat for Federally designated endangered or threatened species Varine Senctuary Varines Perk Designated Federal Wilderness Area Cologically important areas identified under the Coastal Zone Wilderness Act Sensitive Areas identified under the National Estuary Program or Near Coastal Water Program of the Clean Wate Critical Areas Identified under the Clean Lakes Program of the Clean Water Act (subareas in lakes or entire small Vational Monument (air pathway only) Vational Seashors Recreation Area	100
National Lakeshore Recreation Area	
Isbitst known to be used by Federally designated or proposed endangered or threatened species National Preserve National or State Wildlife Refuge Unit of Coestal Barrier Resources System rederal land designated for the protection of natural ecosysteme Administratively Proposed Federal Wilderness Area Spawning areas critical for the maintenance of fish/shellfish species within a river system, bey, or estuary Nigratory pathways and feeding areas critical for the maintenance of anadromous fish species in a river system Ferrestrial areas utilized for breeding by large or dense aggregations of vertebrate animals (air pathway) or semi-equatio foragers (surface water pathway) Istional river reach designated as Recreational	75
tabitat known to be used by State designated endangered or threatened species (abitat known to be used by a species under review as to its Federal endangered or threatened status (castel Barrier (partially developed) (ederally designated Scenig or Wild River	50
itate lend designated for wildlife or geme management itate designated Scenic or Wild River itate designated Natural Area articular areas, relatively small in size, important to maintenance of unique biotic communities	25
	. 5

PA TABLE 6: SURFACE WATER PATHWAY WETLANDS FRONTAGE VALUES

Total Length of Wetlands	Assigned Value
Less than 0,1 mile	0
0.1 to 1 mile	25
Greeter than 1 to 2 miles	50
Greeter than 2 to 3 miles	75
Greeter than 3 to 4 miles	100
Greater than 4 to 8 miles	150
Greater than 8 to 12 miles	250
Greater than 12 to 16 miles	350
Greater than, 16 to 20 miles	450
Greater than 20 miles	500

SURFACE WATER PATHWAY (concluded) WASTE CHARACTERISTICS, THREAT, AND PATHWAY SCORE SUMMARY

	Α	8
WASTE CHARACTERISTICS	Suspected Release	No Suspected Release
14. A. If you have identified any primary target for surface water (pages 12, 14, or 15), assign the waste characteristics score calculated on page 4, or a score of 32, whichever is GREATER; do not evaluate part 8 of this factor.	100	
B. If you have NOT identified any primary target for surface water, assign the waste characteristics score calculated on page 4.	(100,31, as 10)	(100,32 = 10)
wc -	100	

SURFACE WATER PATHWAY THREAT SCORES

Threat	Likelihood of Release (LR) Score (from page 12)	Targets (T) Score (pages 12, 14, 15)	Pathway Waste Characteristics (WC) Score (determined above)	Threet Score LR x T x WG / 82,500
Drinking Water	550	. 55	100	Indoor to a material of 1000
Human Food Chain	550	300	100	100
Environmental	550	304	100	203

SURFACE WATER PATHWAY SCORE	javageet by a maximum of 1005
(Drinking Water Threat + Human Food Chain Threat + Environmental Threat)	

SUSPECTED CONTAMINATION	RESIDENT POPULATION					
	YNU					
	e o n					
	s k					
	0 10 0	Is any residence, school, or daycare facility of or within 200 feet of an area of suspected contamination?				
Surficial contamination can generally be assumed.	000	Is any residence, school, or daycare facility located on adjacent land previously owned or leased by the site owner/operator?				
	0.00.0	Is there a migration route that might spread				
		hazardous substances near residences, schools, or daycare facilities?				
	0 % 0	Have ensite or adjacent residents or students reported adverse health effects, exclusive of apparent drinking water or air contamination problems?				
	080	Does any neighboring property warrant sampling?				
	o*o	Other criteria?				
	o xo	RESIDENT POPULATION IDENTIFIED?				

SOIL EXPOSURE PATHWAY SCORESHEET .

-	,					
Do any	people live on or within 200 ft of	of areas of suspected cor	ntamination?	Yes	No	
Do ally	people attend school or daycare	on or within 200 ft of a	reas			
of st	uspected contamination?			Yes	No	
is the fa	icility active? Yes No	If yes, estimate the r	number of workers:		_	
			. 1	Suspected	1	
LIKELIHO	OD OF EXPOSURE			Contamination	Reference	м
1. SUSPEC	TED CONTAMINATION: Surfice	ial contamination can ge	nerally be assumed.	(1440 to 14		
	core of 550 assigned. Assign ze			1		
contam	nation can be confidently demoi	nstrated.	LE =	550		
				000	J	-
RESIDENT	POPULATION THREAT TA	RGETS			1	
. RESIDE	NT POPULATION: Determine th	e number of people occu	pying residences			
	ding school or daycare on or wit					
	nation (see Soil Exposure Pathw			0		
			people x 10 =	0	and the same	
				les-d		-
. RESIDE	NT INDIVIDUAL: If you have ide	entified a resident popula	tion (factor 2),	0		
assign a	score of 50; otherwise, assign	a score of O.		0	A STATE OF THE REAL PROPERTY.	000
Management	PC 11			HE HE LE A	100	Spr.
	RS: Use the following table to a				A STATE	
workers	at the facility and nearby facility	ies with suspected conta	amination:		Day 1	
		Soore			120000	
- 1	Number of Workson				The same	
F	Number of Workers					
	0	0				
	0 1 to 100	5				
	0 1 to 100 101 to 1,000	5 10		0		
=	0 1 to 100	5		0		
. TERRES	0 1 to 100 101 to 1,000 >1,000	0 5 10	assign a value	0		
	0 1 to 100 101 to 1,000 >1,000 >TRIAL SENSITIVE ENVIRONMEN	0 5 10 15 NTS: .Use PA Table 7 to	STATE OF THE PARTY	0	-	-
for each	0 1 to 100 101 to 1,000 >1,000 >TRIAL SENSITIVE ENVIRONMEN terrestrial sensitive environmen	0 5 10 15 NTS: .Use PA Table 7 to	STATE OF THE PARTY	0	_	-
for each	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation:	0 5 10 15 NTS: Use PA Table 7 to	d .	0	_	
for each	0 1 to 100 101 to 1,000 >1,000 >TRIAL SENSITIVE ENVIRONMEN terrestrial sensitive environmen	0 5 10 15 NTS: Use PA Table 7 to	STATE OF THE PARTY	0		
for each	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation:	0 5 10 15 NTS: Use PA Table 7 to	d .	0		
for each	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation:	0 5 10 15 NTS: Use PA Table 7 to	d .	Market a		
for each	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation:	0 5 10 15 NTS: Use PA Table 7 to	Value	0 25		
for each	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation:	0 5 10 15 NTS: Use PA Table 7 to	d .	Market a		-
for each contami	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN terrestrial sensitive environmen nation: Terrestrial Sensitive Environment Ty	0 5 10 15 NTS: Use PA Table 7 to	Value	25		
for each	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN terrestrial sensitive environmen nation: Terrestrial Sensitive Environment Ty	0 5 10 15 NTS: Use PA Table 7 to	Value	25		
for each contami	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN terrestrial sensitive environmen nation: Terrestrial Sensitive Environment Ty	0 5 10 15 NTS: Use PA Table 7 to	Value Sure =	25		
for each contami	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN terrestrial sensitive environmen nation: Terrestrial Sensitive Environment Ty	0 5 10 15 NTS: Use PA Table 7 to	Value	25 5		
for each contami	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN terrestrial sensitive environmen nation: Terrestrial Sensitive Environment Ty	0 5 10 15 NTS: Use PA Table 7 to	Value Sure =	25 5		
for each contami	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN terrestrial sensitive environmen nation: Terrestrial Sensitive Environment Ty	0 5 10 15 NTS: Use PA Table 7 to it on an area of suspecte	Value Sure =	25 5 30		
on each contami	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN terrestrial sensitive environmen nation: Ferrestrial Sensitive Environment Ty	0 5 10 15 NTS: Use PA Table 7 to it on an area of suspecte	Value Sure =	25 5 30		
ontami	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN terrestrial sensitive environmen nation: Ferrestrial Sensitive Environment Ty	0 5 10 15 NTS: Use PA Table 7 to it on an area of suspecte	Value Sure =	25 5 30		
on each contami	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation: Terrestrial Sensitive Environment Ty ACES HARACTERISTICS the waste characteristics score of	0 5 10 15 15 15 15 15 15 15 16 16 16 16 16 16 16 16 16 16 16 16 16	Value Sure = WC =	25 5 30		
on each contami	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN terrestrial sensitive environmen nation: Ferrestrial Sensitive Environment Ty	0 5 10 15 15 15 15 15 15 15 16 16 16 16 16 16 16 16 16 16 16 16 16	Value Sure =	25 5 30		
on each contami	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation: Terrestrial Sensitive Environment Ty ACES HARACTERISTICS the waste characteristics score of	O 5 10 15 NTS: Use PA Table 7 to at on an area of suspecte calculated on page 4.	Value Sure = WC =	25 5 30		
on each contami	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation: Terrestrial Sensitive Environment Ty ACES HARACTERISTICS the waste characteristics score of	O 5 10 15 NTS: Use PA Table 7 to at on an area of suspecte calculated on page 4.	Value = T = WC =	25 5 30		
NASTE C	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation: Ferrestrial Sensitive Environment Ty RCES THARACTERISTICS the waste characteristics score in	O 5 10 15 VTS: Use PA Table 7 to at on an area of suspecte	Value = T = WC =	25 5 30 100 21 = 19 100		
NASTE C	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation: Terrestrial Sensitive Environment Ty ACES HARACTERISTICS the waste characteristics score of	O 5 10 15 VTS: Use PA Table 7 to at on an area of suspecte	Value = T = WC =	25 5 30		
NASTE C	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation: Ferrestrial Sensitive Environment Ty RCES THARACTERISTICS the waste characteristics score in	O 5 10 15 VTS: Use PA Table 7 to at on an area of suspecte	Value = T = WC =	25 5 30 100 21 = 19 100		
NASTE C	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation: Ferrestrial Sensitive Environment Ty RCES THARACTERISTICS the waste characteristics score in	O 5 10 15 VTS: Use PA Table 7 to at on an area of suspecte	Value = T = WC =	25 30 100 xx = 10 100 20 1 ^{10, x = 10}		
WASTE C	O 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMENT of terrestrial sensitive environment on terrestrial Sensitive Environment Type RCES HARACTERISTICS the waste characteristics score of POPULATION THREAT SCO	O 5 10 15 NTS: Use PA Table 7 to at on an area of suspecte calculated on page 4.	Value = T = WC =	25 5 30 1000 20 0 100 1000 20 1 ^(6, 2, 4, 6)		
NASTE C	0 1 to 100 101 to 1,000 >1,000 >1,000 TRIAL SENSITIVE ENVIRONMEN I terrestrial sensitive environmen nation: Ferrestrial Sensitive Environment Ty RCES THARACTERISTICS the waste characteristics score in	O 5 10 15 VTS: Use PA Table 7 to at on an area of suspecte calculated on page 4.	Value = T = WC =	25 30 100 xx = 10 100 20 1 ^{10, x = 10}		

PA TABLE 7: SOIL EXPOSURE PATHWAY TERRESTRIAL SENSITIVE ENVIRONMENT VALUES

Terrestrial Sensitive Environment	Assigned Value
Terrestrial critical habitat for Federally designated endangered or threatened species	100
National Park	
Designated Federal Wilderness Area	
National Monument	
Terrestrial habitat known to be used by Federally designated or proposed threatened or endangered species	75
National Preserve (terrestrial)	
National or State terrestrial Wildlife Refuge	
Federal land designated for protection of natural ecosystems	
Administratively proposed Federal Wilderness Area	
Terrestrial areas utilized by large or dense aggregations of animals (vertebrate species) for breeding	The state of the s
Terrestrial habitat used by State designated endangered or threatened species	50
Terrestrial habitat used by species under review for Federal designated endangered or threatened status	
State lands designated for wildlife or game management	25
State designated Natural Areas	
Particular areas, relatively small in size, important to maintenance of unique biotic communities	

		SUSPECTED RELEASE	PRIMARY TARGETS
	-		
YN	U n .		
5	k		
O Ø		Are odors currently reported?	
пп	п	Has release of a hazardous substance to the air	
	-	been directly observed?	If you suspect a release to air, evaluate all populations and
- 4			sensitive environments within 1/4 mile (including those
0 0	U	Are there reports of adverse health effects (e.g., headaches, neusea, dizziness) potentially	onsite) as primary targets.
		resulting from migration of hazardous	
		substances through the air?	
- X	_		
ם ט	П	Does analytical or circumstantial evidence suggest a release to the air?	
		2000 at a lateral to the art	
00		Other criteria?	
60		SUSPECTED RELEASE?	
si	tes	. Dust from tailings will trave	ring periods of wind at all three el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.
si	tes	. Dust from tailings will trave	el through means of the air pathway.

AIR PATHWAY SCORESHEET

Pothway Characteristics		
Do you suspect a release (see Air Pathway Criteria List, page 21)? Distance to the nearest individual:	Yes	No
	A Suspensed	No Supposted
LIKELIHOOD OF RELEASE	Releases	Release
 SUSPECTED RELEASE: If you suspect a release to air (see page 21), assign a score of 550. Use only column A for this pathway. 	550	4609
NO SUSPECTED RELEASE: If you do not suspect a release to air, assign a score of 500. Use only column 8 for this pathway.		
LR =	550	
PRIMARY TARGET POPULATION: Determine the number of people subject to exposure from a suspected release of hazardous substances to the air.	. 0	
SECONDARY TARGET POPULATION: Determine the number of people not suspected to be exposed to a release to air, and assign the total population score using PA Table 8.		
NEAREST INDIVIDUAL: If you have identified any Primary Target Population for the air pathway, assign a score of 50; otherwise, assign the Nearest Individual score from PA Table 8.	20	(31/11.00
PRIMARY SENSITIVE ENVIRONMENTS: Sum the sensitive environment values (PA Table 5) and wetland acreage values (PA Table 9) for environments subject to exposure from a suspected release to the air.		
Samulton Employment Type Value Value	0	
Succ = SECONDARY SENSITIVE ENVIRONMENTS: Use PA Table 10 to determine the score for secondary sensitive environments.		N. Best
RESOURCES	5	
T =	25	
A. If you have identified any Primary Target for the air pathway, assign the waste characteristics score calculated on page 4, or a score of 32, whichever is GREATER; do not evaluate part 8 of this factor.	100	
 If you have NOT identified any Primary Target for the air pathway, assign the weste characteristics score calculated on page 4. 	(10L3), o 10	110E.25. ar 180
wc -		0.005
AIR PATHWAY SCORE: LR x T x WC 82 500	17	

PA TABLE 8: VALUES FOR SECONDARY AIR TARGET POPULATIONS

	Mine Story	Nearest	Population Within Distance Category												
Distance from Site	Population	Individual (choose highest)	1 te 10	11 to 30	31 to 100	101 to 300	301 fe 1,000	1,001 te 3,000	2,001 te 10,000	10,001 te 30,000	30,001 Le 100,000	100,001 te 300,000	300,001 fe 1,000,000	Greater EALS 1,000,000	Population Value
Oneite		20	1	2	6	10	52	163	521	1,633	6,214	16,326	62,136	163,246	
>0 to ¼ mile	-	20	1	1	1	4	13	41	130	408	1,303	4,081	13,034	40,811	,
> X to X mile		2	0	0	1.	1	3	9	28	88	282	882	2,815	8,815	
> % to 1 mile		1	0	0	0	1	. 1	3		26	83	261	834	2,612	
1 to 2 miles	-	0	0	0	0	0	1	1	3	8	27	83	266	833	-
> 2 to 3 miles	-	0	0	0	0	0	1	1 1	1_	4	12	38	120	376	
>3 to 4 miles		. 0	0	0	0	0	. 0	1	1	2	7	23	73	229	

PA TABLE 9: AIR PATHWAY VALUES FOR WETLAND AREA

Wetland Area	Assigned Value
Less than 1 acre	0
1 to 50 acres .	25
Greater then 50 to 100 acres	76
Greater than 100 to 150 scree	125
Greater than 150 to 200 acres	176
Greater than 200 to 300 acres	250
Greater than 300 to 400 acres	350
Greater than 400 to 500 acres	450
Greater than 500 acres	500

PA TABLE 10: DISTANCE WEIGHTS AND CALCULATIONS FOR AIR PATHWAY SECONDARY SENSITIVE ENVIRONMENTS

Distance	Distance Weight	Sensitive Environment Type and Value (from PA Table 5 or 9)	Product
Onsite	0.10	×	
		x -	
0-1/4 mi	0.025	x	
		*	
		*	
1/4-1/2mi	0.0054	* 11-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1	
		* The second sec	
		*	
	The state of	x	

Total Environments Score =

SITE SCORE CALCULATION

	s	S ²
GROUND WATER PATHWAY SCORE (Som):	2.06	4.24
URFACE WATER PATHWAY CORE (S,):	100	10,000
OIL EXPOSURE PATHWAY CORE (S,):	21	441
IR PATHWAY CORE (S.):	17	289
TE SCORE:	$\frac{S_{gw^2} + S_{gw^2} + S_{g^2} + S_{g^2}}{4}$	51.8

SUMMARY

	YES	NO
Is there a high possibility of a threat to any nearby drinking water well(s) by migration of a hazardous substance in ground water? A. If yes, identify the well(s).	0	*
B. If yes, how many people are served by the threatened well(s)? Is there a high possibility of a threat to any of the following by hezardous substance.		
migration in surface water?		
A. Drinking water intake 8. Fishery C. Sensitive environment (wetland, critical habitat, others) D. If yes, identify the target(s).	MAG	X00
Is there a high possibility of an area of surficial contamination within 200 feet of any residence, school, or daycare facility?	п	×
If yee, identify the property(lee) and estimate the associated population(s).		
Are there public health concerns at this site that are not addressed by PA scoring considerations? If yes, explain:	*	0
	A. If yes, identify the well(s). B. If yes, how many people are served by the threatened well(s)? Is there a high possibility of a threat to any of the following by hazardous substance migration in surface water? A. Drinking water intake B. Fishery C. Sensitive environment (wetland, critical habitat, others) D. If yes, identify the target(s). Is there a high possibility of an area of surficial contamination within 200 feet of any residence, school, or daycare facility? If yes, identify the property(lee) and estimate the associated population(s). Are there public health concerns at this site that are not addressed by PA scoring considerations? If yes, explain:	Is there a high possibility of a threat to any nearby drinking water well(s) by migration of a hazardous substance in ground water? A. If yes, identify the well(s). B. If yes, how many people are served by the threatened well(s)? Is there a high possibility of a threat to any of the following by hazardous substance migration in surface water? A. Drinking water intake B. Fishery C. Sensitive environment (wetland, critical habitat, others) D. If yes, identify the target(s). Is there a high possibility of an area of surficial contamination within 200 feet of any residence, school, or daycare facility? If yee, identify the property(lee) and estimate the associated population(s). Are there public health concerns at this site that are not addressed by PA scoring considerations? If yes, explain:

PART III

Correspondence (in cronological order)

Forest Service

R-4

2840 Reclamation

... AUG 7 1985

Subject:

Functional Assistance - Pacific Mine

Forest Supervisor, Uinta NF

Enclosed is a short summary and photograph report from Ben Albrechtsen's recent trip to the Uinta National Forest. We hope you find these comments helpful. Please transmit the original to Harry Opfar for his field use.

If you have comments or questions, please contact Ben directly.

E. R. BROWNING

Director

Minerals Area Management

Enclosure

Pleasa t Cri OCT 1	
Fi	# 9
In:	
BUSAIGT	
REC, 1	
ALL	

Jes Jen

. 152 . 152 . 1 to 2

cg D-2

2



Reclamation Recommendations for Pacific Mine Pleasant Grove Ranger District Uinta National Forest

> by Ben Albrechtsen R-4 Reclamation Specialist July 1985

Summary

On July 22, I spent the day on the Pleasant Grove Ranger District reviewing the Pacific Mine property. The purpose of my trip was to make suggestions for reclamation of this area. Accompanying me were:

Harry Ophar - Ranger Ralph McDonald - Forestry Technician

The Pacific Mine is an old gold and silver mine that is still held by active claims. The present claimant is Dan Proctor of Pleasant' Grove, Utah.

Mr. Proctor is interested in spending several thousand dollares in the near future to drill the property, define the ore body, and reopen the mine.

Additionally, he has verbally agreed to provide some reclamation on the previously disturbed site and old tailings pond.

The mine is located in the upper reaches of American Fork Canyon approximately 10 miles above Timpanogus Cave National Monument. American Fork River heads above the mine and is being polluted from water coming off the mine dump and out of the old portal. Recent high water and current beaver activity are worsening this situation, causing further pollution to downstream recreation, irrigation, and fisheries values.

I feel every effort should be made to correct this situation as soon as possible. If the Ranger can negotiate some reclamation through an operating plan, that would be excellent. If this cannot be accomplished quickly, then the Utah State Department of Water Resources should review the situation and corrective action be taken. Part of the property is on private land and the Ranger should get the Utah Division of Oil, Gas and Mining to review the problem. It appears this property would be high on their list for reclamation action.

The following are immediate remedial actions that need to be taken at the Pacific Mine:


- Divert the portal drainage away from the tailings area and into its original channel and then to the main American Fork Creek.
- Obtain both water and soil samples to see if acid mine drainage exists and what contaminants are in the tailings. Send the soil to A&L Laboratory in Omaha, Nebraska. Mention that samples are mine spoils and a complete analysis is needed.
- Close the mine area to off-road vehicle travel to reduce disturbance to the site. This will reduce sediment into the stream until reclamation can be accomplished.
- 4. If suspected acid contaminants are present, call for reclamation assistance from the Intermountain Station in Logan.

5. Determine waste hazards and reclamation objectives before drilling or mining the site again.


The following photograph report will illustrate specific ideas discussed on the ground with Ranger Ophar and Ralph McDonald.

Ben Albrechtsen

Reclamation Specialist




No. 1 The Pacific Mine is a gold and silver mine situated in the upper reaches of the American Fork River, Pleasant Grove RD, Uinta NF. The claimant, Dan Proctor of Pleasant Grove, Utah, would like to reopen the mine after doing some exploratory drillings. Mr. Proctor is willing to do reclamation on the existing disturbance.




No. 2 The water in American Fork Creek above the Pacific Mine appears clean and free of contaminants (photo taken immediately upstream from the mine on July 22, 1985).




No. 3 American Fork Creek, immediately below the mine, is polluted with sediment and possibly acid mine drainage from the portal. Fish populations are limited or nonexistent until the stream is diluted by other tributaries somewhere near Tibble Fork Reservoir downstream.



No. 4 Sediment and contaminants are readily visible downstream from the mine (% mile).




No. 5 The major source of contamination to the stream is this water coming from the old mine portal site. The brown stain on the rocks (yellow boy) indicates the presence of acid mine drainage.




No. 6 Erosion has diverted the water from the vegetated channel on the right of the dump, through the dump (arrows) and into the basin in the foreground. Beaver activity in the basin has ponded the water and diverted it through the mine tailings, polluting American Fork Creek more than would occur normally.



No. 7 The drainage from the portal at Pacific Mine is running through the old tailing dissolving substantial amounts of contaminants and depositing them directly into the stream (see photo No. 8).




No. 8 This situation should b corrected immediately and steps taken to ensure it cannot happen in the future.




No. 9 Immediate relief of most of this problem can be obtained by trapping and removing the beaver and turning the portal drainage out of the tailings and into the established channel shown below.



No. 10 Beaver activity has diverted portal water out of this channel and onto the tailings area. (Note presence of yellow boy on rocks.)




No. 11 The entire mine and tailings area should be reshaped to handle water differently, then be treated and seeded to establish a vegetative cover on it. Vegetation will reduce oxidation as well as control sediment movement. Several species of grass, juncus, sedge, and willow seem readily adaptable to these conditions. Successful revegetation seems likely. Expertise from the Forest Service Research Station in Logan is recommended to determine proper soil ammendments, etc.



No. 12 Slopes of the dump and tailings are immediately adjacent to the creek and need to be stabilized.



No. 13 Off-road vehicle traffic on the tailings is loosening surface material and adding sediment to American Fork Creek. This area should be closed to off-road vehicle use immediately.



No. 14 Miller Hill, another mine in American Fork Canyon, may be contributing similar problems to the Pacific. This property belongs to the Forest Service. It should be examined and future use or treatment of it should be determined.



No. 15 · There are several abandoned and active properties in American Fork Canyon that are possibly contributing to water quality degradation. These should be examined by the Forest Service and/or the State to determine their status.



No. 16 American Fork Canyon and its tributaries have high scenic and recreational values and mine operating plans should reflect future desired conditions after mining.

2840 Reclamation

Date: AUG 7 1985

Subject

Functional Assistance - Pacific Mine

Forest Supervisor, Uinta NF

Enclosed is a short summary and photograph report from Ben Albrechtsen's recent trip to the Uinta National Forest. We hope you find these comments helpful. Please transmit the original to Harry Opfar for his field use. $d \in \mathcal{A} = \{0 - 1 - 1 - 1 \}$

If you have comments or questions, please contact Ben directly.

E. R. BROWNING

Director

Minerals Area Management

Enclosure

se of outer taken.

10:85

cg D-2

20/07/24 C



Reclamation Recommendations for Pacific Mine Pleasant Grove Ranger District Uinta National Forest

> by Ben Albrechtsen R-4 Reclamation Specialist July 1985

Summary

On July 22, I spent the day on the Pleasant Grove Ranger District reviewing the Pacific Mine property. The purpose of my trip was to make suggestions for reclamation of this area. Accompanying me were:

Harry Ophar - Ranger Ralph McDonald - Forestry Technician

The Pacific Mine is an old gold and silver mine that is still held by active claims. The present claimant is Dan Proctor of Pleasant Grove, Utah. Mr. Proctor is interested in spending several thousand dollares in the near future to drill the property, define the ore body, and reopen the mine. Additionally, he has verbally agreed to provide some reclamation on the previously disturbed site and old tailings pond.

The mine is located in the upper reaches of American Fork Canyon approximately 10 miles above Timpanogus Cave National Monument. 'American Fork River heads above the mine and is being polluted from water coming off the mine dump and out of the old portal. Recent high water and current beaver activity are worsening this situation, causing further pollution to downstream recreation, irrigation, and fisheries values.

I feel every effort should be made to correct this situation as soon as possible. If the Ranger can negotiate some reclamation through an operating plan, that would be excellent. If this cannot be accomplished quickly, then the Utah State Department of Water Resources should review the situation and corrective action be taken. Part of the property is on private land and the Ranger should get the Utah Division of Oil, Gas and Mining to review the problem. It appears this property would be high on their list for reclamation action.

The following are immediate remedial actions that need to be taken at the Pacific Mine:


- 1. Divert the portal drainage away from the tailings area and into its original channel and then to the main American Fork Creek.
- 2. Obtain both water and soil samples to see if acid mine drainage exists and what contaminants are in the tailings. Send the soil to A&L Laboratory in Omaha, Nebraska. Mention that samples are mine spoils and a complete analysis is needed.
 - 3. Close the mine area to off-road vehicle travel to reduce disturbance to the site. This will reduce sediment into the stream until reclamation can be accomplished.
- 4. If suspected acid contaminants are present, call for reclamation assistance from the Intermountain Station in Logan.

Ded of the


5. Determine waste hazards and reclamation objectives before drilling or mining the site again.

The following photograph report will illustrate specific ideas discussed on the ground with Ranger Ophar and Ralph McDonald.

Ben Albrechtsen Reclamation Specialist




No. 1 The Pacific Mine is a gold and silver mine situated in the upper reaches of the American Fork River, Pleasant Grove RD, Uinta NF. The claimant, Dan Proctor of Pleasant Grove, Utah, would like to reopen the mine after doing some exploratory drillings. Mr. Proctor is willing to do reclamation on the existing disturbance.



No. 2 The water in American Fork Creek above the Pacific Mine appears clean and free of contaminants (photo taken immediately upstream from the mine on July 22, 1985).




No. 3 American Fork Creek, immediately below the mine, is polluted with sediment and possibly acid mine drainage from the portal. Fish populations are limited or nonexistent until the stream is diluted by other tributaries somewhere near Tibble Fork Reservoir downstream.



No. 4 Sediment and contaminants are readily visible downstream from the mine (½ mile).




No. 5 The major source of contamination to the stream is this water coming from the old mine portal site. The brown stain on the rocks (yellow boy) indicates the presence of acid mine drainage.




No. 6 Erosion has diverted the water from the vegetated channel on the right of the dump, through the dump (arrows) and into the basin in the foreground. Beaver activity in the basin has ponded the water and diverted it through the mine tailings, polluting American Fork Creek more than would occur normally.



No. 7 The drainage from the portal at Pacific Mine is running through the old tailing dissolving substantial amounts of contaminants and depositing them directly into the stream (see photo No. 8).



No. 8 This situation should be corrected immediately and steps taken to ensure it cannot happen in the future.




No. 9 Immediate relief of most of this problem can be obtained by trapping and removing the beaver and turning the portal drainage out of the tailings and into the established channel shown below.



No. 10 Beaver activity has diverted portal water out of this channel and onto the tailings area. (Note presence of yellow boy on rocks.)




No. 11 The entire mine and tailings area should be reshaped to handle water differently, then be treated and seeded to establish a vegetative cover on it. Vegetation will reduce oxidation as well as control sediment movement. Several species of grass, juncus, sedge, and willow seem readily adaptable to these conditions. Successful revegetation seems likely. Expertise from the Forest Service Research Station in Logan is recommended to determine proper soil ammendments, etc.




No. 12 Slopes of the dump and tailings are immediately adjacent to the creek and need to be stabilizer



No. 13 Off-road vehicle traffic on the tailings is loosening surface material and adding sediment to American Fork Creek. This area should be closed to off-road vehicle use immediately.



No. 14 Miller Hill, another mine in American Fork Canyon, may be contributing similar problems to the Pacific. This property belongs to the Forest Service. It should be examined and future use or treatment of it should be determined.



No. 15 There are several abandoned and active properties in American Fork Canyon that are possibly contributing to water quality degradation. These should be examined by the Forest Service and/or the State to determine their status.



No. 16 American Fork Canyon and its tributaries have high scenic and recreational values and mine operating plans should reflect future desired conditions after mining.

OWNA WE

JE17 '89

4300/2840

Subjec

Old Mining Disturbances - American Fork Canyon

Forest Supervisor, Uinta NF

A Lean Roy Mine, and the M

On May 30, 1989, I visited the Pacific Mine, the Lower Bog Mine, and the Mary Ellen Gulch Mines in American Fork Canyon. I was with Paul Skabelund of the SO and two employees of the State of Utah. All of these mines have been inactive for a long time.

The existing water effluent from these mines is not good quality water. Paul Skabelund's data from BYU indicated that these mine effluents are high in one or more of lead, cadmium, or zinc. On the other hand, it is important to note that there are fish in the river only a short distance downstream from any of these mines. The benthic organisms in the streams may suffer a decline during the late summer season.

The Lower Bog Mine is apparently entirely on the Uinta National Forest, but both the Pacific and Mary Ellen Gulch Mines have a mixed federal-private ownership. On these two mines, less than half of the effected area is on National Forest land. This mixed ownership situation probably means that the State of Utah would have to be an active partner if a significant cleanup effort is to be made. The State may have to put legal pressure on the private landowners to spend money to clean up their land. In my view, that is extremely unlikely. Even though these mine water discharges are in violation of State water quality standards, there is little likelihood that the State will act to bring these waters back into compliance with the standards. This is due to the facts that the mine discharges predate the water quality standards law; the mine discharges are in compliance with the State non-degradation clause; and the existing mine discharge is just not all that bad.

The adit flow from the Lower Bog Mine could be collected, piped a short distance, and run through an artifically constructed wetlands. Properly done this treatment could result in improved water quality flowing from the adit to the American Fork River. However, this would be expensive due to the steep rocky terrain, the need to build individual wetland cells, and the need to make significant improvements to the existing road. I estimate the costs at about \$40 to \$75 thousand.

From a technical viewpoint, both the Pacific and Mary Ellen Gulch Mines can be cleaned up and stabilized. It would be expensive, \$40 to \$65 thousand on the Pacific and \$1 to \$2 million on the Mary Ellen Gulch.

These costs make it doubtful that the potential improvement in water quality is worth the cost. However, I do believe that we can affect some material increase in water quality at a relatively low cost by taking steps to keep adit flows or other surface waters from flowing over or through tailings and waste rock piles. This could be done by gathering up these surface waters and



putting them into concrete ditches or plastic pipes and taking them directly to the river. This would prevent further deterioration of surface water quality and would be a useful step at a reasonable cost—a few thousands of dollars at the Pacific Mine and perhaps \$25 thousand at the Mary Ellen Mines. I would suggest that you might consider this action as a fishery improvement project. A hindrance to this mitigation is the mixed ownership; we do not control the lands around these adits.

Every year all of these sites are loosing significant amounts of tailings and waste rock dust through wind erosion.

CONCLUSION AND RECOMMENDATION

We are in a poor position to initiate any cleanup action at the Pacific Mine or the Mary Ellen Gulch Mines because of the ownership situation and the relatively innocuous nature of the problem. On the other hand, it would be useful and relatively cheap to pipe the adit water across the contaminated tailings and waste rock piles. Perhaps that could be done to improve the fisheries.

I recommend that you take no action at the Lower Bog Mine since this mine is a very minor contributor to the overall water quality problems in the North Fork of the American Fork River.

YEUGENE E. FARMER

West-Wide Reclamation

Sen aller to

Specialist

cc:

RW - Stender

MAM - Farmer

Reply to: 2800

Date: March 15, 198

Subject: Minerals-Related Nonpoint Pollution Abatement Needs--Federal Facilities Compliance Program (Your 1tr. 1/28)

To: Regional Forester

We discussed our needs with Gene Former and determined that the Uinta will provide updated wo. tubests for mining related projets at this time. Uinta received \$28,000 of Clean Water Act funding this fiscal year to st the potential problems at mining related sites. With these forms, we plandetermine the water quality impacts at the following:

- 1. Pacific Hine
- 2. Mary Ellan Gulch Mining Area

- 3. Miller Hill Mine
 4. Bog Mine
 5. North Oakbrush Mine
 6. Harker Mine

- 7. Bingelli Mine S. Red Ledges Mine
- 9. Lost Josephine Mine

These studies will also address whether corrective action is needed, what anything should be done to correct the situation, and cost estimates taking corrective action.

/s/ Don T. Nebeker

DON T. NEBEKER Forest Supervisor

CC: U.Stokes J.Reese D-2 D-3



Norman H. Bangerter Governor Dee C. Hansen Executive Director Dianne R. Nielson, Ph.D.

State of Utah

DEPARTMENT OF NATURAL RESOURCES DIVISION OF OIL, GAS AND MINING

355 West North Temple 3 Triad Center, Suite 350 Sall Lake City, Utah 84180-1203 801-538-5340

August 7, 1990

Mr. Paul Skablund Uinta National Forest 100 North 88 West Provo, Utah 84601

Dear Mr. Skablund:

The Utah Abandoned Mine Reclamation Program is concerned with mitigating physical hazards to the public health and safety that occur on abandoned mine sites. We would be able to provide you with plans and specifications for typical closure techniques to secure abandoned mine portals and other work specifications for earthwork, demolition, etc. Possibly a staff member could discuss with you how to draw up reclamation plans for particular sites.

If health hazards are present due to toxic substances, the Utah Department of Health, Environmental Health section, usually takes responsibility.

Possible solutions to the Pacific and Mary Ellen Gulch mines, where extensive tailings dumps are present would be to: 1) route runoff around the dumps and try to stabilize the dumps in place, or 2) remove the dump materials to a lower precipitation site. Removing the dumps would be expensive, logistically difficult and could aggravate the problems present by introducing oxygen into the system. Off-road vehicle use should be prevented at the Pacific Mine tailings area.

From the information in Dr. Merritt's report, elevated levels of cadmium, copper, lead and zinc are present but confined somewhat to localized areas within a mile of the discharge point. Methods to lower these levels are generally prohibitively expensive. It does appear that some of the parameters sampled, particularly copper, lead and zinc increase substantially after flowing through the dump material. Thus, I would recommend preventing, as much as possible, all runoff from flowing over or through the dump.

Please call me if you would like to discuss this further. I would appreciate it if you would keep me informed about the progress of this project.

Sincerely,

Lucia Malin'

Senior Reclamation Specialist

Lucia) mali

an equal opportunity employer AM806/186



Norman H. Bangerter Governor Suzanne Dandoy, M.D., M.P.H. Executive Director Kenneth L. Alkerna

State of Utah DEPARTMENT OF HEALTH DIVISION OF ENVIRONMENTAL HEALTH

288 North 1460 West P.O. Box 16690 Salt Lake City, Utah 84116-0690 (801) 538-6121

August 8, 1990

Mr. Paul H. Skabelund Uinta National Forest 88 North 100 West Provo, Ut 84603

'RE: American Fork River Water Quality

Dear Mr. Skabelund:

This letter is in response to your request for assistance in planning for possible mitigation of water quality problems in the North Fork drainage of the American Fork River. Water quality impairments associated with mine water drainages and mine tailings have been identified on patented and U.S. Forest Service Lands.

Data have shown that impacts are generally localized around water flowing from adits and across tailings piles and cause elevated concentrations of some heavy metals including cadmium, zinc, and lead. Unstabilized tailings piles appear to be the primary cause as metals are dissolved and carried downstream. Substantial dilution at the main stem of the North Fork reduces concentrations and thus reduces the impacts. However, slight impairments have been demonstrated in macroinvertebrate populations and water chemistry samples.

Our primary concern in this drainage is the protection of water quality and public health along the main stem of North Fork which is a high public use system. As mitigation is planned, it would seem appropriate to stabilize tailings piles adjacent to the stream which are subject to runoff and erosion. Secondary measures could include the diversion of mine drainage away from the tailings, and the construction of wetlands to reduce sediment and metals loads to receiving streams.

Attempts should be made during the planning phase to identify and contact patent holders within the drainage system and determine if cooperative agreements can be established. Without such contacts, any attempts at mitigation may only partially succeed since mine flow often originates on these private land holdings.

We would like to remain involved in your planning process as well as assist, as resources allow, in additional data collection and analysis including water quality parameters, fish tissue analysis, and macroinvertebrate community studies. Please contact Reed Oberndorfer of my staff at 538-6146 for additional information.

Sincerely,

Don A. Ostler, P.E., Director

In a. Osten

Bureau of Water Pollution Control

RYO:pb skabblund.ttr

UINTA NATIONAL FOREST

JUL 9 1991 RIGHT OF ENTRY CONSENT TO THE PACIFIC MINE SITE

by

Euro-Nevada Mining Company W. Dan Proctor-Agent

ACTION FS SEC	ACTION
_ AO	-B&F-
E/MN	_ PERS_
FM	_ PIO
R&L_	DIST D-2
_RWL	/ Paul 98-15
I IVI	

I, the undersigned, W. Dan Proctor as agent for the Euro-Nevada Mining Company, do hereby consent to the study and determination of hazards to the publics health, safety and general welfare at the site known as the Pacific Mine by the Utah Division of Oil, Gas & Mining, Department of Natural Resources (Division) and its agents, employees or contractors.

The Pacific Mine site is particularly described as in the:

SE4 of Section 22 (unsurveyed), Township 3 South, Range 3 East, SLBM.

The Division expressly assumes liability for any and all injuries sustained by its employees. Furthermore, the Division expressly waives liability of the Landowner for any and all injuries sustained by Division employees.

Except as herein set forth in this right of entry consent, neither the Division nor Landowner shall undertake any activity, either expressed or implied, nor make any representation which purports to bind the other.

It is expressly understood that all costs incurred for studies and tests shall be the sole liability of the Division.

This consent will expire December 31, 1993.

Dated this 27th

day of

,199/

DIVISION OF OIL, GAS & MINING

() ()

Name: Dianne R. Nielson

Title: Director

Euro-Nevada Mining Co.

Name: W. Dan Proctor

Title: Landowner/Agent

Mary Ann Wright, Administrator Abandoned Mine Reclamat, on Program

Euro-Nevada Mining Corporation, Inc.

6121 Lakeside Drive, Suite 240 Reno, Nevada 89511 (702) 825-8890 / Fax (702) 825-4994

October 4, 1991

Mr. Robert R. Easton District Ranger United States Forest Service Pleasant Grove Ranger District P. O. Box 228 Pleasant Grove, Utah 84062

Reclamation Measures in the Vicinity of the Pacific Mine

Dear Mr. Easton:

Thank you for your letter of September 24, 1991 regarding the rehabilitation of certain matters in the vicinity of the Pacific Mine. We have discussed your letter with Mr. W. Dan Proctor, and have this date authorized him to undertake the measures requested in your letter. We have asked Mr. Proctor to complete this work before winter sets into the area and to provide us with a report upon completion. We will provide you with a copy of the report and, at that time, request that you inspect the property and provide us with your acceptance of the work. Of course, should you have any questions or additional concerns, please contact us at your convenience. We thank you for keeping us informed.

Very truly yours,

M. Craig Haase **Executive Vice President**

Mr. W. Dan Proctor

CC:

Mr. Pierre Lassonde

RECEIVED Pleasant Grove Flanger District OCT 0 7 1991 RESASG RECREATION SECTION

Paul -Here are my comments in the vater quility and North fork of the American fork Nich First a couple of general comments. its al-years difficult to make I contifusive statements about water quality when there are only two or linee samples. This is especially live I in that of here larguers comments about here; in being flushed out of the system at the first of the snowmell period. Secondly, the forest needs to delive where it wants to go with this study. for the F.S. to undertake some rehald projects (eg Pacific Mine tailings statistication). On the other hand if you want to prove that patented lands are polleting NFS strowns youll need a lot more intersive studies including more chancial analyses, macroinvertebrate sumpling and perhaps even fish bioassays. I personally inite well Lest leave water quality violetions on private lands up to the water Pollu tion Control Bureau My Mind Comment has to do with the way I losted at deta only in terms of impacts to beneficial water uses. In these avainages cold water Isharies is the only recognized beneficial use Thus, small tributary streams without fish are not polluted even if there are extremely bight levels of toxic chemicals.

Specific Comments

Fine from lower bog mine is violating standards in No Fk American Fk River assuming there are fish in this section of the River. It I seems that I recall you saying there is a waterfall that acts as a barrier to fish downstream. If this is the case and there's no fisheries here, then, by definition, there's no pollution.

Lead and Zinc appear to be in high concentrations in the effluent from Pacific Mine (portals & runoff from tailings). However, data from site #8 (No Fk American Fix R below Pacific Mino) does not clearly indicate standards violation with this data, I believe you'd have a hard time convincing anyone that lead and zinc from Pacific Mine are impucting beneficial use

Zinc from Mary Ellen Mine shows a high concentration in Mary Ellen Bulch immediately below the mines. However, by the time it gets to the mouth of Mary Ellen Gulch its diluted and meets standards. The question is how much of Mary Ellen Cycloh contains a fishery. That will determine whether there's pollution is the strict sense.

Jummary I don't see a great deal of impact on beneficial uses as a result of contaminants from the mines. In most cases it appears that mine effluent is diluted by the time it gets to sections of the River that support a beneficial use. Additional sampling during the spring flush may be warrented. However, I'd suggest limiting samples to sites supporting a beneficial use.

Post Script

I talked to Mike Reichert about the data and my strict interpretation of water quality standards and what constitutes violations. I found that Mike takes a much more liberal view than I do. He feels that a stream doesn't necessarily have to be able to support a beneficial use for the standards to apply. For example, the No Ft American the above the waterfall might be in violation of state standards even though fish would not normally be found there.

He suggested that you get someone from his office (and maybe Dwk as well) to look at the streams and make their judgements.

WATER QUALITY STANDARDS FOR 3A STREAMS

Arsenic 360 mg/l (1 hour) Cadmium 3.9 ug/l (1 hour) Copper 18 ug/l (1hone) Lead 82 ug/l (1 tions) Mercury 2.4 ugle (16our) Selver 4.1 ug/l (1hour) Barium Chromum 1700 ugll (1 hour) Iron 1000 ugk (1hour) Manguiresc Selenium 20 right (thour) Zinc 120 mg/l (thour) Sulfate + -

* No standard found for 3A streams (cold Water fisheries)



State of Utah

DEPARTMENT OF NATURAL RESOURCES DIVISION OF OIL, GAS AND MINING

Dee C. Hansen Executive Director Dianne R. Nielson, Ph.D. 355 West North Temple 3 Triad Center, Suite 350 Salt Lake City, Utah 84180-1203 801-538-5340

August 7, 1990

Mr. Paul Skablund Uinta National Forest 100 North 88 West Provo, Utah 84601

Dear Mr. Skablund:

The Utah Abandoned Mine Reclamation Program is concerned with mitigating physical hazards to the public health and safety that occur on abandoned mine sites. We would be able to provide you with plans and specifications for typical closure techniques to secure abandoned mine portals and other work specifications for earthwork, demolition, etc. Possibly a staff member could discuss with you how to draw up reclamation plans for particular sites.

If health hazards are present due to toxic substances, the Utah Department of Health, Environmental Health section, usually takes responsibility.

Possible solutions to the Pacific and Mary Ellen Gulch mines, where extensive tailings dumps are present would be to: 1) route runoff around the dumps and try to stabilize the dumps in place, or 2) remove the dump materials to a lower precipitation site. Removing the dumps would be expensive, logistically difficult and could aggravate the problems present by introducing oxygen into the system. Off-road vehicle use should be prevented at the Pacific Mine tailings area.

From the information in Dr. Merritt's report, elevated levels of cadmium, copper, lead and zinc are present but confined somewhat to localized areas within a mile of the discharge point. Methods to lower these levels are generally prohibitively expensive. It does appear that some of the parameters sampled, particularly copper, lead and zinc increase substantially after flowing through the dump material. Thus, I would recommend preventing, as much as possible, all runoff from flowing over or through the dump.

Please call me if you would like to discuss this further. I would appreciate it if you would keep me informed about the progress of this project.

Sincerely, Sweed Malin

Lucia Malin'

Senior Reclamation Specialist

an equal opportunity employer
IIII
AM806/186

No.110418109106

PARTICIPATING AGREEMENT

between

THE UTAH DIVISION OF OIL, GAS & MINING

and

THE USDA FOREST SERVICE, UINTA NATIONAL FOREST

THIS AGREEMENT is made and entered into by and between the Utah Division of Oil, Gas and Mining (Division, also known as the cooperator) under state and federal law and implementing regulations (Section 40-10-1 et seq. U.C.A. and P.L. 95-87, and the USDA Forest Service (FS), under the provisions of the Act of December 12, 1975, 16 U.S.C. 565A.

WITNESSETH:

WHEREAS, the Division is the designated state agency responsible for implementation of a statewide program for the reclamation of abandoned mines, and

WHEREAS, both parties are interested in the conservation of our nation's natural resources, and inasmuch as that interest extends to the protection and management of lands that are threatened by the adverse effects of abandoned mines, and

WHEREAS, the abandoned Pacific Mine, the Lower Bog Mine, the Miller Hill Drain Tunnel site, the Mary Ellen Gulch Mine and other smaller unnamed sites in American Fork Canyon are FS Region 4 areas of concern for the abatement of pollution in the Wasatch Mountains of Utah, and

WHEREAS, the Uinta National Forest contains a number of acres of mineral mine wastes from these abandoned mines which have high levels of lead, zinc, cadmium and other heavy metals, causing highly acidic drainage to enter the American Fork River, and

WHEREAS, to manage and protect the lands and waters in the vicinity of abandoned mines, their effects need to be studied and remedies need to be proposed, and

WHEREAS, it is mutually advantageous to the parties herein to share in the study and correction of abandoned mine impacts on the environment and propose remedies for affected areas within the Uinta National Forest and publish a report thereon.

NOW THEREFORE, in consideration of the above, the parties agree to cooperate on a reclamation project the total direct value of which is estimated at \$4,750, to be equally divided between the parties as shown in the Attachment.

A. THE FS SHALL:

- Provide initial advice and assistance as needed to formulate and meet project objectives.
- Provide aerial photography and other supplies at their (FS) costs to meet project needs efficiently.
- 3. Reimburse the Division for the cost of the project, up to \$2,375 as provided for in the attached financial project plan, upon delivery of the completed report and upon receipt of an itemized listing of project expenditures. Advance payments shall not be made.
- Designate Paul H. Skabelund or his designated replacement to represent its interest in this effort.
- Provide the Division with permission to study the areas, of abandoned mine problems which it manages, for solutions to the acid mine drainage problems.

B. THE DIVISION SHALL:

- Provide manpower, equipment and supplies through its Abandoned Mine Reclamation Program to complete the selection and management of abandoned mine investigation and remediation proposals as outlined in the project work plan.
- Provide funding and services up to \$7,886 as shown in the attached financial project plan for completion of the project as outlined in the project work plan.
- 3. Prepare and present to the FS a report that summarizes the results of the field investigations and proposes plans and cost estimates for remediation of the abandoned mine problems as outlined in this agreement.
- 4. Be alert to the presence of other abandoned mine problems known or suspecced to exist on the Uinta National Forest in the process of performing the work plan.

- Provide the Uinta National Forest with any other information collected incidental to the completion of this project.
- Designate Mary Ann Wright or her replacement as the Division employee responsible for carrying out its part of this agreement.
- 7. Bill the Forest Service upon completion of the project.
- 8. Coordinate with any landowners as necessary to obtain permission to enter and study the past effects of abandoned mine lands on private lands adjacent to the FS lands.
- Coordinate with other state agencies as necessary to accomplish the proposed work.
- 10. Give USDA Forest Service or the Comptroller General, through any authorized representative, access to and right to examine all records, books, papers or documents related to the award.

C. IT IS MUTUALLY AGREED AND UNDERSTOOD THAT:

- The work under this agreement shall be completed no later than December 31, 1992.
- The FS and the Division shall work together to reach a mutually acceptable reclamation plan for the American Fork sites.
- 3. The FS or the Division, in writing, may terminate the agreement in whole, or in part, at any time before the date of expiration, whenever it is determined that the other party has materially failed to comply with the conditions of this agreement. The other party shall not incur any new obligations for the terminated portion of the agreement after the effective date, and shall cancel as many obligations as is possible. Full credit shall be allowed for the FS share of the obligations incurred to the effective date and all non-cancelable, properly incurred obligations by the cooperating party (Division) prior to termination.

- 4. Any monies that are payable from the United States under this agreement to any person or legal entity not an agency or subdivision of a State or local government may be subject to administrative offset for the collection of any debt the person or legal entity owes to the United States. Information on the person's or legal entity's responsibility for a commercial debt owed the United States shall be disclosed to consumer or commercial credit reporting agencies.
- 5. The cooperator (Division) shall comply with Title VI of the Civil Rights Act of 1964, that no person in the United States shall, on the ground of race, color, handicap or national origin, be excluded from participation, be denied the benefits of, or be otherwise subjected to discrimination under any program or activity for which the recipient receives Federal financial assistance and will immediately take any measures necessary to effectuate this agreement.
- 6. Improvements placed on National Forest land at the direction of either of the parties, shall thereupon become the property of the United States, and shall be subject to the same regulations and administration of the FS as all other National Forest improvements of a similar nature.
- 7. This agreement in no way restricts the FS from participating with other public or private agencies, organizations, and individuals or from accepting contributions and/or gifts for the improvement, development, administration, operation and preservation of this or any other project.
- 8. No part of this agreement shall entitle the cooperator (Division) to any share or interest in the 'project other than the right to use and enjoy the same under the existing regulations of the FS.

Page 5 Agreement

- 9. No member of, or Delegate to Congress, shall be admitted to any share or part of this agreement, or any benefits that may arise therefrom; but this provision shall not be construed to extend to this agreement if made to a cooperation for its general benefit.
- 10. Nothing herein shall be considered as obligating the FS to expend or as involving the United States in any contract or other obligations for the future payment of money in excess of appropriations authorized by law and administratively allocated for this work.
- Persons provided as contributed labor under this agreement shall be considered as federal employees for the purposes of tort claims and compensation for work injuries.
- 12. This agreement may be revised as necessary by mutual consent of both parties, by the issuance of a written amendment, signed and dated by both parties.

Page 6 Agreement

The parties hereto have executed this agreement as of the last date written below:

DIVISION OF OIL, GAS & MINING

USDA FOREST SERVICE

Dianne R. Nielson, Director

Date: 9/18/91

DEPARTMENT OF NATURAL RESOURCES

Dee C. Hansen Executive Director

DIVISION OF OIL, GAS & MINING

Carl Roberts, Budget Officer

DIVISION OF FINANCE

Gordon L. Crabtree, Director

nate: 9/35/41

Agreement Attachment Page 1

AMERICAN FORK CANYON ABANDONED MINE STUDY MOA PROJECT BUDGET AND PLAN

PHASE I

Task 1	Contracted hydrolog sites.	gic analysi	is: Site analyses f	or three
	Hydrologist	20 hours	@ \$50/hour	\$1,000
	Technician	25 hours	@ \$25/hour	\$ 625
	Travel		lump sum	\$1,000
	Supplies and equip	ment	lump sum	\$ 400
Task 2	Contracted hydrolog of technology asses development.			
	Hydrologist	20 hours	@ \$50/hour	\$1,000
	Drafter	5 hours	@ \$25/hour	\$ 125
Task 3	Hydrologic analysis	s report.		
	Hydrologist	8 hours	@\$50/hour	\$ 400
	Clerical	10 hours	@ \$20/hour	\$ 200
			TOTAL	\$4,750

PHASE II

USFS and Division coordination to reach an acceptable reclamation plan.

PHASE III

Task 4 Division design and construction specification development.

Program Administrator	60 hours	@\$20/hour	\$1,200
Sr Reclamation Spec'lst	160 hours	@13/hour	\$2,080
Clerical	20 hours	@8/hour	\$ 160
		SUBTOTAL	\$3,440
Benefits		0 33%	\$1,135
Travel		lump sum	\$ 250
Other Direct			\$ -0-
Current Expense	15% of Sa (\$4,575)	laries & Benefits	\$ 686
		TOTAL	\$5,511

PHASE IV

Review by both parties and implementation of plan.

Agreement Attachment Page 3

BUDGET SUMMARY

FOREST SERVICE CONTRIBUTION		
Value of contributed labor		\$1,000
50% share of contracted cos	sts ¹	\$2,375
SUBTOTAL DIVISION CONTRIBUTION	\$3,375	
Value of contributed labor		\$5,511
50% share of contracted cos	sts	\$2,375
SUBTOTAL	\$7,886	
	PROJECT TOTAL	\$11,261

Total FS Share 50 % of Direct Consulting Costs Division Share 50 % of Direct Consulting Costs

¹Estimated payment to the Division. Reimbursement will be made only upon actual expenses incurred by the Division, not to exceed estimated payment.

Agreement Attachment Page 3

BUDGET SUMMARY

FOREST SERVICE CONTRIBUTION		
Value of contributed labor		\$1,000
50% share of contracted cos	ts ¹	\$2,375
SUBTOTAL DIVISION CONTRIBUTION	\$3,375	
Value of contributed labor		\$5,511
50% share of contracted cos	ts	\$2,375
SUBTOTAL	\$7,886	
	PROJECT TOTAL	\$11,261
建筑的		

Total FS Share 50 % of Direct Consulting Costs Division Share 50 % of Direct Consulting Costs

¹Estimated payment to the Division. Reimbursement will be made only upon actual expenses incurred by the Division, not to exceed estimated payment.



Norman H. Bangerter Oovernor Dee C. Hansen Executive Director Dianne R. Nielson, Ph. D. Dotsion Director

DATE:

FAX #

ATTN:

an equal opportunity amployer

State of Utah

DEPARTMENT OF NATURAL RESOURCES DIVISION OF OIL, GAS AND MINING

355 West North Temple 3 Triad Center, Suite 360 Sell Lake City, Ureh 84180-1203 801-538-5340

5-8-92

1-375-0821

Paul Skelbland

UTAH DIVISION OF OIL, GAS AND MINING FACSIMILE TRANSMISSION COVER SHEET

COMPANY	U.S. F.S
FROM:	Mark Mesch
DEPARTMENT:	oil Gas + ming
NUMBER OF PA	GES BEING SENT (INCLUDING THIS ONE): 8
	t receive all of the pages, or if they are illegible, (801) 538-5340.
We are sendi number is (8	ng from a Murata facsimile Machine. Our telecopier 01) 359-3940.
MESSAGES:	
	Review and comment on the Scop
	nk in this contract.

ATTACHMENT C

SCOPE OF WORK

- C.1 GENERAL: The Scope of Work includes a hydrological analysis of water quality associated with three metal mine sites; development of design recommendations with cost proposals, to reduce impacts of mine water entering the American Fork River, and preparation of a hydrological report discussing the water quality analysis and the preferred design recommendation. Each segment of the WORK is divided into the following Tasks:
 - Task I Hydrological Analysis (water quality)
 - Task II Hydrological Design: Application of technology assessment, design and cost proposal development
 - Task III Hydrological Report, Recommended Design
- C.1.2 THE WORK: The objective of the WORK is: I) to conduct a hydrological analysis (water quality) to determine impacts of mine water to off site water sources; II) based on Task I findings, develop a mite low cost, aesthetically pleasing, and hydrological stable design recommendations, with cost proposals; and III) submit a written hydrological report disoussing the analyses performed, data collected and results obtained as well as preferred design recommendations.
- C.1.3 If technical deficiencies or engineering related problems are encountered within the Scope of Work and technical specifications during future bidding or construction activity, the OWNER will require further information or verification of assumptions from the CONTRACTOR. It is expected that if such deficiencies are found, the CONTRACTOR will act to alleviate and resolve any conflicting, missing or unsubstantiated information found within the Construction Specifications.
- C.1.4 It is not the intent of the OWNER to bind the CONTRACTOR to work not included as part of the WORK. The intent of the OWNER is to require the CONTRACTOR to complete the specifications accurately and in the detail sufficient to perform the construction work.

C.2 TASK I - HYDROLOGICAL ANALYSIS, WATER QUALITY

The purpose of Task I is to conduct the necessary hydrological analyses to determine the current water quality conditions at the Pacific Mine, Mary Ellen Gulch, and the Lower Bog Mine and their associated impacts to the American Fork River. This will involve sampling for the following Total metals: arsenic, barium, cadmium, chromium, lead, mercury, nitrate (as N), selenium, silver, copper, iron, manganese, sulfate, zinc, and total dissolved solids (TDS). Additionally, the following field measurements will be taken: pH, conductivity, dissolved oxygen, temperature, color and rate of flow. Samples will be taken at the following locations: A) portal openings (3 samples); B) as water leaves the tailings piles (3 samples); C) at the beaver pond adjacent to the Pacific Mine (2 samples); D) locations above and below mine sites along the American Fork River (3 samples). It is estimated that eleven points will be sampled during the low flow period. Exact sampling points will be identified on a quad map of the area.

C.3 TASK II - HYDROLOGIC DESIGN, APPLICATION OF TECHNOLOGY ASSESSMENT, COST PROPOSAL

CONTRACTOR will prepare a suite of site specific design alternatives based on TASK I findings to reduce the impacts of mine water entering the American Fork River. Low cost and an aesthetic, natural appearance are critical design factors in addition to effective performance.

- C.3.1 Structural design specifications of the recommended design will be presented in a manner to fit the OWNER's contract bid specifications.
- C.4 TASK III HYDROLOGICAL ANALYSIS REPORT, DESIGN RECOMMENDATIONS
- C.4.1 CONTRACTOR will submit to the OWNER a written report discussing the nyurorogical findings, (fact 1), and the options, with associated costs (Task II), and the recommended design, containing, if necessary, formatted Technical Specifications and Site Specific Requirements.

DIV UIL UNO 6 HAHAMO

- C.4.2 CONTRACTOR will provide these site specific specifications to be compatible with the 0300 Section format used by the OWNER. When the WORK specified is not work previously conducted by the OWNER, CONTRACTOR will provide technical directions, in standard ASTM or comparable format and compatible with the 0200 Section format used by the OWNER. As part of the WORK Specifications, the following is included as part of the WORK:
- C.4.3 Detailed areas for specific construction and grading will be drawn at a scale and with contour intervals as appropriate and approved by the OWNER.
- C.4.4 When 95% of the final design WORK is complete, CONTRACTOR will accompany the OWNER'S representative to the project site for the purpose of field verification of the design plans.

End of Attachment C

ATTACHMENT D SCHEDULE OF PRICES & COST SHEETS

- D.1 FIXED NOT-TO-EXCEED PRICE AMOUNT
- D.1.1 CONTRACTOR hereby agrees to perform the WINK, TASK T, TI, and III, as described in this Agreement and the OWNER agrees to pay CONTRACTOR In the amount not to exceed \$ 4.750.00 for said WORK, as a FIXED NOT-TO-EXCEED AMOUNT.
- D.1.2 OWNER shall pay amounts invoiced only at UNIT COST PRICES submitted by the CONTRACTOR on a regular basis.
- D.2 UNIT COST PRICES
- D.2.1 Unit prices provided by the CONTRACTOR and approved by the OWNER are attached to and considered part of this agreement. These are considered UNIT COST PRICES for each line item and the FIXED NOT-TO-EXCEED PRICE for the total sum of all items contained in the WORK.

D.2.2 COST SUMMARY SHEETS

D.2.3 The following cost summary sheets were completed from information supplied by the CONTRACTOR in negotiations with the OWNER and are binding as part of the AGREEMENT upon execution.

D.2.4 TOTAL FIXED-PRICE COST

Task I	HYDROLOGICAL ANALYSIS, WATER QUALITY \$3,025.00
Task II	HYDROLOGIC DESIGNS, APPLICATION OF TECHNOLOGY ASSESSMENT, COST PROPOSALS \$1,125.00
Task III	HYDROLOGICAL ANALYSIS REPORT, DESIGN RECOMMENDATION \$ 600.00

Total Fixed Not-To-Exceed Price

\$4,750.00

D.2.5 BREAKDOWN OF COSTS BY TASK

TAGE T HEPROLOGICS	I ANALYSIS. WATER	QUALITY	
<u>Item</u>	Units	Rate	Extension
Hydrologist Technician Travel Supplies and squipment Total Task I	20 hr 25 hr	\$50 \$25 lump sum lump sum	\$1,000.00 \$ 625.00 \$1,000.00 \$ 400.00 \$3,025.00

TASK 11 - HINKLING DESIGN, APPLICATION OF TROUNDINGY ASSESSMENT,

Item	Units	Rate	Extension
Hydrologist Drafter Total Task II	20 hr 5 hr	\$50 \$25	\$1,000.00 \$ 125.00 \$1,125.00

TASK III HYDROLOGICAL ANALYSIS REPORT, DESIGN RECOMMENDATIONS

Item	Units	Rate	Extension	
Hydrologist Clerical Total Task III	8 hr 10 hr	\$50 \$20	\$ 400.00 \$ 200.00 \$ 600.00	

TOTAL ALL TAEXS \$4,750.00

End of Attachment D

ATTACHMENT E

TIME SCHEDULE

- E.1 TIME SCHEDULE: Schedule is shown in Attachment E.1.1 in calendar form, on a daily schedule, the activities required for the WORK described in this Agreement for each activity based on actual start date for the WORK.
- E.1.1 Constraints and float allowed for in the calendar are identified and explained for in the schedule. Considerations for weather, access and normal delays are accounted for in these constraints. Deviations from this schedule must be approved in writing by OWNER. All deliverables are due by August 31, 1992.

June - July - August

Task I 1 - 30

Task II 1 - 31

Task III 1 - 31

End of Attachment E

ATTACHMENT F

SERVICES AND FACILITIES PROVIDED BY OWNER

- F.1 OWNER shall provide the following services for the CONTRACTOR with respect to the WORK required:
- F.2 Any site-specific information that the OWNER has that would be helpful to the CONTRACTOR in accomplishing the WORK.
- F.3 OWNER'S representative will be available to accompany the CONTRACTOR to facilitate location of the site features, assist with sample collection, collaborate on design, and respond to questions.

- F.4 Access to resources in the OWNERS possession that may aid in the completion of the WORK, such as:
 - a) All project and site files, project correspondence, slides, photographs, and any maps relating to the project area.
- F.5 Funds for the analysis of water samples at the State of Utah Water Laboratory will be the responsibility of the OWNER and are not included in this contract amount.


End of Attachment F

ATTACHMENT G

AMENDMENTS TO THE AGREEMENT

- G.1 OWNER shall require amendments to the Agreement to be in the form of a change order, signed by both parties and similar in form to the change order shown in this Attachment.
- G.2 Change orders shall become attached to and part of the Agreement under the terms of the Agreement with changes as stipulated on the change order and shall not release the CONTRACTOR from any other terms or conditions that apply and are a part of the Agreement.

End of Attachment G





Norman H. Bangerter Governor Dee C. Hansen Executive Director Dianne R. Nicison, Ph. D.

State of Utah

DEPARTMENT OF NATURAL RESOURCES DIVISION OF OIL, GAS AND MINING

355 West North Temple 3 Triad Center, Suite 350 Salt Lake City, Utah 84180-1203 801-538-5340

January 20, 1993

Mr. Paul Skabelund Uinta National Forest 88 West 100 North Provo, Utah 84601

JAN 25 1993

cc: D-Z

Dear Mr. Skabelund:

Re: American Fork Mine Site Analyses and Reclamation Recommendations

Enclosed please find one bound original and one unbound copy of Lidstone & Anderson's American Fork Hydrology and Water Quality Study. The report, in addition to supplementing the earlier water quality work of Merritt, examines the geochemistry of the area, identifies biological, geochemical, and hydrological controls at each site, and develops mitigation alternatives and recommendations. The report also acknowledges a need for further data collection and analysis. Below I have attempted to summarize the salient issues in the report.

ABIOTIC AND BIOTIC FACTORS

Two abiotic factors, geological and hydrological, are operating to reduce the severity of the off site impacts of the Pacific Mine, mines in Mary Ellen Gulch, and the Lower Bog Mine:

1) high buffering capacity due to a host rock rich in carbonates; and 2) high dilution ratios, up to 33:1 at the Lower Bog Mine. These factors result in a change in pH values measured at the mine portals and downstream of 5.1 to 7.52 at the Lower Bog Mine, 6.5 to 8.02 at the Pacific Mine, and 6.95 to 7.95 at Mary Ellen Gulch.

The beaver pond at the Pacific mine appears to play a significant biotic role in removing trace elements from the portal effluent, specifically, zinc, cadmium, and lead. The effectiveness of the beaver pond clearly identifies its potential role in any reclamation activity undertaken at the Pacific Mine.

MITIGATION ALTERNATIVES

Pacific Mine

Based on the analysis of the data collected, reclamation at the Pacific mine is the highest priority. Two sources of

Page 2 Mr. Paul Skabelund January 20, 1993

These problems could be addressed in two or three phases. Phase one would route all portal drainage off the tailings pile and via a riprap ditch into the beaver pond. Phase two would isolate, recontour and treat, topsoil, and revegetate the tailings dumps. A third phase would develop a wetland above the beaver dam to provide additional treatment to the portal effluent if water sampling after completion of phase one indicated a decrease in the ability of the beaver pond to treat the portal discharge adequately.

Lower Bog Mine

Due to the inaccessibility of the Lower Bog mine, the limited magnitude of the problem it presents, and the high dilution ratio (33:1), no reclamation action is recommended.

Mary Ellen Gulch Mines

A suite of problems exist at the Mary Ellen Gulch site ranging from trace metal contamination in the creek to active mining exploration in the Belorophan mine and at the Yankee dumps. Samples taken in Mary Ellen Creek identified contamination but an insufficient number of samples were collected to fully characterize the source. The sample identified as AF#7 taken from the most northerly portal on the mine bench did not show elevated metals except for zinc, suggesting some other source of contamination exists. This could possibly be from the tailings piles or the mining activity occurring at the Belorophan mine. The Utah Division of Oil, Gas and Mining's Minerals Program issued a Small Mining Operations permit for the "Yankee Project" in August of 1992, after we noticed mining activity taking place while sampling in the area. The operator, James Warr, was advised by DOGM's Minerals Program in a July 27, 1992 letter of the following issues: 1) that the Forest Service was very concerned about any off site impacts to Mary Ellen Creek and surrounding areas; 2) that if old workings were developed a UPDES permit would be required from the Division of Water Quality; and 3) that the mine dumps had been placed on the CERCLIS list and that the operators might be responsible for some expensive CERCLA cleanup.

Due to the complexity of the situation at the Mary Ellen Gulch site, further study is warranted to identify the specific source(s) of contamination and allow for some resolution to occur with regards to the mining activity prior to developing reclamation alternatives for this site.

Page 3 Mr. Paul Skabelund January 20, 1993

Miller Hill Tailings

Water sampling was not performed at this site. No portal discharge is occurring, and the adverse effects are more a result of erosive conditions along the toe of the pile during high water stream flows. Tailings samples taken by the Forest Service and analyzed by Utah State University Soil Testing Laboratory for crop production/vegetation success are within the range for plant establishment and growth. Revegetation, however, does not remedy the problem that the location of the tailings presents to the North Fork of American Fork. Based on the rather small areal extent of the tailings pile and the ease of access, removal may be the best alternative.

Utilizing the tailings as road surface material may be an effective form of disposal. However, the following precautions are warranted. Testing for total soluble metals is recommended. Soluble metals leaving road surfaces during rain storms or as snowmelt could be problematic and affect off site areas. Spreading the tailings out over a large area (i.e. roadway) would also increase the oxidation rate by increasing the surface area of the tailings, as opposed to keeping the tailings confined as a single deep pile. Tailings should be mixed with locally obtained limestone material prior to placement as road surface. This would continue to buffer the tailings material once in place on the road surface. Finally, any road sites selected for tailings placement should be situated away from water courses.

Using the report's recommendations for the Pacific mine, I will develop construction costs for the work phases. After you have had time to review the report we can arrange for a meeting to discuss the report and the direction the Forest Service wishes to take in addressing the reclamation at these sites.

Sincerely,

Mark Mesch

Reclamation Specialist

Abandoned Mine Reclamation Program

MRM Enclosures WP.Skabelun.Let

PART IV

Sample Results From Water, Soil, and Tailings at the Pacific Mine, Mary Ellen Gulch Mine and the Lower Bog Mine (cronological order)

TABLE 2.14.2

NUMERIC CRITERIA FOR AQUATIC WILDLIFE

Parameter		Wildlife		
	S.B.A.L.	3B · · ·	3C	30
PHYSICAL				
TOTAL DISSOLVED GASES	(1)	(1)		
DISSOLVED OXYGEN (MG/L) (2)				
30 DAY AVERAGE	6.5	5.5	5.0	5.0
7 DAY AVERAGE	9.5/5.0	6.0/4.0		
1 DAY AVERAGE	8.0/4.0	5.0/3.0	3.0	3.0
MAX. TEMPERATURE (C)	20	27	27	
MAX. TEMPERATURE CHANGE (C)	2	4	4	
pH (RANGE)	6.5-9.0	6.5-9.0	6.5-9.0	6.5-9.0
TURBIDITY INCREASE (NTU)	10	10	15	15
METALS (3)				
(ACID SOLUBLE, UG/L) (4)				
ARSENIC (TRIVALENT)				
4 DAY AVERAGE	190	190	190	190
1 HOUR AVERAGE	360	360	360	360
CADMIUM (5)				
4 DAY AVERAGE	1.1	1.1	1.1	1.1
1 HOUR AVERAGE	3.9	3.9	3.9	3.9
CHROMIUM (HEXAVALENT)				
4 DAY AVERAGE	11	11	11	11
1 HOUR AVERAGE	16	16	16	16
CHROMIUM (TRIVALENT) (5)				
4 DAY AVERAGE	210	210	210	210
1 HOUR AVERAGE	1700	1700	1700	1700
COPPER (5)				
4 DAY AVERAGE	12	12	12	
1 HOUR AVERAGE	18	18	18	18
CYANIDE (FREE)				
4 DAY AVERAGE	5.2	5.2	5.2	
1 HOUR AVERAGE	22	22	22	22
TRON (MAXIMUM)	1000	1000	1000	1000
LEAD (5)				
4 DAY AVERAGE	3.2	3.2	3.2	3.2
1 HOUR AVERAGE	82	82	82 .	82

TABLE 2.14.2, CONTINUED

Manage	c Wildlife			
3A .	38	3C	3D	
				61 3
0.012	0.012	0.012	0.012	
2.4	2.4	2.4	2.4	
160	160	160	160	
1400	1400	1400	1400	
5.0	5.0	5.0	5.0	
20	20	20	20	
0.12	0.12	0.12		
4.1	4.1	4.1	4.1	
110	110	110	110	
120	120	120	120	
(6a)	(6a)			
(6b)	(6b)	(6b)	(6b)	
0.011	0.011			17.00
0.019	0.019	0.2	(8)	
2.0	2.0	2.0	2.0	
0.01	0.01	0.01	0.01	
	0.012 2.4 160 1400 5.0 20 0.12 4.1 110 120	0.012 0.012 2.4 2.4 160 160 1400 1400 5.0 5.0 20 20 0.12 0.12 4.1 4.1 110 110 120 120 (6a) (6a) (6b) (6b) 0.011 0.011 0.019 0.019 2.0 2.0	3A 3B 3C 0.012 0.012 0.012 2.4 2.4 2.4 160 160 160 1400 1400 5.0 5.0 5.0 20 20 20 0.12 0.12 0.12 4.1 4.1 4.1 110 110 110 120 120 (6a) (6b) (6b) (6b) 0.011 0.011 0.019 0.019 0.2 2.0 2.0 2.0	3A 3B 3C 3D 0.012 0.012 0.012 0.012 2.4 2.4 2.4 2.4 160 160 160 160 160 1400 1400 1400 5.0 5.0 5.0 5.0 20 20 20 20 0.12 0.12 0.12 4.1 4.1 4.1 4.1 110 110 110 110 120 120 120 (6a) (6b) (6b) (6b) (6b) 0.011 0.011 0.019 0.019 0.2 (8) 2.0 2.0 2.0 2.0 2.0

HT-NUMBER

A & L MID WEST AGRICULTURAL LABORATORIES, INC.

13611 "B" Street • Omaha, Nebraska 68144-3693 • Phone: (402) 334-7770

19/85

ACCOUNT NO.

3613 . . . GRC

PO 43-8460-5-259



UINTA NATIONAL FOREST 88 WEST 100 NORTH 80X 1428 PROVO UT 84603

SUBMITTED BY:

SOIL ANALYSIS REPORT

(SEE EXPLANATION ON BACK

ORGANIC	SECTION SECTION	HOSPHORUS	Bussell .	POTASSIUM	MAGNESIUM:	E CALCIUM A	SODIUM .	ASSE PH	SANTE HAL		T. Lautette	13 3 d S H	PERCENT	PRIME	1415
	(WEAK BRAY)	ISTRONG BRAY)	BICARBONATE	ppm 7-2 RATE	Mg Mg PATE	ppm RATE	Na RATE	SOIL 19	BUFFER 1 GI INDEX meq/	EXCHANGE CAPACITY C.E.C.	E VIE	BASE SA	TURATION (C	OMPUTEDI TEL MAR	1 20
12285 0.40[4.41	100	200		39M	6VL	110M	22H	3.9	7.1 0.	1 1.0	10.5	5.2	57.6	16.71	0.0
12286 1.2(56 12287 1.01 52	3VL	101		31VL	8VL 45VH	30VL	20VL	Your earlier	7.1 0.		4.7	22.4	59.7	94.4	15167
12288 2.84 82	110	16L		44VL	26VL	160VL	28VL			* 20.5	0.5	1.1		10.13	0.6
	7:1			D Tage	1					4	17/				a sud
	TPA EX	TRACT	ton	2 1 1 1 1 T				***	COMM	Committee of the commit					
	MANGA NESEE	IRON S	COPPER Cu	BORON?	EXCESS SOLUB LIME SALT	Company of the Compan					7.7				
	opin RATE		ppm RATE		mmhos/	A		15 Table		A Maria	111	in the	445		
		T 8 A V H	3.2VH	0.400	L 0.4			1	- Av						
	301	AUL	14.00H	B. K.	L 0.81	-				es con de la compania			1	.1.	
4 9900 11901	200	176VH	0.91	0.4VL	L 2.0	44		1		ort applies only t	o the sample	(a) tested Sa	mples are rel	ained a maxir	mum of
			50			13/		- Andrews			WEST AG	RICULTURAL	LABORATO	ORIES, INC.	
ING VERY LOW (VL), LOW (L), MEDRUM (M), H TED NITROGEN RELEASE FRISH IS N NOR BY	IGH (H), VERY HIS	GH (VH), AND NO	NE (N)	MULTIPLY THE MULTIPLY THE	RESULTS IN ppm BY RESULTS IN ppm BY 0 (2) MILLION POLINE	4.6 TO CONVERT TO 2.4 TO CONVE	LBS. PER ACRE P20 S. PER ACRE K20 ACRE OF SOIL 6-	5				Toh)		1	



SOIL SAMPLE INFORMATION SHEET

	A & L MID WE									
FALER ACCOUNT	GROWE	n - 1 - 1		i e	OPY TO		30	w		E
_ Uinta National Forest	Po#43-8	4A0-5-2	\$9			K) (4	41 M + 14			
P.O-BOX 228										
Pl. Grove Ut. ZIP84062 PHONE 1801 1 785-3563		ZIP				ZIP		0 00	FOFAS	
	CROP PLUS (\$1.00 PER REP	ORTI						2 03	5854 ^s	
MINARES STANDARD MICRONUTRIENT	NOIVIDUAL ANALYSIS	IF FERTILITY RECOMMEND	REGON ATION	MMENDA1 IS CHARG	IONS E INC	ARE R LUDES	EQUIRED TWO CR	COMPL OPS OR	ETE THIS SEC YIELD LEVELS	TION.
/xx × V	12285	CROPIBI TO BE BROWN YIELD	PALISI PA	EVIOUS CROPISI	PREV. YLD.	BAMPLE DEPTH	PERTILIZER APP			C SST CSTADISKI
			-			TO		1	O BOLLONE O PLANTED	SPERIFICIAN D THES
		CROPISI TO SE GROWN (VIELD I	DOALISI PR	EVIOUS CROPIS)	PREV. YLO.	SAMPLE DEPTH	SARY GO		COVET	, IMPORTED TES O
2 ×× ×	12286					10		K ₂ O	O HILLY O UPLANS	SRAVITY D. OTHER SPRINKLER D. C
			1/8		Tyler.	REMARK	Any G	round	manufacture and the latter to	
		CROPIES TO BE GROWN VIELD	DOALIER PR	EVIOUS CROPISI	PREV. YLD.	BAMPLE DEPTH	PERTICIZER APP	THED THE		C SET GSTABIBAL;
3 x x x V	12287					10		K ⁸ O	O MILLY O UPLANO	ENEVERSE D I GALVER
						REMARK		roune		
444 2 2	12288	CROPES TO SE GROWN VIELD	SOALISI PR	EVIOUS CROP(S)	PREV. YLD.	BAMPLE DEPTH TO	FENTILIZER APP	K ⁶ O AS-	TOPOGRAPHY OMELY DUPLAND DESTROM DUPLAND	SASVITY O STINES
						REMARK	s.Any (STOK	10	
		CROPES TO BE BROWN YIELD	SOALISI PE	HVIOUS CROP(E)	PREV. YLD.	SAMPLE DEPTH	PERTYLIZER APP	K _E O		F SHEWLATED YES CO
					10 10 13	70			DHILLOR DINEWO	BASSALLA D 2 GAMES
1011	1			,	- 1	REMARK				
EMARKS: Need Complete 2 Certifizer needs (PLANATION OF TESTS: Organic matter, estimated nitrogen release, phosph hydrogen, exchange capacity, percent base saturet Same as 1A except sodium bicarbonate P in place of Soluble stits, excess time and sodium run in combin 3 Gulfur, boron, zinc, manganese, iron, copper	orus (P1, P2), potassium, magnes cion. of P2.	ium, calcium, soil pH, salt		S	= Sulfur = Zinc = Manga = Iron = Copper = Boron	nese	Signature Of Signature Samples S	Shipped O	Ish A. Wes	Vondl
Zinc, manganese, iron, copper 5 Suffur, zinc. Suffur,zinc, boron esa.	LABORATOR	Y COPY	0	Mo CI NO3-N	= Molybo = Chlorid = Nitrate	0			O Omeha, Nebreska) INC.

A & L MID WEST AGRICULTURAL LABORATORIES, INC.

13611 "B" Street • Omaha, Nebraska 68144 • Phone: 402-334-7770

Secretary to the second

FEDERAL ID NUMBER 47-0564465



UINTA NATIONAL FOREST BB WEST 100 NORTH BOX:1428 PROVO: UT 84603

INVOICE NUMBER B6622

INVOICE

PORT NO.	GROWER OR SUBMITTED BY	NUMBER OF SAMPLES — TYPE OF ANALYSIS	TOTAL
		4.82/4.83/4.Nitrote Nitrogen Subtotal Less 25.00% discount	95.20 95.20 23.60
	BOND SERVICE CONTRACTOR OF THE SERVICE CONTR		正江西江 隆
			T I
			- 1 Z
of the experiment			HUNGAN
326			
7		CONTROL CONTROL CONTROL OF THE CONTR	THE PURE
24307			
			.1120 12
41.75.0			LETT

BILLING D

RTMENT

INTEREST WILL BE CHARGED ON OVER DUE BALANCES AT THE MAXIMUM RATE ALLOWABLE BY LAW, PAYABI COUNTY OF ISSUE.

UTAH STATE DEPARTMENT OF HEALTH
Office of Administrative Services
Bureau of Finance
P.O. Sox 16700
Salt Lake City, Utah 84116-0700

U.S. DEPARTMENT OF AGRICULTURE NATIONAL FINANCE CENTER PO BOX 60075 NEW ORLEANS LOUISIANA 70160

Date: September 1, 1988

Reference:

QUANTITY	DESCRIPTION	PRICE	AMOUNT
	LABORATORY TESTS PERFORMED:		Lament-2
7 ea.	Water-SO4, T HDNS, T ALK, TDS, Metals	\$95.00	\$665.00
1 ea.	Water-SO4, T HDNS, As, Pb, T ALK, TDS, Cd, Zn	65.00	65.00
1 ea.	Water-T ALK, As, Cu, Hg, TDS, Cd, Pb, Zn	75.00	75.00
1 ea.	Water-SO4, T HDNS, As, Cu, Zn, T ALK, TDS, Cd, Pb	75.00	75.00
1 ea.	Water-As, Cd, Cu, Zn, Pb	45.00	45.00
	Laboratory #s 8803932-8803942		
	TOTAL AMOUNT DUE		\$025.00
	PLEASE RETURN DUPLICATE COPY OF INVOICE WITH REMITTANCE.		\$925.00

2815-2475-A30700



NORTH FORK AMERICAN FORK ABOVE BOG MINE Site # 1 UINTA NATIONAL FOREST

88 W 100 N

PROUG

UT 84603

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: NORTH FORK AMERICAN FORK ABOVE BOG MINE

Site ID:

Source: 00

Cost Code: 3508 Lab Number: 8803937 Type: 04

Date of Review and QA Validation Inorganic Review: 88/08/22

Sample Date: 88/07/20 Time: 10:25

Tot. Cations: 41
Tot. Anions: 83 me/l Cations:

2.4 Radiochemistry Review: 124 me/l Anions: 2.4 Microbiology Review:

Organic Keview:

Laboratory Analyses

d lotal:

Sulfate	29	mg/l	Tot, Alk.	91	mg/1
1. Hardns.	115.1	mg/1	TDS @ 180C	128	mg/1
l'-Arsenic	<1.0	ug/l	T-Barium	0.037	mg/1
1-Cadmium	<1	ug/l	T-Chromium	<5.0	ug/1
I-Copper	<20.0	ug/l	T-Iron	0.16	mg/1
I-Lead	(5.0	ug/l	I-Manganes	24.0	ug/1
Mercury	<0.2	ug/l	T-Selenium	(0.5	ug/1
1-Silver	<2.0	ug/l	1-Zinc	28.0	ug/l

UTAH STATE WATER QUALITY SYSTEM MONITORING RUN PROGRAM

Ju. 2700 003935

MC ORING RUN:[

1

STORET: [

1

SOURCE:[] COUNTY:[] USE:[] COST CODE:[]

DESCRIPTION: [North Fork American Fork above Bog Mine

COLLECTOR: [P][A][U][L][][H][][5][K][A][B][E][L][U][N][D][][][][]

DATE: [8] [8] [0] [7] [2] [0] YYMMDD

TIME:[1][0][2][5]

TYPE:[][]

FIELD TESTS

TEMPERATURE: pH: D.O.:

[1][0].[0] [][6].[7]

CO2: DEPTH: C1 RESID .: FLOW (MGD): SP.COND.: SP. GRAUITY: TRANSPARENCY:

[], [], [] [][][][][][[][][][], [] 10 10 10 10 FLOW (GPM): [][].[] FLOW (CFS):

0 10 10

SAMPLE BOTTLES NEEDED

TRY: [+) TOS, T-ALK, ARDNS, BOY

NUTRIENTS: [7]

TOTAL METALS: [#] AS, BA, CD, CR, CU, FE, PB, MN, 146, SE, AG, ZN

FIELD COMMENTS:

TAKE FLOW

Jump = .07' Width = 3.6' Depth = -3'

ADIT OF LOWER BOG MINE Site # 2 UINTA NATIONAL FOREST 88 W 100 N

PROVO UT 84603

> UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: ADIT OF LOWER BOG MINE

Site ID:

lot. Cations:

Anions:

Source: 00

me/1 Cations:

me/l Anions:

Cost Code: 3508

Lab Number: 8803945 Type: 04

Sample Date: 88/07/20 Yime: 11:35

Date of Keview and QA Validation Inorganic Review: 88/08/22

Organic Keview:

Radiochemistry Review: Microbiology Keview:

.d lotal: Laboratory Analyses

I-Arsenic T-Lead

3.0 ug/1 5.0 ug/1

T-Cadmium T-Zinc

控13mug/1 \$530.0 ug/1

NORTH FORK AMERICAN FORK BELOW LOWER BOG MINE Site # 3 UINTA NATIONAL FOREST 88 W 100 N UT 84603 · PROVO

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: NORTH FORK AMERICAN FORK BELOW LOWER BOG MINE

Site 10: Source: 00

Site ID: Source: 00
Cost Code: 3508
Lab Number: 8803933 Type: 04
Sample Date: 88/07/20 Time: 11:50

Tot. Cations: 35
Anions: 74 me/l Cations: Anions:

109 me/l Anions:

Date of Review and QA Validation Inorganic Review: 88/08/22 Organic Review:

2.0 Radiochemistry Review: 2.1 Microbiology Review:

Sulfate	31 1	ng/1	lot. Alk	73	mg/1
T. Hardns.	96.91	ng/I	TDS @ 180	C 120	mg/l
T-Arsenic	2.5 1	1/21	T-Cadmium	1	ug/1
T-Lead	(5.0)	ug/1	1-Zinc	77.0	ug/l

NORTH FORK AMERICAN FORK RIVER ABOVE PACIFIC Site # 34 UINTA NATIONAL FOREST 88 W 100 N . PROVO U1 84603

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: NORTH FORK AMERICAN FORK RIVER ABOVE PACIFIC

Site IU:

Source: 00

Cost Code:

350B

Lab Number: 8803939 Type: 04

Sample Date: 88/07/20 Time: 14:05

Date of Keview and QA Validation Inorganic Review: 88/08/22

Tot. Cations: lot. Anions:

nd Total:

42

80 me/l Cations:

122 me/l Anions:

Organic Keview: 2.5 Radiochemistry Review: 2.4 Microbiology Review:

1.0	- 11			F . A11	100	
18	mg/1			lot. Alk.	104	mg/1
120.9	mg/l			TUS @ 180C	130	mg/1,
1.0	ug/l			T-Barium	0.044	mg/1
<1	ug/1	1		T-Chromium	(5.0	ug/l
<20.0	ug/1			T-Iron	0.033	1/gm
(5.0	ug/1			1-Manganes	(5.0	ug/I
<0.2	ug/1			T-Selenium	(0.5	ug/1
(2.0	ug/l			1-Zinc	<20.0	ug/I
	120.9 1.0 <1 <20.0 <5.0 <0.2	18 mg/l 120.9 mg/l 1.0 ug/l <1 ug/l <20.0 ug/l <5.0 ug/l <0.2 ug/l <2.0 ug/l	120.9 mg/l 1.0 ug/l <1 ug/l <20.0 ug/l <5.0 ug/l <0.2 ug/l	120.9 mg/l 1.0 ug/l <1 ug/l <20.0 ug/l <5.0 ug/l <0.2 ug/l	120.9 mg/l 1DS @ 180C 1.0 ug/l T-Barium <1 ug/l T-Chromium <20.0 ug/l T-Iron <5.0 ug/l T-Manganes <0.2 ug/l T-Selenium	120.9 mg/l 1DS @ 180C 130 1.0 ug/l T-Barium 0.044 <1 ug/l T-Chromium <5.0 <20.0 ug/l T-Iron 0.033 <5.0 ug/l T-Manganes <5.0 <0.2 ug/l T-Selenium <0.5

UTAH STATE WATER QUALITY SYSTEM MONITORING RUN PROGRAM	4:27e.003939	
ORING RUN:[.]		
STORET:[] SOURCE:[] COUNTY:[] USE:[,]	COST CODE:[]	
DESCRIPTION: [Worth Fork American Fork River Above Pacific 1	Vine]	
COLLECTOR: [P][A][U][L][][A][][S][K][A][B][E][L][U][M][D][][10 10 1	
DATE: [8][8][0][7][2][0] TIME: [/][4][0][5] TYPE: [][1	
FIELD TESTS		
TEMPERATURE: [/][6].[0] CO2: pH: [][6].[7] DEPTH: [] D.O.: [][].[] C1 RESID.: SP.COND.: [][][][][][] FLOW (MGD): SP. GRAVITY: [].[][][] FLOW (GPM): [] TRANSPARENCY: [][][][] FLOW (CFS): [][][][]	[][][][] [][][].[] [][][].[] [][][][]	
SAMPLE BOTTLES NEEDED		
C STRY: [1] TDS, T-AUK, HROWS., SOY NUTRIENT: [7] TOTAL METALS: [1] AS, BA, CD, CR, CU, FE, PB, MN, HG,	SE, AG, ZN	
FIELD COMMENTS: TAKE FLOW Jump = .10' Width = 5.5'	Depth = .51	

PACIFIC MINE -MAIN ADIT Sile # 4 UINTA NATIONAL FOREST 88 W 100 N PROVO UT 84603

> UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PACIFIC MINE -MAIN ADIT

Site ID:

Cost Code: 3508 Lab Number: 8803947

Type: 04

Date of Review and QA Validation 88/08/22

Inorganic Review: Organic Keview: ®

Sample Date: 88/07/20 Time: Tot. Cations:

Anions:

me/1 Cations: me/1 Anions:

Source: 00

Radiochemistry Review: Microbiology Review:

Laboratory Analyses

'I-Arsenic 1-Copper Mercury

20.0 ug/1 442.0 .ug/1: <0.2 ug/1

T-Cadmium T-Lead T-Zinc

13 ug/1; 15.0 ug/1; 1600.0 ug/1

FIELD COMMENTS: TAKE FLOW Jump = 04' . width=1ft Depth = -15'

PACIFIC MINE N.W. PORTAL 5te #5 UINTA NATIONAL FOREST 88 W 100 N PROVO UT 84603

> UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PACIFIC MINE N.W. PORTAL

Site ID:

Source: 00

Cost Code:

350B

Lab Number: 8803943 Type:

04

Sample Date: 88/07/20 Time: 13:00

Date of Review and QA Validation

Inorganic Review: 88/08/22 Organic Review:

lot. Cations:

Lat. Anions: id Total:

me/l Cations: me/l Anions:

Radiochemistry Review: Microbiology Review:

Laboratory Analyses

T-Arsenic

2.0 ug/1

T-Lead

(5.0 ug/1

oi,	-	R-DATE/SAMPL	E-NUMBER.:[7	45 19 27 88 0 0 3 9	43
UTAH STATE WATER QUALITY SYS	STEM				
M ORING RUN:[)				
STORET:[] SOURCE:[] (COUNTY:[]	USE:[]	COST CODE:[]
DESCRIPTION: [PACIFIC MI	NE - N	JW PORTAL]	
COLLECTOR: [P][A][U][L][][H]	[][3][A	[A][B][E][L]	[][[][[][[]	[][][]	
DATE: [8][8][0][7][2][0]	TIME:[[0][0][1]	TYPE:[]	[]	
Y Y M M D D		FIELD TESTS			
pH: [][D.O.: [][SP.COND.: [][][][] SP. GRAVITY: [].[]].[]	CO2: DEPTH: C1 RESID.: FLOW (MGD): FLOW (GPM): FLOW (CFS):	С :		
		MPLE BOTTLES N			
CI STRY:[1] TOS, T-AC	LK - (CAN'T DO	FROM E	IN ACIDIFIC	D
TOTAL METALS: [4] AS , PB					
FIELD COMMENTS: TAKE FLOW	Sam	e as from	Mine Port	al at sample	<u>e</u>

PACIFIC MINE CENTER OF TAILINGS Site # 6 UINTA NATIONAL FOREST 88 W 100 N UT 84603 PROVO

> UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PACIFIC MINE CENTER OF TAILINGS

Site ID:

Source: 00

Cost Code: 350B

Lab Number: 8803944 Type:

04

Sample Date: 88/07/20 Time: 13:10

Inorganic Review:

Date of Keview and QA Validation 88/08/22

Organic Review:

Tot. Cations:

[at. Anions: d lotal:

me/1 Cations:

me/l Anions:

Radiochemistry Review: Microbiology Review:

Laboratory Analyses

T-Arsenic T-Copper I-Linc

13.0 ug/1 : 30.0 ug/1 1000.0 ug/1,

I'-Cadmium I-Lead

175.0 ug/1

PACIFIC MINE AT LOWER EDGE OF TAILINGS Sile #7 UINTA NATIONAL FOREST 88 W 100 N PROVO UT 84603

> UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PACIFIC MINE AT LOWER EDGE OF TAILINGS

Source: 00

Site 10:

3508

Cost Code:

Lab Number: 8803946 Type: 04

Sample Date: 88/07/20 Time: 14:25

lot. Cations:

Anions: d lotal:

me/l Cations: me/1 Anions:

Date of Keview and QA Validation Inorganic Review: 88/08/22

Organic Review:

Radiochemistry Review: Microbiology Review:

Laboratory Analyses

I-Arsenic 1-Copper Mercury

22.0 ug/1 30.0 ug/1 1 0.29 ug/1

T-Cadmium T-Lead

T-Zinc

9. ug/l . 850.0 ug/l · 1000.0 ug/1

51	R-DATE/SAMPI.	E-NUMBER.:[#7 .m 27000 39	46
UTAH STATE WATER QUALITY SYSTEM MONITORING RUN PROGRAM				
VITORING RUN:[]				
STORET:[] SOURCE:[] C	OUNTY:[]	USE:[]	COST CODE:[]
DESCRIPTION: [PACIFIC MINE AT LO	WEREDOE (OF TAILING	45]	
COLLECTOR:[P][A][U][L][][H][][5][X				
YYMMDD	FIELD TESTS	TYPE:[]	[]	
PH: [][6].[7] D.O.: [][][][] SP.COND.: [][][][][]	CO2: DEPTH: C1 RESID.: FLOW (MGD): FLOW (GPM): FLOW (CFS):			
SAM	IPLE BOTTLES N	EEDED		
MISTRY: [7] 705, FAUS - (CAW FAST	AKE DE	FROM AN	
I NUTRIENT: [7]	011/ 7.1			
TOTAL METALS: [1] AS, CD, CU, PE	5,179,00			
FIELD COMMENTS: TAKE FLOW Jump	=.03' Dept	h = .15' h); dth = 2.75	-
	NEX CIERT			

]

NORTH FORK AMERICAN RIVER BELOW PACIFIC MINE Sile # 8 UINTA NATIONAL FOREST 88 W 100 N PROVO UT 84603

> UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: NORTH FORK AMERICAN RIVER BELOW PACIFIC MINE

Site ID:

Source: 00

Cost Code:

350B

Lab Number: 8803934 Type: 04 Sample Date: 88/07/20 Time: 14:50 Date of Keview and QA Validation Inorganic Review: 88/08/22

Organic Keview:

Tot. Cations:

Anions: (d Total:

69 me/1 Cations: 69 me/l Anions:

Radiochemistry Review: 2.3 Microbiology Review:

Laboratory Analyses

lot. Alk. TDS @ 180C 134 mg/1 115 mg/1 1-Arsenic 4.5 ug/l T-Cadmium <1 ug/1 I-Copper <20.0 ug/1 I-Lead 20.0 ug/1 Mercury (0.2 ug/1 1-Zinc 81.0 ug/1

SAMPLE BOTTLES NEEDED

CISTRY:[1] 705, TALK

NUTRIENTS:[7]

TOTAL METALS:[] AS, CD, CU, PB, HG, ZN

FIELD COMMENTS:	TAKE FLOW Jump = .04	Depth = .5' Width = 6ft'
Site Located	Just upstream from M	outh of Dry Fork
\		. 0.

NORTH FORK AMERICAN FORK RIVER @ DUTCHMAN FLA Sile# 9 UINTA NATIONAL FOREST 88 W 100 N PROVO U1 84603

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: NORTH FORK AMERICAN FORK RIVER @ DUICHMAN FLA

Site ID: Source: 00

Cost Code: 3508 Lab Number: 88039 8803941 Type: 04

Sample Date: 88/07/20 Time: 15:35

Tot. Cations: 54

Date of Review and QA Validation Inorganic Review: 88/08/22

Organic Review:

Anions: 101 me/l Cations: 3.2 Radiochemistry Review: d Total: 155 me/l Anions: 3.2 Microbiology Review:

Sulfate	16	mg/1	Fot. Alk.	43 mg/1
1. Hardns.	157.3	mq/1	TDS @ 180C 1	74 mg/1
T-Arsenic	<1.0	ug/1	T-Barium 0.0	53 mg/1
T-Cadmium	<1	ug/1	T-Chromium <5	.0 ug/1
T-Copper	<20.0	ug/l	T-1ron 0.0	29 mg/1
T-Lead	5.0	ug/l	T-Manganes 7	.0 ug/1
Mercury	(0.2	ug/l	T-Selenium <0	.5 ug/1
T-Silver	(2.0	ug/l	1-Zinc 43	.0 ug/1

R-DATE/SAMPLE-NUMBER .: [#9 UTAH STATE WATER QUALITY SYSTEM Ju 2732003941 MONITORING RUN PROGRAM TORING RUN: [STORET:[] SOURCE:[] COUNTY:[] USE:[] COST CODE:[] DESCRIPTION: [North Fork American Fork River a Dutchman Flat] COLLECTOR: [P][A][U][L][][H][][S][K][A][B][G][L][U][M][D][][][][] DATE: [8] [8] [0] [7][2][0] TIME: [/][4][3][5] TYPE: [][] YYMMDD FIELD TESTS TEMPERATURE: [][][][][] [/][6].[5] CO2: [][6].[8] DEPTH: D.O.: [][].[] C1 RESID.: [].[][]

SP.COND.: [][][][][] FLOW (MGD): [].[][]

SP. GRAVITY: [].[][][] FLOW (GPM): [][][][][][]

TRANSPARENCY: [][][].[] SAMPLE BOTTLES NEEDED STRY: [4] TOS, T. ALR, HRDNIS, SOY W; HG, SE, AG, Z. TOTAL METALS:[1] AS, BA, CD, CA, CU, PE, PB, MAN, TAKE FLOW Jamp = . 15' Width = 8' Depth = . 67' FIELD COMMENTS:

MARY ELLEN GULCH CREEK AT MOUTH Sile # 10 UINTA NATIONAL FOREST 88 W 100 N

PROVO UT 84603

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: MARY ELLEN GULCH CREEK AT MOUTH

Site 10:

Cost Code: 350B Lab Number: 8803940 Type: 04 Sample Date: 88/07/20 Time: 16:10 Date of Keview and QA Validation 88/08/22 Inorganic Review:

Tot. Cations: Organic Review: 48

Source: 00

Tot. Anions: 98 me/l Cations: 146 me/l Anions: 2.8 Radiochemistry Review: 146 me/l Anions: 2.9 Microbiology Review:

33	mg/1	Tot, Alk.	109	mg/l
135.8	mg/1	TDS @ 180C	154	mg/1
2.5	ug/l	l'-Barium	0.046	mg/1
<1	ug/l	1-Chromium	(5.0	ug/l
<20.0	ug/l	T-Iron	0.059	mg/1
(5.0	ug/1	1-Manganes	6.0	ug/1
(0.2	ug/l	T-Selenium	(0.5	ug/1
(2.0	ug/1	1-Zinc	72.0	ug/1
	135.8 2.5 <1 <20.0 <5.0 <0.2	33 mg/1 135.8 mg/1 2.5 ug/1 <1 ug/1 <20.0 ug/1 <5.0 ug/1 <0.2 ug/1 <2.0 ug/1	135.8 mg/1 TDS @ 180C 2.5 ug/1 Y-Barium <1 ug/1 T-Chromium <20.0 ug/1 Y-Iron <5.0 ug/1 T-Manganes <0.2 ug/1 Y-Selenium	135.8 mg/l TDS @ 180C 154 2.5 ug/l Y-Barium 0.046 <1 ug/l Y-Chromium <5.0 <20.0 ug/l Y-Iron 0.059 <5.0 ug/l Y-Manganes 6.0 <0.2 ug/l Y-Selenium <0.5

J

NORTH FORK AMERICAN FORK BELOW MARY ELLEN GUL Site # 11 UINTA NATIONAL FOREST 88 W 100 N PROVO UT 84603

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: NORTH FORK AMERICAN FORK BELOW MARY ELLEN GUL

Site ID:

Source: 00

Cost Code: 350B Lab Number: 8803936

Lab Number: 8803936 Type: 04 <u>Date of Review and QA Validation</u>
Sample Date: 88/07/20 Time: 16:30 Inorganic Review: 88/08/22
Tot. Cations: 52 Organic Review:

Sample Date: 50707.

Tot. Cations: 52

Anions: 101 me/l Cations: 3.1 Radiochemistry Neview: 3.1 Microbiology Review: 3.1 Radiochemistry Review:

Sulfate	20	mq/l	Tot. Alk.	135	mq/1
1. Hardns.	150.7		TDS @ 180C		mg/1
l'-Arsenic		ug/1	T-Barium	0.053	mg/1
1-Cadmium	<1	ug/1	1-Chromium	(5.0	ug/1
I-Copper	<20.0	ug/l	1-1ron	0.039	mg/1
T-Lead	10.0	ug/l	I-Manganes	7.0	ug/l
Mercury	(0.2	ug/l	T-Selenium	<0.5	ug/1
T-Silver	(2.0	ug/l	1-Zinc	40.0	ug/1

R-DATE/SAMPLE-NUMBER . : [# // 01 " -UTAH STATE WATER QUALITY SYSTEM Jet 2785003936 MONITORING RUN PROGRAM TORING RUN:[STORET:[] SOURCE:[] COUNTY:[] USE:[] COST CODE:[] DESCRIPTION: [North Fork American Fork below-Mary Ellen Gulch] COLLECTOR: [P][A][U][L][][H][][5][K][A][B][E][L][U][N][D][][][][] DATE: [8][8][0][7][2][0] TIME:[1][6][3][0] TYPE:[][] YYMMDD FIELD TESTS TEMPERATURE: [1][7].[0] CO2: pH: DEPTH: [][6].[9] D.O.: [][][] Cl RESID.: [][][][][][] SP. COND.: [] [] [] FLOW (MGD): FLOW (MGD): SP. GRAVITY: [][][] FLOW (GPM): [][][][].[] [][].[] FLOW (CFS): [][][][][][][] TRANSPARENCY: SAMPLE BOTTLES NEEDED CH STRY: [] TDS, T-ALIK, SOY-HADNS NUTRIENT: [7] TOTAL METALS: [\$] AS, BA, CD, CR, OC, FE, PB, NKN, 46, SE, AG, ZN TAKE FLOW Jump = .03' Depth = .625' Width =7' FIELD COMMENTS:

MARY ELLEN GULCH CREEK ABOVE MINE TAILINGS 5-12-412 UINTA NATIONAL FOREST 88 W 100 N

UT 84603

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: MARY ELLEN GULCH CREEK ABOVE MINE TAILINGS

Site IU:

Source: 00

PROVO

Lab Number: 88020

Lab Number: 8803938 Type: 04 Date of Review and QA Validation
Sample Date: 88/07/21 Time: 09:40 Inorganic Review: 88/08/22
Tot. Cations: 39 Organic Review:

Tot. Cations: 39 Organic Review:
Tot. Anions: 77 me/l Cations: 2.3 Radiochemistry Review:
nd Total: 116 me/l Anions: 2.3 Microbiology Review:

Sulfate	22	mg/1	lot. Alk.	92	mg/1
1. Hardns.	110.1	mg/l	TDS @ 180C	120	mg/1
T-Arsenic -	1.5	ug/l	f-Barium	0.041	mg/1
1-Cadmium	<1	ug/1	T-Chronium	(5.0	ug/1
T-Copper	<20.0	ug/1	T-Iron	0.025	mg/1
1-Lead	<5.0	ug/l	T-Manganes	(5.0	ug/1
Mercury	<0.2	ug/l	Y-Selenium	(0.5	ug/1
T-Silver	<2.0	ug/1	T-Zinc	(20.0	ug/1

01 " •	R-DATE/SAM	MPLE-NUMBER . : E#	12	
UTAH STATE WATER QUALITY SYSTEM MONITORING RUN PROGRAM			1.17 0039	38
MCORING RUN:[,]				
STORET:[] SOURCE:[]	COUNTY:[]	USE:[.]	COST CODE:[J
DESCRIPTION: [MARY ELLEN GULCH	CREEK	ABOUG MINE T	TAILINGS	
COLLECTOR: [P][A][V][V][][H][][5][k][A][B][E][()[v][N][D][]	[][][]	
DATE: [8][8][0][7][2][/] TIME: Y Y M M D D	[0][9][4][0] FIELD TESTS		[]	
	FIELD TESTS			
	CO2: DEPTH: Cl RESID.: FLOW (MGD) FLOW (GPM) FLOW (CFS)	:		
SA	MPLE BOTTLES	S NEEDED		
CH STRY: [4] TDS, T. ALIC, HR	* /			
TOTAL METALS: [1] AS, BA, CD, CR	CU, FE,	, PB, MN, I	19,5E,AG,Z	N
	,			
FIELD COMMENTS: TAKE FLOW Jump	, =.02'	width = 1.51	Depth = . 1	31
			THE STATE OF	

MARY ELLEN MINE PORTAL Sile # 13 UINTA NATIONAL FOREST

88 W 100 N

PROVO

U1 84603

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: MARY ELLEN MINE PORTAL

Site 10: Cost Code:

350B

Lab Number: 8803942 Type: 04

Sample Date: 88/07/21 Time: 10:00

Source: 00

Date of Review and QA Validation Inorganic Review: 88/08/22

Organic Review:

Tot. Cations: me/l Cations: me/l Anions: Tot. Anions: nd Total:

Radiochemistry Review: Microbiology Review:

Laboratory Analyses

97.0 ug/l T-Arsenic 1-Copper <20.0 ug/l 570.0 ug/1 T-Zinc

T-Cadmium T-Lead 1 ug/1 <5.0 ug/1

	R-DATE/SAME	LE-NUMBER . : [4	#13	
or			#13 77000394	12
UTAH STATE WATER QUALITY SYSTEM MONITORING RUN PROGRAM				
MOTORING RUN:[]				
SIGKET:[] SOURCE:[]	COUNTY:[] .	USE:[]	COST CODE:[)
DESCRIPTION: [MARY ELEN	MINE POR	TAL	1	
COLLECTOR: [P][A][V][L][][H][][5][A][A][B][G][L][v][N][Þ][]	[][][]	
DATE: [8][8][0][7][2][1] TIME:	[1][0][0][1]	TYPE:[]	[]	
1 1 4 4 5 5	FIELD TESTS			
PH: [][6].[5] D.O.: [][].[] SP.COND.: [][][][][] SP. GRAVITY: [].[][][]	CO2: DEPTH: C1 RESID.: FLOW (MGD): FLOW (GPM): FLOW (CFS):	C	[][][][][][][][][][][][][][
Sf	MPLE BOTTLES	NEEDED		
CH STRY: [1] TOS, T-ALK, HANNORMENT: [7] TOTAL METALS: [1] AS,CD,CU,PB, 2	NONES), BE SI	4-CAN'T I ACTOLFIE	DO FROM A D BUHLE	٨
FIELD COMMENTS: TAKE FLOW Jump	=.03' Wi	dth = 1.0'	Dep th = .21'	
		WAY THE WAY IN THE		

WEST FORK MARY ELLEN GULCH CREEK & MOUTH Sile # 13 A UINTA NATIONAL FOREST 88 W 100 N PROVO UT 84603

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: WEST FORK MARY ELLEN GULCH CREEK @ MOUTH

Site 10:

Source: 00

Site 1D: Source: 00

Cost Code: 3508

Lab Number: 8803932 Type: 04

Sample Date: 88/07/21 Time: 11:00

Date of Review and QA Validation Inorganic Review: 88/08/22
Organic Review:
2.5 Radiochemistry "oview:

Tot. Cations: 43
Tot Anions: 103 me/l Cations: 146 me/l Anions:

2.6 Microbiology Review:

Sulfate	67	mq/l	Tot. Alk. 60	mq/1
1. Hardns.	121.7			mq/1
T-Arsenic	14.5	ug/1	1-Barium 0.022	mg/1
1-Cadmium	2	ug/l	1-Chromium <5.0	ug/1
T-Copper	53.0	ug/l	Y-Iron 1.2	mg/1
T-Lead	10.0	ug/1	T-Manganes 74.0	ug/1
Mercury	(0.2	ug/l	Y-Selenium <0.5	ug/1
T-Silver	<2.0	ug/l	T-Zinc 450.0	ug/1

MARY ELLEN GULCH CREEK BELOW MINE TAILINGS UINTA NATIONAL FOREST

Sile #14

N 001 M 88

PROVO

UT 84603

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: MARY ELLEN GULCH CREEK BELOW MINE TAILINGS

Site 10:

Source: 00

Cost Code:

350B

Lab Number: 8803935 Type: 04 Sample Date: 88/07/21 Time: 11:35

Date of Keview and QA Validation

Inorganic Review: 88/08/22

lot. Cations:

48

98 me/l Cations:

Organic Keview: 2.8 Radiochemistry Review:

Tot Anions: G | Total:

146 me/l Anions:

2.8 Microbiology Review:

Sulfate	38	mg/1	Tot. Alk.	100	mg/1
1. Hardns.	137.4	mg/1	TDS @ 180C	138	mg/1
1-Arsenic	3.0	ug/1	Y-Cadmium	<1	ug/l
1-Copper	(20.0	ug/1	T-Lead	(5.0	uq/1
T-/inc	110 0	110/1			10,000

PACIFIC MINE MAIN PORTAL AT ADIT UINTA NATIONAL FOREST 88 W 100 N U1 84603 PROVO

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PACIFIC MINE MAIN PORTAL AT ADIT

Site 10:

Source: 00

Cost Code:

350B

Lab Number: 8802854 Type: 04

Sample Date: 88/05/18 Fime: 10:00

Date of Keview and QA Validation

Inorganic Review:

Organic Review:

Tot. Cations:

Tot. Anions: nd lotal: me/l Cations:

me/1 Anions:

Radiochemistry Review: Microbiology Review:

Laboratory Analyses

Tot. Alk.	163	mg/1	TDS @ 180C	202 m	ng/1
1-Arsenic	22.0	ug/l	1-Barium	0.069 n	ng/1
I'-Cadmium	6	ug/l	I-Chromium	(5.0 t	ug/1
1-Copper	34.0	ug/l	1-1ron	4.0 n	ng/1
I'-Lead	25.0	ug/l	l'-Manganes	11.0 0	ug/1
Mercury	0.2	ug/1	1-Selenium	(0.5 L	1/2
I-Silver	(2.0)	ug/l	1-Zinc	800.0 u	1/2

UINTA NATIONAL FOREST

SEP 2 6 1988

5	TI	CUP	-
853		PIO	-
40		CS	-
ENG		APE AND	-
645		878/1699	+
MIN		FIRE ISTAL	E
Y: folgra. et	1	Filterstra	-
13-1	1_	700	-
D-3	_	-	-
D-3		-	-

xact Description (of Sampling Point	AD!	THE CASE WAS ARREST TO THE	NG	MAIN TO	oR Iric
ollector: Phu					County:	Cost Code: 35
end Report to:	linta N	AT.	FORCS +			
		-	VO, UTAH			zip: <u>8460</u>
/_tcpc	<u>/</u> TMPM	800	NutB	act	_PestTHM	RadSp
BOD	Cyan	ide			MPN Total Colifor	rmc /100 ml
TSS		olics		790	MPN Fecal Colifor	
TKN	Sulf	ide		7.2	MF Total Coliforn	ns/100 ml
TOC	pH				MF Fecal Coliforn	ns/100 m1
000	0i1 &	& Grease			Fecal Streptococo	
				-	Plate Count - Org]./ml
CATIONS		. 4	ANIONS		TOTAL	METALS
			Disabanda		*1	X Lead
Armonia Arsenic	Lead Magnesium	-	Bicarbonate Carbon Dioxide		Aluminum Arsenic	X Manganes
Barium	Manganese		Carbonate		× Barium	× Mercury
Boron	Nickel		Chloride		Beryllium	Molybden
Cadmium	Potassium	AMES AND	CO3 Solids		X Cadmium	Nickel
Calcium	Selenium		Fluoride		X Chromium	✓ Selenium
Chromium	Silver		Hydroxide	- 0	Cobalt	λ Silver
Chromium, Hex	Sodium		Nitrate		∠ Copper	Vanadium
Copper	Zinc		Nitrite		lron	_≺_Zinc
_lron			Phosphorus, Ortho	0 .		
			Silica			
			Sulfate	* 07		
	1			•		
Total I	Phosphorus				RADIOLOGICS	
	Alk. as Catto				TO TO LOUI US	
	lardness as CaCO	,		Alpl	na, Gross	228 _{Radium}
	ity as NTU			Beta	a, Gross	Uranium
	nd. (umhos/cm)			226	Radium	
X TDS @ 1						
Other:						

NORTH FORK AMERICAN RIVER AT DUTCHMAN FLAT UINTA NATIONAL FOREST

88 W 100 N

PROVO

U1 84603

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: NORTH FORK AMERICAN RIVER AT DUTCHMAN FLAT

Site 1D:

Source: 00

Cost Code: 3508 Lab Number: 8802855 Type: 04

Date of Keview and QA Validation

Inorganic Review: Organic Keview:

Sample Date: 88/05/18 fime: 16:10 lot. Cations:

lot. Anions: nd lotal:

me/l Cations:

me/1 Anions:

Radiochemistry Review: Microbiology Keview:

lot, Alk.	83	mg/1	TDS @ 180C	102	mg/1
1-Arsenic	2.5	ug/l	1-Barium	0.056	mg/1
1-Cadmium	<1	ug/1	I-Chromium	(5.0	ug/1
1-Copper	<20.0	ug/1	1-lron	0.45	mg/1
I-Lead	60.0	ug/1	l'-Manganes	31.0	ug/1
Mercury	<0.2	ug/1	1-Selenium	<0.5	ug/1
I-Silver	<2.0	ug/1	I-∠inc	77.0	ug/1

1-/1 02855

Water System No	Source	No Date C		5-18 Time Co	11ected /6. /C
	of Sampling Point:	LOLU E			
					TOTCK
· K	rIVER At	DitcHM	IAN	FLAT	PEREE
ollector: PAL	L H. 5K,	ABELUNI)	County:	_cost code: 35
end Report to:	linta NA	AT. FORE:	って	Te1ephone	No: 377-57
idress: 680	₩ /00 N	Provon	ITAH		
<u>/</u> тсес	/THPH	_BODRut	Bact	PestTHM	RadSpe
BODTSS	Cyanide Phenolic		-	MPN Total Colifor	
TKN	Sulfide			MF Total Coliforn	
TOC	pH			MF Fecal Coliforn	
C00	0i1 & Gr	ease .		Fecal Streptococc	
				Plate Count - Org	J./MI
CATIONS		ANIONS		TOTAL	METALS
Ammonia	Lead .	Bicarbonate		Aluminum	X_Lead
Arsenic	Magnesium .	Carbon Dioxi	de .	✓ Arsenic	<u>X</u> Manganese
Barium	Manganese .	Carbonate		⊻Barium	Mercury
Boron	Nickel .	Chloride		Beryllium	Molybden
Cadmium	Potassium .	CO3 Solids		K_Cadmium	Nickel
Calcium _	Selenium .	Fluoride		Chromium	Selenium
Chromium _	Silver .	Hydroxide		Cobalt	
Chromium,Hex	Sodium .	Nitrate		X Copper	Vanadium
Copper	Zinc .	Nitrite			X_Zinc
_lron		Phosphorus,(ortho .		
		Silica			
	٠.	Sulfate			
	Phosphorus			RADIOLOGICS	
	Nlk. as CaCO3				200
	lardness as CaCO3			a, Gross	_228Radium
AND REAL PROPERTY AND ADDRESS OF THE PARTY AND	ity as NTU		Beta	, Gross	_Uranium
	nd. (umhos/cm)		ZZ6R	adium	
Other:_					

PAC.MINE NW PORTAL PIPED OUT OF MINE UINTA NATIONAL FOREST 88 W 100 N PROVO UI 84603

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PAC.MINE NW PORTAL PIPED OUT OF MINE

Site 10:

Source: 00

Cost Code: 3508

Lab Number: 8802856 Type: 04

Sample Date: 88/05/18 Fime: 10:10

Date of Review and QA Validation Inorganic Review:

Organic Keview:

lot. Cations: Tot. Anions:

me/1 Cations:

me/1 Anions:

Radiochemistry Review: Microbiology Review:

Fot. Alk.	198	mg/1	TDS @ 180C	208	mg/1
1-Arsenic	1.0	ug/1	1-Barium	0.15	mg/l
I-Cadmium	<1	ug/1	l'-Chromium	(5.0	ug/1
1-Copper	<20.0	ug/1	1-1ron	0.091	mg/1
I'-Lead	60.0	ug/1	l'-Manganes	19.0	ug/1
Mercury	(0.2	ug/1	1-Selenium	<0.5	ug/1
I-Silver	(2.0	uq/1	I-Zinc	78.0	ug/1

ter System No	Source No	Date Collect	ed 88 5 /8 Time	Collected 10:10
	77	AC. MINE	NW Pa	011
act Description o	of Sampling Point:	1C. PIINE	Nº 10	NETHE
	PIDED (out at a	1/NE	
11ector: PAC	OL H. 51	KABELIUND	County:	Cost Code: 350
nd Report to:	Vinita ALA-	NONAL FOR		one No: 377-57
				8
dress: 8 C	w-100 N	1 PROV	10, UTAH	zip: <u>8460</u>
/	1			
PCPC _	/TMPH	BODNutBac	tPestT	HMRadSpec
B00	Cyanide		MPN Total Col	iforms/100 ml
TSS	Phenolics		MPN Fecal Col	iforms/100 ml
TKN	Sulfide		MF Total Coli	forms/100 ml
TOC	pH		MF Fecal Coli	
000	Oil & Grea	se .	Fecal Strepto	
			Plate Count -	Org./ml
CATIONS		ANIONS	. TO	TAL METALS
				,
_Ammonia _	Lead .	Bicarbonate	Aluminu	m
Arsenic	Magnesium .	Carbon Dioxide	Arsenic	
Barium	Manganese .	Carbonate	Barium	<u></u> ✓ Hercury
Boron	Nickel .	Chloride	Berylli	[2] [1] [1] [2] [2] [3] [3] [4] [4] [4] [4] [4] [4] [4] [4] [4] [4
Cadmium	Potassium .	CO3 Solids	Cadmium	
_Calcium _	Selenium .	Fluoride		
Chromium	Silver .	Hydroxide	Cobalt	<u>x</u> Silver
Chromium,Hex	Sodium .	Nitrate	Copper	Vanadium
_Copper _	Zinc .	Nitrite	lron	Zinc
_lron		Phosphorus,Ortho		
		Sulfate		
	, , , , , , , , , , , , , , , , , , ,	Surrace		
	THE THE BOTTON OF THE			
	hosphorus		RADIOLO	GICS
	llk. as CaCO3			220
Total H	lardness as CaCO3		Alpha, Gross	228 _{Radium}
	tu ac MTH		Beta, Gross	Uranium
Turbidi			226	
Turbidi	id. (umhos/cm)		226Radium	

PORTAL LOWER BOG MINE UINTA NATIONAL FOREST 88 W 100 N UI 84603 PROVO

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PORTAL LOWER BOG MINE Site 10: Source: 00

3508 Cost Code:

Lab Number: 8802857 Type: 04 Sample Date: 88/05/18 Time: 12:30

lot. Cations:

lot. Anions: ond lotal:

me/1 Cations: me/1 Anions: Date of Keview and QA Validation

Inorganic Review: Organic Keview:

Radiochemistry Review: Microbiology Review:

Tot. Alk.	0	mg/1	TDS @ 180C	90	mg/1
1-Arsenic	1.5	ug/l	1-Barium	0.037	mg/1
I-Cadmium	12	ug/1	l'-Chromium	(5.0	ug/1
1-Copper	<20.0	ug/1	1-1ron	7.9	mg/1
I-Lead	(5.0	ug/l	I-Manganes	270.0	ug/1
Mercury	<0.2	ug/1	1-Selenium	(0.5	ug/l
I-Silver	(2.0	ug/l	l'-∠inc	510.0	ug/1

2857

	Source No.		yy/mm/dd		24 hr	cloc
xact Description	of Sampling Point:	ORTAC 102	JER 1	20(7/	Wille	4
	L H. SKAB					
end Report to: (JINTA NAT	TONAL FOR	2657	Te1ephone	No: 377-	57
	W 100N					
<u>/</u> TCPC		300NutBa	actPest	THM	Rad	Spe
BOD	Cyanide		MPN To	otal Colifor	ms/100 m1	
TSS	Phenolics		-	ecal Colifor		
TKN	Sulfide		MF Tot	al Coliform	s/100 ml	
TOC	pH		The state of the s	cal Coliform		
000	Oil & Great	ie .		Streptococc		
			Plate	Count - Org	./m1	
					Salayet S	
CATIONS		ANIONS		TOTAL	METALS	
Ammonia	Lead .	Bicarbonate		Aluminum	XLead	
Arsenic	Magnesium .	Carbon Dioxide	. X	Arsenic	X Manga	nese
Barium	Manganese .	Carbonate	. X	Barium	入 Mercu	гу
Boron	Nickel .	Chloride		Beryllium	Molyb	deni
Cadmium	Potassium .	003 Solids	· <u>Y</u>	Cadmium	Nicke	
Calcium	Selenium .	Fluoride	Account to	Chromium	× Selen	
Chromium	Silver .	Hydroxide		Cobalt	Silve	
Chromium,Hex	Sodium .	Nitrate		Copper	Vanad	ium
Copper	Zinc .	Nitrite		Iron	^_Zinc	
lron		Phosphorus,Ortho				
		Silica				
		Sulfate				
Total I	Phosphorus			RADIOLOGICS		
	llk. as CaCO ₃			INDIOLOGICS		
	lardness as CaCO3		Alpha, Gro	oss	228 _{Radium}	
	ity as NTU		Beta, Gros	The second second	Uranium	
	nd. (umhos/cm)		226 _{Radium}			
X TDS @ 1						
Other:		The state of the s				

MARY ELLEN PORTAL UINTA NATIONAL FOREST 88 W 100 N PROVO UI 84603

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: MARY ELLEN PORTAL

Source: 00

Site 10: Cost Code:

350B

Lab Number: 8802858 Type: 04

Date of Keview and QA Validation

Sample Date: 88/05/16 Fime: 15:00

Inorganic Review:

lot. Cations:

me/l Cations:

Organic Keview: Radiochemistry Review:

Tot. Anions: nd lotal:

me/l Anions:

Microbiology Keview:

Tot. Alk.	36	mg/1	TDS @ 180C	206	mg/1
1-Arsenic	100.0	ug/1	1-Barium	0.019	mg/1
I'-Cadmium	4	ug/l	1-Chromium	(5.0	ug/1
1-Copper	40.0	ug/1	1-1ron	9.9	mg/1
I-Lead	10.0	ug/1	l'-Manganes	140.0	ug/1
Mercury	<0.2	ug/1	1-Selenium	(0.5	ug/1
I-Silver	(2.0	uq/1	I-∠inc	1200.0	uq/1

1.7-1-1-2858

Water System No	Source No.	Date Collect		Collected 15.0
	A1	ARY ELLE	yy/mm/dd	
xact Description	of Sampling Point: / /	May GLLE	NICHOTA	<u> </u>
		All and the second		
ollector: PAV	L H. SKAR	BELLOND	County:	Cost Code: 350
end Report to: U	ININ NATI	ONAL FORE		
		PROVO		
idress: (· () (S FOOTA	TROVE	10.1111	
/TCPC .	PMB	000NutBac	et Pest T	HMRadSpe
BOO	Cyanide		MPN Total Col	i forms /100 ml
TSS	Phenolics		MPN Fecal Col	
TKN	Sulfide		MF Total Coli	
TOC	pH		MF Fecal Coli	forms/100 m1
COD	Oil & Greas	se .	Fecal Strepto	cocci/100 ml
			Plate Count -	Org./ml
CATIONS		ANIONS	· <u>10</u>	TAL METALS
Ammonia	Lead	Bicarbonate	Aluminu	n / Lead
Arsenic	Magnesium .	Carbon Dioxide	. Arsenic	\ Manganese
Barium	Manganese .	Carbonate	. X Barium	\ Mercury
Boron	Nickel .	Chloride	. Berylli	
Cadmium	Potassium .	CO3 Solids	. A' Cadmium	
Calcium	Selenium .	Fluoride	. Y Chromiu	n . Y Selenium
Chromium	Silver .	Hydroxide	. Cobalt	Silver
Chromium, Hex	Sodium .	Nitrate		Vanadium
Copper	Zinc .	Mitrite	lron	
lron		Phosphorus,Ortho		
		Silica		
		Sulfate		
Total f	hosphorus		RADIOLO	ctos
	ilk. as CaCO ₃		IONUTULU	aroj
	lardness as CaCO ₂		Alpha, Gross	228 _{Radium}
The second secon	ty as NTU		Beta, Gross	Uranium
	id. (umhos/cm)		226 _{Radium}	
X TDS @ 1	80°C			

PACIFIC PORTAL AT CREEK (MARKINGS WIPED OFF UINTA NATIONAL FOREST 88 W 100 N U1 84603 PROVO 377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PACIFIC PORTAL AT CREEK (MARKINGS WIPED OFF

Site 10:

Source: 00

Cost Code:

350B

Lab Number: 8802859

Type: 04 Time:

Date of Keview and QA Validation

Inorganic Review: Organic Keview:

Sample Date: Tot. Cations:

lot. Anions: nd lotal: me/l Cations: me/l Anions:

Radiochemistry Review: Microbiology Keview:

Tot. Alk.	164	mg/1	TDS @ 180C	200	mg/1
1-Arsenic	22.5	ug/l	1-Barium	0.28	mg/1
f-Cadmium	31	ug/1	I-Chromium	(5.0	ug/1
1-Copper	60.0	ug/l	1-1ron	5.3	mg/1
I'-Lead	4000.0	ug/l	l'-Manganes	23.0	ug/1
Mercury	0.63	ug/I	1-Selenium	<0.5	ug/1
T-Silver	5.0	ug/l	l'-∠inc	1600.0	ug/1

Water System No	Source No	Date Collec	ted 85-5- Tin	ne Collected		
			yy/mm/dd	24 hr clo		
Exact Description o	of Sampling Point: PAC	Tic	PORTAL .	17 CK'EEK		
collector: PA	UL H. SKNS	BECUMO	County:	Cost Code: 35		
11:	WHA NATIONAL	C-21		377-5		
end Report to:	WH MATIONAL	- TOIC ()	Telep	phone No: 3//		
iddress: 650	W HOON	Micera	1,011/	1 zip: 6.46.		
800	Cyanide		MPN Total Co	oliforms/100 ml		
TSS	Phenolics		MPN Fecal Co	oliforms/100 ml		
TKN	Sulfide			iforms/100 ml		
TOC	pH		MF Fecal Coliforms/100 ml			
000	Oil & Grease			Fecal Streptococci/100 ml		
			Plate Count	- Org./ml		
CATIONS		ANIONS	: 1	TOTAL METALS		
				1		
Ammonia	Lead	Bicarbonate	Alumin			
Arsenic Barium	Magnesium Manganese .	Carbon Dioxide Carbonate	/_Arseni			
Boron	Nickel .	Chloride	. Beryll			
Cadnium	Potassium .	CO3 Solids	. Cadmiu	The second secon		
Calcium	Selenium .	Fluoride	. A Chromi			
Chromium	Silver	Hydroxide	Cobalt			
Chromium, Hex	Sodium	Nitrate	. Copper	Control of the Contro		
Copper	Zinc .	Nitrite	. Iron	✓. Zinc		
lron		Phosphorus, Ortho				
		Silica				
		Sulfate				
Total 9	Phosphorus		RADIOL	ogics		
	ilk. as CaCO ₃		The same of the sa			
	lardness as CaCO ₃		Alpha, Gross	228 _{Radium}		
	ty as NTU		Beta, Gross	Uranium		
Sp. Con	nd. (umhos/cm)		226Radium			
	80°C					
Other:						

PACIFIC N TAILING UINIA NATIONAL FOREST 88 W 100 N PROVO U1 84603

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PACIFIC N TAILING

Site 10: Source: 00

350B Cost Code:

Lab Number: 8802860 Type: 04

Sample Date: 88/05/18 Time:

lot. Cations:

lot. Anions:

me/l Cations: and lotal: me/l Anions:

Date of Keview and QA Validation

Inorganic Review: Organic Keview:

Radiochemistry Review: Microbiology Review:

lot, Alk.	21	mq/1	TDS @ 180C	140	mq/1
1-Arsenic		ug/l	1-Barium		mg/1
I-Cadmium		ug/1	l'-Chromium		ug/1
1-Copper	260.0		1-1ron	13.0	mg/1
I'-Lead	20000.0	ug/1	l'-Manganes	48.0	ug/1
Mercury	3.24	ug/1	1-Selenium	1.0	ug/1
I-Silver	45.0	ug/1	l'-Zinc	7700.0	ug/1

....2660

UTAH STATE HEALTH LABORATORY

Water System No	Sc	ource No	Date Colle	cted Pr	5/8. Time Co	
				уу	/mm/dd	24 hr cloc
xact Description	of Sampling Poi	nt: TA	COFIC A	!	TAILING	
ntike is						
ollector: PAC	ic H.	SKAB	ELUNIA		County:	_Cost Code: 35
end Report to: (livita	NIAT	lowner F	C.20:) Telephone	No: 377-5
89	(3/- (1)	0.1	PRO	103	LITALL	. 00/
idress:	10 10	0.0	1 10 01	1		Zip:/L.

<u>l</u> тс	PH	800	NutB	act _	PestTHM	RadSpe
		, Ju	1,17			
800		nide nolics		_	MPN Total Colifor MPN Fecal Colifor	
TSS		fide			MF Total Colifor	
TOC	DH DH	riue		1/3	MF Fecal Coliforn	
COD	-	& Grease		1	Fecal Streptococo	
			Kill International	100	Plate Count - Or	
			•			
CATIONS			ANIONS		TOTAL	METALS
165			mit-		Aluminum	× Lead
Ammonia	Lead		Bicarbonate Carbon Dioxide		Arsenic	- Lead - Manganese
Arsenic Barium	Magnesium Manganese		Carbonate		∠ Arsenic	Hercury
Boron	Nickel	-	Chloride	1	Beryllium	Molybdenu
Cadmium	Potassium		_CO3 Solids .		× Cadmium	Nickel
Calcium	Selenium		Fluoride		Chromium	< Selenium
Chromium	Silver	The second second	Hydroxide		Cobalt	< Silver
Chromium, Hex	Sodium		Nitrate		Copper	Vanadium
Copper	Zinc		Nitrite		- Iron	×Zinc
lron			Phosphorus, Ortho	0 .		
			Silica			
			Sulfate			
	1				2001	
Total	Phosphorus				RADIOLOGICS	
	Alk. as CaCO ₃					
	Hardness as CaC	02		Alpi	na, Gross	228 _{Radium}
	ity as NTU	3			a, Gross	Uranium
-	nd. (umhos/cm)				Radium	
XIDS @						
Other:						

MARY ELLEN CREEK 1/4 MILE BELOW MINE AREA UINTA NATIONAL FOREST 88 W 100 N 377-5780 UT 84603 PROVO

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: MARY ELLEN CREEK 1/4 MILE BELOW MINE AREA

Source: 00

Site 1D: Source: 00 Cost Code: 350B Lab Number: 8802861 Type: 04 Sample Date: 88/05/18 Time:

Date of Review and QA Validation Inorganic Review: 88/06/22 Organic Review: Radiochemistry Review:

lot. Cations:

fot. Anions: 55 me/l Cations: 55 me/l Anions:

1.8 Microbiology Review:

fot. Alk.	92	mg/1	FDS @ 180C	132	mg/L
1-Arsenic	<1.0	ug/l	1-Barium	0.039	mg/1
T-Cadmium	2	ug/l	1-Chromium	(5.0	ug/1
T-Copper	42.0	ug/l	T-Iron	1.1	mg/l
T-Lead	40.0	ug/l	i-Manganes	46.0	ug/l
Mercury	(0.2	ug/l	1-Selenium	<0.5	ug/l
I-Silver	<2.0	ug/l	ľ-Zinc	310.0	ug/l

Ja7ta.0u2861

Water System No	So	urce No	Date Collec			
		4.7			mm/dd	24 hr clos
xact Description	of Sampling Poi	nt: /1/1)[RY ELLER	VI (CREEK	
	1/4	MI C-	BELOW	M	. 1/: 1	DC-A
0						
						Cost Code: 35
end Report to:_(), wha	NATIO	UNAL FO	TZG5	/ Telepho	ne No: 377-57
iddress:	8 W	100 1	J PROY	0,0	71111	Zip:
1 тс рс	/ _{TM} PM	BO0	NutBac	ct	TH	MRadSpe
BO0TSSTKNTOCCOO	Phe Su1 pH	nide nolics fide & Grease			MPN Total Coli MPN Fecal Coli MF Total Colif MF Fecal Colif Fecal Streptoc Plate Count —	forms/100 ml orms/100 ml orms/100 ml occi/100 ml
CATIONS			ANIONS	:	<u>101</u>	AL METALS
Armonia	Lead		Bicarbonate		Aluminum	\ Lead
Arsenic	Magnesium		Carbon Dioxide		X Arsenic	Manganese Mangane
Barium	Manganese		Carbonate		X Barium	Mercury
Boron	Nickel		Chloride		Berylliu	-
Cadmium	Potassium		CO ₃ Solids		× Cadmium	Nickel
Calcium	Selenium		Fluoride		Chromium	
Chromium	Silver		Hydroxide		Cobalt	Silver
Chromium, Hex	Sodium		Nitrate		X Copper	Vanadium
Copper	Zinc		Nitrite		^ 1ron	Zinc
lron			Phosphorus, Ortho			
			Silica			
			Sulfate			
	,					
		V		7	T CRAC	1817 12 CANADA
the state of the s	Phosphorus				RADIOLOG	ICS
Company of the contract of the	Nik. as Caccog					200
	Hardness as CaC	3			a, Gross	228 _{Radium}
	ity as NTU				, Gross	Uranium
The second secon	nd. (umhos/cm)		a complete	ZZ0R	adium	
Other:						

PACIFIC MINE PORTAL FLOW 200 YDS. BELOW PORTA UINTA NATIONAL FOREST 88 W 100 N PROVO UT 84603 377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PACIFIC MINE PORTAL FLOW 200 YDS. BELOW PORTA

Site ID: Source: 00
Cost Code: 3508
Lab Number: 8802862 Type: 04
Sample Date: 88/05/18 Time:

Date of Review and QA Validation Inorganic Review: 88/06/22

Organic Review:

lot. Cations:

G d Total:

Tot. Anions: 91 me/1 Cations:

91 me/l Anions:

Radiochemistry Review: 3.0 Microbiology Review:

Tot. Alk.	152	mg/l	TDS @ 180C	202	mg/1
1-Arsenic	24.0	ug/l	T-Barium	0.11	mg/l
Γ-Cadmium	9	ug/l	Γ-Chromium	(5.0	ug/1
1-Copper	62.0	ug/l	T-lron	6.6	mg/l
Γ-Lead	180.0	ug/l	I-Manganes	23.0	ug/1
Mercury	<0.2	ug/l	7-Selenium	<0.5	ug/l
[-Silver	<2.0	uq/1	Γ−Zinc	1300.0	ug/1

Water System No	Sour	Source No					
		45%			yy/mm/dd	24 hr cloc	
Euret Description	of Compline Daint	Par	Tr M'	116	PORTAL	E/W	
xact Description	or sampling Point	1/1(.16	0	10.1	Jer Inc	1600	
10	10 you 1.	Clou	i'oR tr	9			
collector: PA	UL H.	SKAB	ELUNI)	County:	Cost Code: 35	
					> 7 Telephon		
iddress: EX	W 100 K)	MROVE	(1)	7114	zip: 5 1/60	
1 тс рс		BOD	Nut	Bact	PestTHM	RadSpe	
BOD	Cyani				MPN Total Colife		
TSS	Pheno			-	MPN Fecal Colife		
TKN TOC	Sulfi	oe .		-	MF Total Colifor		
		Grease		-	Fecal Streptoco		
	011 &	urease	80	-	Plate Count - 0		
			<u> Parisson</u>				
CATIONS			ANIONS		TOTAL	L METALS	
Ammonia	Lead		Bicarbonate		Aluminum	Lead	
Arsenic	Magnesium		Carbon Dioxide		Arsenic	- V Manganese	
Barium _	Manganese		Carbonate		Barium	Mercury	
Boron	Nickel	The second second	Chloride		Beryllium		
Cadmium	Potassium Selenium		CO ₃ Solids Fluoride		Cadmium Chromium	Nickel Selenium	
Calcium Chromium	Silver		Hydroxide		Cobalt	V Silver	
Chromium, Hex	Sodium		Nitrate		V Copper	Vanadium	
Copper	Zinc		Nitrite		U Iron	Zinc	
copper 1ron	21110	_	Phosphorus,Ort	ho .	IIIII	EIIIC	
			Silica				
		-	Sulfate	8			
	1						
PENG.				R. S.			
	Phosphorus				RADIOLOGIC	<u>CS</u>	
	Alk. as CaCO3					220	
	lardness as CaCO3				lpha, Gross _	228 _{Radium}	
	ity as NTU		5 - 16	B	eta, Gross	Uranium	
	nd. (umhos/cm)				²⁶ Radium		
.X_TDS @							
Other:	OF ACTOR VIEW	UARRE	William Co.				

LOWER PAC.MINE PORTAL ACROSS STREAM FROM BAKE UINTA NATIONAL FOREST 88 W 100 N

PROVO

U1 84603

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: LOWER PAC.MINE PORTAL ACROSS STREAM FROM BAKE

Site 10:

Source: 00

Cost Code: 3508

Lab Number: 8802863 Type: 04 Sample Date: 88/05/18 Fime: 10:45

Date of Review and QA Validation Inorganic Review: 88/06/22

lot. Cations:

Tot. Anions: Gr lotal:

109 me/l Cations:

109 me/l Anions:

Organic Review: Radiochemistry Review: 3.6 Microbiology Review:

Tot. Alk.	183	mg/1	TDS @ 180C	204	mg/1
1-Arsenic	<1.0	ug/l	1-Barium	0.036	mg/1
I-Cadmium	<1	ug/l	1-Chromium	<5.0	ug/1
T-Copper	<20.0	ug/l	1-1ron	0.048	mg/1
[-Lead	<5.0	ug/l	I-Manganes	6.0	ug/1
Mercury	<0.2	ug/l	1-Selenium	<0.5	ug/1
T-Silver	<2.0	ug/1	ſ-∠inc	<20.0	ug/1

;;===:am2863

Water System No	Source	No	Date Collected		Collected / / /
		1 - 15 -	2	yy/mm/dd	
Exact Description	of Sampling Point:	Lawete	- TAC	1011101	PERHAL
A	TRUSS 5	TREAM	1 FROM	M BAKER	FORK DONK
Collector: PAL		CABEL			Cost Code: 350
Send Report to:	JINTA NI	TIONA	K FOIZ	(S) Teleph	one No: 377-57
Address: 8	#25 V	NON			11.7- zip: 84605
		FE 401 CAL 100 EL		*************	
PCPC	PMPM _	B00	NutBact	PestT	HMRadSpe
	Cyanide			MPN Total Col	iforms/100 ml
TSS	Phenoli			MPN Fecal Col	
TKN	Sulfide			MF Total Coli	
TOC	pH			MF Fecal Coli	
	0i1 & G	rease		Fecal Strepto Plate Count -	
					0.9.7
CATIONS		AN	IONS	. <u>10</u>	TAL METALS
			Particular 14		V
Ammonia	Lead .		rbonate	Aluminu	-
Arsenic	Magnesium .		on Dioxide	Arsenic	- William William
Barium Boron	Manganese .	Carb	onate	. <u>×</u> Barium . Berylli	Mercury um Molybdenu
Cadmium	Potassium .			. Z Cadmium	
Calcium	Selenium .	Fluo	Solids	. X Chromiu	
Chromium	Silver		oxide	Cobalt	k Silver
Chromium, Hex	Sodium .	Nitr		. Copper	Vanadium
Copper	Zinc .	Nitr		. Iron	₹ Zinc
1ron		- management	phorus,Ortho		
		Sili			
		Sulf			
	٠.				
	Name to the second			DADTOLO	otes
	Phosphorus			RADIOLO	uics
	Nlk. as CaCO ₃ Hardness as CaCO ₃			Alpha, Gross	228 _{Radium}
	ity as NTU		- 550	Beta, Gross	Uranium
	nd. (umhos/cm)		1 650	226 _{Radium}	oranian
又TDS @			The Part		
Other:			S. 150 C. S. S.		
W. W. S. S. S.					

01	R-DATE/SAMPLE-NUMBER.:[#/
UTAH STATE WATER QUALITY SYSTEM MONITORING RUN PROGRAM	
MO RING RUN:[]	T.
STORET:[] SOURCE:[]	COUNTY:[] USE:[] COST CODE:[]
DESCRIPTION: [North Fork American	Folk above Bog Mina]
COLLECTOR: [P][A][U][L][][H][][S][A	· ·
DATE: [8][8][0][7][2][0] TIME: [[][0][2][5] TYPE:[][]
	FIELD TESTS
TEMPERATURE: [1][0].[0] pH: [][6].[7] D.O.: [][][][][][] SP. COND.: [][][][][][] SP. GRAVITY: [].[][][][] TRANSPARENCY: [][][][]	CO2: [][][][] [] DEPTH: [][][][].[] C1 RESID.: [].[][] FLOW (MGD): [].[][][][].[] FLOW (GPM): [][][][][][].[] FLOW (CFS): [][][][][][][][][][]
SA	MPLE BOTTLES NEEDED
CI. STRY: [2]	
NUTRIENTS:[7]	
TOTAL METALS:[2]	
FIELD COMMENTS: TAKE FLOW	Junep = -07' Width = 3.6' Depth = -3'

01:			R-DATE/SAMPL	E-NUMBER.:[#3)
UTAH STATE WATER MONITORING RUN	QUALITY SYS	TEM			
MOI RING RUN:[3			
STORET:[]	SOURCE:[] co	[]:YTNUC	USE:[] COS	ST CODE:[]
DESCRIPTION:[No	RTH FORK	AMERIC	CAN FORK BEL	OW LONER BOGMI	(E)
COLLECTOR: [P][A]	[1][1][1][1]	[][5][*][A][B][E][L]	[][][][D][][][11 1
DATE: [8][8][0][7 Y Y M M			/][/][5][0] FIELD TESTS	TYPE:[][]	
TEMPERATURE: pH: D.O.: SP.COND.: SP.GRAVITY: TRANSPARENCY:	11].[] [][] [][]	DEPTH: C1 RESID.: FLOW (MGD): FLOW (GPM):	נ זנ ז	
		SAM	PLE BOTTLES N	EEDED	
CH TRY:[2]					
NUTRIENT:[7]					
TOTAL METALS:[2]					
				011 - 11 - 2	//
FIELD COMMENTS:	TAKE FLOW	Jump	= .03' W	id:1/ :5' De	p Ph = . 351

TAKE FLOW Jump = . 10' Width = 5.5' Depth = . 51

FIELD COMMENTS:

01	*	R-DATE/SAMPLE-NUMBER .: [#5]
UTAH STATE WATER MONITORING RUN			
MO RING RUN:[]		
STORET:[]	SOURCE:[]	COUNTY:[] USE:[] COST CODE:[1
DESCRIPTION: [PA	CIFIC MINE -	NW PORTAL	
COLLECTOR: [P][A]	[U][L][][H][][S][[A][A][B][E][L][U][M][D][][][][]	
DATE: [8] [8] [0] [7] Y Y M M][각[0] TIME: D D	[[][3][O][O] TYPE:[][] FIELD TESTS	
	/		
TEMPERATURE: pH: D.O.: SP.COND.: SP.GRAVITY: TRANSPARENCY:	[0].[0]\$\frac{1}{2}\$\] [1].[1][1] [1].[1][1] [1].[1][1][1] [1].[1][1][1] [1].[1][1][1]	CO2: [][][][] [] DEPTH: [][][][][][] Cl RESID.: [].[][] FLOW (MGD): [].[][][][][] FLOW (GPM): [][][][][][][] FLOW (CFS): [][][][][][][][]	
	s	AMPLE BOTTLES NEEDED	
CH TRY:[2]			
NUTRIENTS: [7]			
TOTAL METALS: [2]			
FIELD COMMENTS:	TAKE FLOW Size	ne as from Mine Portal at sang	le

01	R-DATE/SAMPLE-NUMBER . : [#-4
UTAH STATE WATER QUALITY SYSTEM MONITORING RUN PROGRAM	ω
MO RING RUN:[]	
STORET:[] SOURCE:[]	COUNTY:[] USE:[] COST CODE:[]
DESCRIPTION: [Pacific Mine - C	enter of Tailings
COLLECTOR: [P][A][U][L][][H][][S][A	· ·
DATE: [8][8][0][7][2][0] TIME:	[/][3][/][0] TYPE:[][] FIELD TESTS
SP. COND.: [][][][][] SP. GRAVITY: [].[][][]	CO2: [][][][] DEPTH: [][][][][] CI RESID.: [].[][] FLOW (MGD): [].[][] FLOW (GPM): [][][][][][].[] FLOW (CFS): [][][][][][][].[]
SA	MPLE BOTTLES NEEDED
CH STRY: [2]	
NUTRIENT: [7]	
TOTAL METALS: [2]	
FIELD COMMENTS: TAKE FLOW	
<u> </u>	

01.	R-DATE/SAMPLE-NUMBER.:[#7
UTAH STATE WATER QUALITY SYSTEM MONITORING RUN PROGRAM	
MC RING RUN:[]	
STORET:[] SOURCE:[]	COUNTY:[] USE:[] COST CODE:[]
DESCRIPTION:[1
COLLECTOR:[P][A][U][L][][H][][5]	[K][A][B][E][L][V][N][D][][][][][][]
DATE: [8][8][0][7][2][0] TIME	:[/][4][5][5] TYPE:[][]
1 1 2 2 5	FIELD TESTS
TEMPERATURE: [2][0].[0] pH: [][6].[7] D.O.: [][].[] SP.COND.: [][][][][][] SP. GRAUITY: [].[][][] TRANSPARENCY: [][].[]	CO2: DEPTH: C1 RESID.: FLOW (MGD): FLOW (GPM): FLOW (CFS): [][][][][][][]. []
S	SAMPLE BOTTLES NEEDED
CHL_STRY: [2]	
NUTRIENT: [7]	
TOTAL METALS: [2]	
FIELD COMMENTS: TAKE FLOW June	p = .03' Depth = .15' Width = 2.75'
<u> </u>	

]

R-DATE/SAMPLE-NUMBER.:[#8
COUNTY:[] USE:[] COST CODE:[]
~ Fork River Below Pacific Mine]
[K][A][B][E][L][V][N][D][][][][]
:[/][4][5][0] TYPE:[][]
FIELD TESTS
CO2:
SAMPLE BOTTLES NEEDED
p = .04' Depth = .5' Width = 6ft'
from Month of Dry Fork

01	R-DATE/SAMPLE-NUMBER.:[#9
UTAH STATE WATER QUALITY SYSTEM MONITORING RUN PROGRAM	
MC RING RUN:[
STORET:[] SOURCE:[]	COUNTY:[] USE:[] COST CODE:[]
DESCRIPTION: [North Fork Americ	an Forh River a Dutchman Flat]
COLLECTOR: [P][A][U][L][][A][][.	S][x][A][B][G][L][U[N][D][][][][][]
DATE: [8] [8] [0] [7] [2] [0] TIN	ME:[/][差][3][5] TYPE:[][]
TEMPERATURE: [/][6].[3 pH: [][6].[8 D.O.: [][]].[SP.COND.: [][][][][][][SP. GRAUITY: [].[][][][TRANSPARENCY: [][].[].[
	SAMPLE BOTTLES NEEDED
CH TRY: [2] NUTRIENTS: [7]	
TOTAL METALS:[2]	
FIELD COMMENTS: TAKE FLOW J4	mp = .151 Width = 81 Depth = .67'

11 14	40 14	h a	PART THE 18	and an area areas and a second	.4
DESCRIPTION: [MAK	EY ELEEN GULC	H CREEK A	T MOUTH]	
COLLECTOR: [P][A][v][L][][H][][s][k][A][B][E][L]	[U][N][D][]	[][][]	
DATE: [8][8][0][7]		[/][6][/][0]	TYPE:[]	[]	
		FIELD TESTS			
TEMPERATURE: pH: D.O.: SP.COND.: SP.GRAVITY: TRANSPARENCY:	[][][][]	CO2: DEPTH: C1 RESID.: FLOW (MGD): FLOW (GPM): FLOW (CFS):][][][][]]]][][][] [].[][] [].[][]]]][][][].[7]	
	SA	MPLE BOTTLES	NEEDED		
CHEMISTRY: [2]					
NUTRIENT:[7]					
TOTAL METALS:[2]					
FIELD COMMENTS:	TAKE FLOW Jump	=.14' Wid	th =3.2'	Depth = -18'	

01	R-DATE/SAMPLE-NUMBER.:[#//
UTAH STATE WATER QUALITY SYSTEM MONITORING RUN PROGRAM	
MO. RING RUN:[]	
STORET:[] SOURCE:[] CO	UNTY:[] USE:[] COST CODE:[]
DESCRIPTION: [North Fork American	Fork below-Mary Ellen Gulch]
COLLECTOR: [P][A][U][L][][A][][5][K]	[A][B][C][L][U][N][D][][][][]
DATE:[8][8][0][7][2][0] TIME:[1][6][3][0] TYPE:[][]
	IELD TESTS
SP.COND.: [][][][][][]	CO2: DEPTH: C1 RESID.: FLOW (MGD): FLOW (GPM): FLOW (CFS): [][][][][][][][][][][][][]
SAMI	PLE BOTTLES NEEDED
CH. JTRY: [2]	
NUTRIENT: [7]	
TOTAL METALS: [2]	
TOTAL PICTALS. [2]	
FIELD COMMENTS: TAKE FLOW Jump :	: 103' Depth = .625' Width =7'

]

01.	R-DATE/SAMPLE-NUMBER .: 14/2
UTAH STATE WATER QUALITY SYSTEM MONITORING RUN PROGRAM	
MC RING RUN:[]	
STORET:[] SOURCE:[] CO	DUNTY:[] USE:[] COST CODE:[]
DESCRIPTION: [MARY ELLEN GULCH	CREEK ABOUG MINE TAILINGS
COLLECTOR:[P][A][V][V][][H][][5][K	[][][][][0][N][N][][][][
	0][9][4][0] TYPE:[][]
Y Y M M D D	FIELD TESTS
pH: [][6].[7] D.O.: [][][][] SP.COND.: [][][][][]	CO2:
SAM	PLE BOTTLES NEEDED
CH TRY:[2] NUTRIENTS: [7]	
TOTAL METALS: [2]	
FIELD COMMENTS: TAKE FLOW Jump	=.02' Width = 1.5' Depth =.13'

FIELD COMMENTS:	TAKE FLOW	Jump = .03'	width = 1.0'	Depth = . 21'	
"三"				1/2	

1 7 0 1103932

DESCRIPTION: [West Fork Mary Ellen Gulch Creek a Mouth]

COLLECTOR: [P][A][V][V][][H][][5][K][A][B][G][L][V][N][P][][][][]

DATE: [8][8][0][7][2][/]

TIME:[/][/][0][0]

TYPE:[][]

FIELD TESTS

TEMPERATURE: [1][3].[0] CO2: [][6].[7] DEPTH: D.O.: [][].[] C1 RESID.:

SP.COND.: [][][][][] FLOW (MGD):

SP. GRAVITY: [].[][][] FLOW (GPM):

TRANSPARENCY: [].[][] FLOW (GPM): [][][][].[] FLOW (CFS): [][][][][][][TRANSPARENCY:

SAMPLE BOTTLES NEEDED

CHETTRY: [#] -T-ALK, HRONS, SOY, TIDS

NUTRIENT: [2]

TOTAL METALS: [2]1 - AS, BA, CO, BCR, CU, FE, PB, MN, Hg, SE, AG, ZN

TAKE FLOW Jump = .07' Width = 1.5' Depth = .3' FIELD COMMENTS:

FIELD COMMENTS:	TAKE FLOW	Jump = .01	Wid 14 = 1.5	Depins

01, .,	R-DATE/SAMPLE-NUMBER .: [-# /4]
UTAH STATE WATER QUALITY SYSTEM MONITORING RUN PROGRAM		
MC DRING RUN:[]		
STORET:[] SOURCE:[]	COUNTY:[] USE:[] COST CODE:[]	
DESCRIPTION: [MARY ELLEN GU	LCH CR. BELOW MINE TAILINGS	
COLLECTOR: [P][4][U][4][][H][][5][K][A][B][G][L][[V][M][D][][][][][][]	
DATE: [8][9][0][7][2][1] TIM	E:[/][/][3][5] TYPE:[][]	
YYMMDD	FIELD TESTS	
pH: [][6].[7 D.O.: [][][][].[SP.COND.: [][][][][][] C1 RESID.: [].[][]] FLOW (MGD): [].[][]] FLOW (GPM): [][][].[]	
	SAMPLE BOTTLES NEEDED	
CHL STRY: [2] NUTRIENT: [7] TOTAL METALS: [2]		
FIELD COMMENTS: TAKE FLOW Ju	imp = .04 Width = 3.0' Depth =	43' -

	2 RECEIVING OFFICE INC.	3 CONTRACT NUM	MER	and the second second	ONDER DATE	5 SF-2				B ORDER NUMBE		9 SUB.
1 OF DI	CHASE (Check one)	OH	-		05/20/88	AT		cone	175	43-8462	2-8-1447	7
	HASE ORDER	DELIVERY	ORDER									
AN	STATE HEAL	State, Eip Code, and	Phone No.)	* * -	11 Cens			NAT ROL		of Inspection and A	Acceptance	
WATE	R LAB-STATE	HEALTH LA			SHIP					TT (PAUL SI	KABEL UN	33
40 H	EDICAL DRIV		Passa		TO	88	WES'	T 100		TFF		
	1533-6131	0.1	84113			PRO	7.75	586-57	***		84601	
1 12	1202-0131		14			& No.1		17	18	Check > FI	TS ▶ A	COMM ▶
ACT CODE		- WATER SA	SCRIPTION		BUD		NE C	YTITHAUC	UNIT	UNIT PRICE	201	AMOUNT 11904
	F. IRON;	ALS: A. AU M; D. CHRCS G. LEAD; N. RY; J. SELI ND ALSO FOI SCLIDS.	Kiuh; e. (. Pabganes Enium; K.	OPPER; SILVER;	TAL							
		- WATER SAI INITY AND 1				500 0	1	1	EA	15.0	00	15
	TO HELD TO SELECT				200						1	
1	3000											- 1
5												
			and the make	e e e e e e e e e e e e e e e e e e e	and the state of t	6 1 64	*	nikarist.	i eiti	on the second	# 15 PE 150	heriosi.
	Harest S	A. 2460	mi s vije			10 m			被整	et tricke.	24 K.S	Marin .
	Hours of a	Kan dia Kana	est of which					ray to	de la company	a ville	24 156	Marie .
	tente de		18 C 18 18			\$/3 %			福裝			M. Marie
	eto alconoto		us (x x x			10		en e	福宏	di Tibler	A 16.8	E.W.
PHIS PUR	CHASE ORDER NE	GOTIATED PURS	UANT TO AUTE	ORITY OF 41 U				/x:-:-	福裝			Marie
O.B. POINT	r vz.	GOTIATED PURS	22 DISCOUNT A	ND/OR NET PAYME	S.C. 252(e)		TYPE	COMMODIT).		25	
DESTI	MATION F.O.B. POINT ON OR I	BEFORE (Date)	22 DISCOUNT A	A STATE OF THE PARTY OF THE PAR	S.C. 252(e)	22a	TYPE	COMMODITED TO CO). ryi	Sub-Total	25	1205.
DESTI	MATTON F.O.B. POINT ON OR I PER CONTRACT	BEFORE (Data)	22 DISCOUNT A	ND/OR NET PAYME	S.C. 252(e)	22a	TYPE	COMMODIT ENT O). ryi	Sub-Total TOTAL	25	1205.
DESTI	MATION F.O.B. POINT ON OR I	BEFORE (Data)	22 DISCOUNT A	ND/OR NET PAYME	S.C. 252(e)	22a	TYPE	COMMODITED TO CO). ryi	Sub-Total	> 25 > 27	1205.
DESTI	MATTON F.O.B. POINT ON OR I PER CONTRACT	BEFORE (Data)	22 DISCOUNT A	ND/OR NET PAYME	S.C. 252(e)	228	TYPE PAYM CODE ESTIM	COMMODITION OF THE PROPERTY OF). ryi	Sub-Total TOTAL	> 25 > 27	1205.
DESTI	MATTON F.O.B. POINT ON OR I PER CONTRACT	BEFORE (Data)	22 DISCOUNT A	ND/OR NET PAYME	S.C. 252(e)	22a	TYPE PAYM CODE ESTIM	COMMODITED TO CO). ryi	Sub-Total TOTAL	> 25 > 27	1205.
O.B. POINT DESTIP ELIVER TO	FO.B. POINT ON OR I	BEFORE (Data) ON B 10	22 DISCOUNT A	ND/OR NET PAYME	S.C. 252(e)X NT TERMS	22e 26 1-1-0 D BY (Near	TYPE PAYMEN CODE ESTIM	COMMODITION OF THE PROPERTY OF).). (1) (1) (2) (3) (4)	Sub-Total TOTAL 30 DISTRIBUTION 2	> 25 > 27 > 3	1205. 1205. 1205. 1205.
O.B. POINT DESTINATION OF THE POINT OF THE P	FOR POINT ON OR I	BEFORE (Date) ON B 10	22 DISCOUNT A NO	ND/OR NET PAYME	S.C. 252(cX) NT TERMS. 0 4	22s 26 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	TYPE PAYME COLOR ESTIMATE OF C	COMMODIFIED OF ATED FREE 1-062 1). (YY) (O)	Sub-Total TOTAL 30 DISTRIBUTION 2	25 27 27 A	1205 1205 31 AMOUNT
O.B. POINT DESTU	FOR POINT ON OR I PER CONTRACT UNTING CLASSIFICATI A SCUTE ACQUITE SCUTE ACQUITE TO REST SERTI	BEFORE (Date) ON B 10	22 DISCOUNT A NO	ND/OR NET PAYME	S.C. 252(e)X NT TERMS	22s 26 26 GIAL PHO	TYPE PAYMEN CODE ESTIM	COMMODITION OF THE PROPERTY OF).),)(GHT O	Sub-Total TOTAL 30 DISTRIBUTION 2	25 27 27 A	1205. 1205. 1205. 1205.
O.B. POINT DESTU	FOR POINT ON OR I	BEFORE (Date) ON B 10	22 DISCOUNT A NO	ND/OR NET PAYME	S.C. 252(cX) NT TERMS. 0 4	22s 26 To D BY (New A FO) (SO) (SO)	TYPE PAYME OF STIME O	COMMODIFIED OF OR ATED FREE .00).),)(GHT O	Sub-Total TOTAL 30 DISTRIBUTION 2	25 27 27 A	1205 1205 31 AMOUNT

Fund Skabelund WEST FORK-MARY-ELLEN GULCH CREEK @ MOUTH UINTA NATIONAL FOREST 88 W 100 N PROVO UT 84603

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: WEST FORK MARY ELLEN GULCH CREEK @ MOUTH

Site 10:

Source: 00

Cost Code: 3508 Lab Number: 8803932

Lab Number: 8803932 Type: 04 Sample Date: 88/07/21 Time: 11:00

Date of Review and QA Validation

Inorganic Review: 88/08/22

Organic Review:

Tot. Cations: 43

Tot. Anions: 103 me/l Cations: and Total: 146 me/l Anions:

2.5 Radiochemistry Moview: 2.6 Microbiology Review:

Laboratory Analyses

Sulfate	67	mg/l	Tot. Alk.	60	mg/1
1. Hardns.	121.7	mg/1	TDS @ 180C	190	mg/1
I-Arsenic	14.5	ug/1	T-Barium	0.022	mg/1
1-Cadmium	2	ug/1	1-Chromium	(5.0	ug/1
T-Copper	53.0	ug/1	I-Iron	1.2	mg/1
1-Lead	10.0	ug/l	T-Manganes	74.0	ug/1
Mercury	(0.2	ug/1	T-Selenium	<0.5	ug/l
T-Silver	<2.0	ug/l	T-Zinc	450.0	ug/l

sife.		Ar senic ug 1.1. 22.5 12.0	Cal - run ug 1.1. 31 9	apper ug / 1.	4,000 850.0	Mercury ug [1. .63 .29	silver u.g/l. 5.0	garium ng 1/1. 0.28	chamilan ag/L 450	#\$77. 5.3	manganese ug 61. 23.0	scientary ag// 40.5	1,600.0	Sultice Mg/1.
8	North FK American FK Einer Along Below Pacific Mine April 18	4.5	21 21	<20.0 <20.0	20.0 45.0	≥0.2							81.0 < 20.0	23
								2 kg 2 kg						
9	North Fork American Fork spill River a Dutch man Flat youls 9/22/18	2.5 41.0	۷۱ ۷۱ ۷۱	< 20.0 < 20.0 < 20.0	60.0 5.0 5.0	< 0.2	<2.0 <2.0	0.05%	< 5.0 < 5.0	0.45	31.0	40.5	77.0 43.0 37.0	16 21
10	Mary Eller Ereck 2 Mondh foots	2.5	<1	<20.0 <20.0	2 5.0 5.0	< 0.2	< 2.0	0.046	45.0	0.059	4.0	< 0.5	72.0 41.0	23 26
11	North Fork American Fork thatse There below Mary Ellen 9/2/81 Soulch	3.5	41 41	<20.0 <20.0	10.0 < 5.0	۷.2	42.0	0.053	<5.0	0.089	7.0	40.5	46.0 99.0	25 52
12	Mary Eller Gulch Creek 7/2/18 Above Mixed Areas 9/2/18	1.5	41	<20.0 <20.0	< 5.0 < 5.0	< 0.2	< 2.0	0.041	<5.0	0.625	45.0	40.5	22.0	22 30

site 7		Arsinic ug/1. 22.5 22.0	Ordinizem ug (1. 31	aniper ug/1. 60.0 30.0	3	Sulface My 11.	10 mes Mall. 1124	77 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -	TG8 - mg//.	Flows GFS.
8	North FX American PR River 10088 Below Pacific Min & 9/2/8	4.5	41	420.0 420.0		23	14.6	/ 15 /:9	134 156	
9	North Fork American Fork \$18 88 River a Dutch man Flat 1/20/88	2.5	۷۱ ۷۱	< 20.0 < 20.0 < 20.0		16 21	157.3 166.4	83 /43 /43	102 174 168	
10	Mary Eller Creek 2 Month 1/20/78	2.5	<	<20.0 <20.0		33 26	135.8 162.3	109 136	154 190	
	North Fork American Fork 1/20/88 River be low Mary Ellen 9/22/88 Gulch	3.5	<u> </u>	<20.0 <20.0		20	150.7 156.5	135 89	188	
12	Mary Eller Gulch Creek pulls Above Mined Areas 9/2/188	1.5	۷	<20.0 <20.0		22 30	111.0	92 80	/20 /32	

Sife No.	Station Mary Ellen Mine Parful	Date Arsenic Slate 100 7/2/18 97 9/2/18	Calmium ng/1.	Copper 19/1. 40.0 <20.0 <20.0	49 / 1. 10.0 65.0 65.0	Mercury ug 11.	\$1 (ver ng/1. < 2.0	Barrio - mg / 1	Chromitation 48.71.	Iron mg/1. 9.9	Manganese ug/1- 140	511 mum ug/1- 40.5	2176 1,200.0 570.0 140.0	sulfate mg/1.
/3A	Month West Fork Mary 5Then Gulch	112/188 14.5 9/2/188	2 2	53,0 <20.0	/0.0 < \$.0	4.2	<2.0	0.022	< 5.0	1.2	74.0	<0.5	450 390	67
14	Mary Eller Gulch Greek Downstream from Mines	- shifts = 1.0 - shifts = 3.0 - shifts = 3.0	z <1 <1	42 ~20.0 < 20.0	40.0 < 5.0 < 5.0	<42	<2.O	0.039	4 5.0	1.1	46.0	< 0.5	310 110 92	38 69
3A	North Fork American For & River Above Rich Mine	2/20/88 1.0	٤/	< 20.0	< 5.0 < 5.0	20.2	< 2.0	0.044	< 5.0	0.033	4 5.D	< 0.5	<20.0 <20.0	18
-	Miller Hill Mine downstru From Sheific Mine	-56/st <1.0	۷.	<26.0	<5.0	< 0.2	<2.0	0.034	45.0	0.048	6.0	<0.5	<20.0	

చ

+ =										
ité 13	Station Mary Elien Mine Portal	Date 4152 5/1/18 100 7/2/188 97 5/2/188 97	enic Calman	20.0	40.0	Sulfate mg/.(-	mg/1- Total- Hundriss	Total Alkalinity 19917-	TD5 Mg/1.	Flow CFS.
		Service of		~20.0	0.0	73	115.1	40	196	
13A	Month West Fork Many	Thus		TO HER LEVEL TO						
	Month West Fork Mary Eller Gulch	olulis 14.5	2 2	53.0 <20.0	1	67	121-7	60	190	
		41.	~ ~	<20.0	<	84.	121-7	33	182	
162					1					
14		. 108								
14	Mary Eller Gulch Greek Downstream from Mines	5/18/2 × 1.0	2 2	42				92	/22	
	DOWNSTREAM Trom Mines	7/2/108 3.0	> <1	42 ~20.0 < 20.0	<	38	137-4	100	/32 /38	
		distife	41	< 20.0	<	38 69	136.8	62	166	
5A	Worth Fork American	2/20/86 1.0	<1	< 20-0	20	18	120.9	104	120	to the second season.
	Fork River Above Pacific	" Ist its	< 1	< 20.0	<.0	25	134.9	109	130 148	
	-ine						00.00.00.22.0	100	170	
	Miller Hi / Mine downstrum, From Facific Mine	sisis <1.0	41	<20.0	40			183	204	
	No.				1					

WATER QUALITY SAMPLING - UPPER AMERICAN FORK CANYON

North Fork America FRRIVER Above Bog Mine	e 7/20/28	<1.0	<1	<20.0	2 < 5.0	40.2	<2.0	0.037	Chromiun < 5.0	n Iron 0.16	Manganese 24.0	50 wishim <0.5	Zinc 28.0	5.1 fale 29
Lower Eng. Mine	जीतहरू ग/काश्च नृह्य हिंह	1.5 3.0	/Z /3 8	<20.0 <20.0	< 5.0 5.0 < 5.0	< 0.2	< 2,0	0.057	< s.o	7.9	270	<0.5	510.0 530.0	
North Fk. American Fork below Lower Bog Mine	7/20/88 Y20/88		ı , y	<20.0	< 5.0 < 5.0	1 V							77.0 190.0	51 31 44
Parlal Pacific Mine	5/18/28 7/20/78 9/52/88	22.0	6 13 5	34.0 42.0 <20.0	25.0 /5.0 < 5.0	0.2 <0.2	42.0	0,063	45.0	4.0	11.0	4.0.5	900.0	
NW Fortal Pacific Mine	shrise 71=0 00	/.o 2.0	د ا	< 20.0	60.0 < 5,0	<0.2	<2.0	0.15	< €.0	0.091	19.6	4 ೩೯	73.0	40
Center of Tailings below Main Facific Mine Fortal (from Runoff)	- HOISE	13.0	9 8 (51)	62.0 30.0 (260.0)	180.0	< 0.2	<2.0	0.11	< 5.0	6.6	23.0		1,306.0	
				(200.0)	(20,000.0)	(3.24)	(45.0)	(0.15)	(4 E.O)	(13.0)	(48.6)		1,000.0 7,700.0	




				SJATER	2 GUSL					
200	Station North Fork America FEEiver Abon Bog Mire	in July	freenic 49/1 <1.0	, dad mit	(20) ×20.	7 file 29	Harders	# 1 / P	783 128	Flor FAS.
2	Lower Boy Mine	डीतांचर ग/२०/ख	1.5 3.0	12	<200			c	90	
		35, 131		/3 8	<20.C	51	39. 3	0	72	
3	North FR. American Fork below Lower Bog Mine		Z.5	1		3/	96.9	73	120	
	or a sold sof mine	9/20/28		4	<20.0 1	45	106.0	53	.38	
4	Portal Pacific Mine	7/20/88	22.0	6	54.0 42.0			/43	202	
		9 22 88		5	<20.0 4	0	204-4	0	356	
2111										
5	NW Fortal Pacific Alms	5/13/28 7/20/83	1.0	<1	< 20.0			198	208	
	4 7									10
6	Center of Tailings below	5 11/98	24.0	9	62.0			/52	202	
	(from Runoff) -	7/20/86	13.0		30.0					
	(nom nunott) -	Dielos (30.0)	(51)	(260.0)			21	140	

40.00

AMERICAN FORK HYDROLOGY AND WATER QUALITY STUDY

Lidstone & Anderson, Inc.

Water Resources and Environmental Consultants



DIVIDION OF

AMERICAN FORK HYDROLOGY AND WATER QUALITY STUDY

PREPARED FOR:

Utah Division of Oil, Gas and Mining Abandoned Mine Reclamation Program 355 W. North Temple Triad Center, Suite 350 Salt Lake City, UT 84180-1203

and

United States Forest Service Uinta National Forest 88 West 100 North Provo, UT 84601

PREPARED BY:

Lidstone & Anderson, Inc. 736 Whalers Way, F-200 Fort Collins, CO 80525

February 3, 1993

TABLE OF CONTENTS

1.0	INTE	RODUCTION	1
	1.1 1.2	Site Conditions	
2.0	PHY	SIOGRAPHIC AND GEOLOGIC SETTING	3
	2.1	Geochemical Setting of the Project Area	6
3.0	SAM	PLING RESULTS	6
	3.1 3.2 3.3	Lower Bog Mine	7
4.0		ER QUALITY IMPACTS TO THE NORTH FORK OF THE RICAN FORK	15
	4.1 4.2 4.3	Summary of Site Water Quality and Recommended Action	15
		FIGURES/TABLES/APPENDICES	
		FIGURES	
Figure Figure Figure Figure Figure	2. 3. 4.	Location Map: American Fork Canyon Water Quality Study. American Fork Canyon Sampling Sites Schematic. Lower Bog Sampling Program. Pacific Mine Sampling Program. 1 Trilinear Diagram Characterizing the	4 8
Figure Figure	6.	Water Quality at Pacific Mine	13
Figure	8.	Trilinear Diagram Characterizing the Water Quality at Mine Partals	
Figure	9.	Conceptual Design of Reclamation at the	2

TABLE OF CONTENTS (CONTINUED)

TABLES

Table 1.	American Fork Canyon Water Sampling Program.	5
Table 2.	Water Quality Samples Which Exceed Class 3A Aquatic Fisheries Standards (4-Day Average)	18

APPENDICES

Appendix A: Water Quality Data Appendix B: Gaging Data

1.0 INTRODUCTION

A water quality investigation was conducted at several abandoned mine sites in the American Fork Canyon, Utah County, Utah during the three day period of July 7th through the 9th, 1992. The project was cooperatively funded by the Utah Division Oil, Gas and Mining (DOGM), Abandoned Mine Reclamation Program (AMR) and the U.S. Forest Service, Uinta National Forest. Several previous studies had been conducted in the area including:

- (1) Merritt, Lavere B., 1988; "Preliminary Survey of Water Quality in Mine Drainage in Sheeprock Mountains and North Fork of the American Fork River." (Water Quality).
- (2) Mangum, Fred, 1988; "Aquatic Ecosystem Inventory, Macroinvertebrate Analysis; Annual Progress Report, Uinta National Forest". (Water Quality and Macroinvertebrates).
- (3) Kastning-Culp, Nancy, et.al., 1992; "Year End Report On Mitigation Systems for Hard Rock Mine Effluent in Utah". (Soils, Water Quality, Vegetation, Ecosystems).


1.1 Site Conditions

The American Fork Canyon Mining District is characterized by inactive underground mine workings, shafts, portals, spoils and tailings located in the Uinta National Forest. The majority of these workings are associated with valid mining claims. A number of abandoned mine sites have been inventoried by the Utah DOGM in the past. The scope of the current sampling study was to specifically investigate three mine drainage problem areas: the Pacific Mine, the Lower Bog Mine and the Mary Ellen Gulch Mines (Figure 1).

In many cases the underground workings of inactive mines are flooded by ground water. This ground water comes in contact with the mineralized rock, spent ore and/or tailings, which results in changes in water chemistry. Typically this change manifests itself as lower pH conditions and higher concentrations of trace metals. Where there is sufficient ground water "head" or gradient, the mine water is discharged to the surface and enters area streams. If toxic levels of trace metals are present in these mine waters, an adverse impact to area streams or aquatic life can occur.

1.2 Site Investigation

The purpose of this study is to investigate the hydrology, geochemistry and water quality impacts of mine drainage on receiving waters within the National Forest Lands. Following the analysis of the water quality impacts, a conceptual "action plan" will be developed. This report documents the sampling study, the laboratory analyses, and a mass balance analysis of the water quality in the vicinity of the three study sites.



Field investigations were conducted by Lidstone & Anderson, Inc. and a representative of the Utah Division of Oil, Gas & Mining, AMR Program to determine existing conditions. These investigations included water sample collection, flow estimates, measurements of field water quality and soil pH parameters. Additional analyses included observations of geological and mineralogical conditions, natural biological and geochemical controls or hydrochemical barrier conditions present at each site.

Flow estimates were made at each portal and in the vicinity of sample points using a bucket and stop watch. Flow estimates were made along major drainages (Mary Ellen Gulch and the North Fork of the American Fork) using a Pygmy Current Meter. Field water quality parameters included field pH (Orion Research Model No. 200), field conductivity and temperature (YSI Model No. 33) and color. Water samples were collected and handled using standard EPA sampling protocol. Samples were unfiltered, preserved in the field, packed in ice and delivered to the Utah Department of Health laboratory within 24 hours of collection. Laboratory analysis included major anions and cations, total dissolved solids, total alkalinity and selected acid soluble trace metals.

Figure 2 presents the sample sites in relationship to the mine portals and receiving streams. Field pH and laboratory TDS characterize the water quality at each sampling point. Flow discharge measurement points and estimates are presented on this figure. Table 1 documents the field sampling program, a description of each sample site and the field parameters measured at each site. The analytical results and a conceptual sketch of each site showing the relative locations of sample sites are presented in Appendix A. Gaging measurement data sheets are presented in Appendix B.

2.0 PHYSIOGRAPHIC AND GEOLOGIC SETTING

The American Fork Canyon study area is situated within the upper headwaters of the North Fork of the American Fork River in Utah County, Utah. The locations of the American Fork River and its various tributaries are shown on Figure 1. The North Fork is a south west-flowing drainage tributary to the American Fork River, which drains into Utah Lake, the Jordan River and eventually into the Great Salt Lake. The headwaters of the American Fork Canyon in the vicinity of the project area range in elevation from 9,200 to over 10,000 feet above sea level. The drainage originates in a glaciated cirque basin, known as Mineral Basin at the base of Mount Baldy. Mary Ellen Gulch is a southeast draining tributary to the North Fork of the American Fork, entering the American Fork Canyon at Dutchman's Flat. The project area within Mary Ellen Gulch ranges in elevation from 8,800 to 9,400 feet above sea level. This drainage originates in a glaciated cirque basin, known as Merril Flat at the base of Twin Peaks.

The streams draining the divide are steep gradient cobble- to boulder-bed streams. The flow conditions of the streams range from rapid to turbulent along most of the project area reaches. The drainage pattern is dendritic with most tributaries sustaining a base flow throughout most of the year.

The geologic setting of the project area is extensively fractured and mineralized carbonate and metasedimentary rocks of Paleozoic or Precambrian Age. The oldest rocks within the




FIGURE 2. AMERICAN FORK CANYON SAMPLING SITES SCHEMATIC

Table 1. American Fork Canyon Water Sampling Program.

			Discharge		Field P	arameters		
Sample No.	Data Collected	Time	Discharge	pH (s.u)	TDS (ppm)	EC (umhos)	Temp.	Location
AF#1	7/8/92	12:56 PM	6.5 gpm*	7.8	_	325	18.3	Discharge from Pacific Mine ab. confluence w/American Fork, through tailings
AF#2	7/8/92	- 1:20 PM	9.5 cfs ^b	8.0	130	170	13.3	American Fork bl. Pacific Mine
AF#3	7/8/92	12:20 PM	9.2 cfs	8.4	140	150	11.3	American Fork ab. Pacific Mine
AF#4	7/8/92	12:25 PM	12 gpm	8.0	-	280	11.7	Discharge from Pacific Mine after treatment in Beaver Pond ab. confluence
AF#5	7/8/92	2:25 PM	44.5 gpm	5.1	80	-	10.1	Discharge form Lower Bog Mine Portal
AF#6	7/8/92	3:30 PM	144 gpm	6.5	180	230	7.8	Discharge from Pacific Mine Portal
AF#7	7/8/92	5:05 PM	70 gpm	5.9	140	180	8.0	Discharge from North Portal Mary Ellen Gulch
AF#8	7/8/92	5:50 PM	0.55 cfs	8.1	-	140	9.1	Mary Ellen Gulch us. of AMR and active mine disturbance
AF#9	7/8/92	7:15 PM	1.50 cfs	7.9	-	170	10.4	Mary Ellen Gulch ds. of AMR and active mine disturbance
			1989 B	Miscel	laneous Sampl	ing Sites	00000	
-	7/7/92	-	2.5 gpm°	7.2	-	205	7.0	Mary Ellen Gulch South Portal
-	7/7/92	-	0.6 cfs ^e	7.7	-	105	10.2	Trib. North of North Portal Mary Ellen Gulch Mis us. of AMR disturbance
_	7/6/92	-	0.0	6.9	260	-	22.2	Ponded water on tailings at Pacific Mine
-	7/8/92	2:45 PM	5-9 cfs*	7.9	110	-	10.0	N. Fork American Fork ab. Lower Bog Mine discharge.
-	7/8/92	2:55 PM	5-9 cfs*	7.5	100	-	11.2	N. Fork American Fork bl. Lower Bog Mine discharge

gpm measured utilizing a stopwatch and bucket cfs measured utilizing a pygmy meter flow visually estimated

immediate project area compromise the Late Precambrian Big Cottonwood Formation, which consists of quartzites, shales and metasedimentary rocks. The formation is approximately 16,000 feet thick (James, L. P., 1979) and is well exposed on steeply dipping exposures along the American Fork Canyon and Mary Ellen Gulch. It is exposed along the American Fork channel immediately below the Lower Bog Mine, as well as along the steeper reaches of Mary Ellen Gulch. The Paleozoic sequence within the project area consists of the Cambrian Age Tintic Quartzite, Ophir Formation and Maxfield Limestone, and the Mississippian Age Fitchville Formation, Deseret and Gardison Limestones. The Pacific Mine portals lie within a fault graben block of Gardison Limestone. The Mary Ellen Gulch mine portals are situated in Cambrian Age Maxfield Limestone and dolomites of the Mississippian Fitchville Formation. The Lower Bog Mine portal was "driven into" the Precambrian Big Cottonwood Formation.

2.1 Geochemical Setting of the Project Area

Mineralization and ore trends within the project area are closely associated with the Miocene age emplacement of silicic, intermediate and aplite dikes of the Alta Stock (James, L.P. 1979). The rocks of the Alta Stock are typically granodiorite to quartz monzonite in composition. Mineralization and alteration trends are concordant with the extensive faulting and fracturing of the host rocks. Historical mining in the area generally followed these ore trends. The chemistry of the Alta Stock and the mineralization within the American Fork Canyon is high in copper, lead, zinc and iron. The high arsenic and cadmium concentrations present in the mineralized zones are associated with accessory minerals, which occur as the sulfides, arsenates and carbonate minerals.

The characteristics of the mine drainage chemistry are a reflection of the relationship of host rock chemistry, the surrounding equilibrium conditions of waters in contact with the mineralized or "mined zone" and upgradient ground water quality. The "mined" or mineralized zone is high in both primary sulfides, secondary sulfates and hydrous sulfates. Because of the high sulfide content of the mineralized rock, one would typically anticipate acid mine drainage from the American Fork portals. Of the three sites investigated, two sites are characterized by nearly neutral pH conditions: the Pacific Mine and the Mary Ellen Gulch Mines. In both cases the host rocks are limestones or dolomites and are rich in carbonates. Although the oxidation of the sulfides within the mineralized zones continues to occur and generate acid pH conditions, the buffering capacity of the upgradient ground water quality is such that the water is neutralized upon exiting the mine portal. Acid drainage is present at the Lower Bog Mine (pH ranges from 3.9 to 5.1). The host rock at the Lower Bog Mine is predominantly quartzites, siltstones and shales of the Big Cottonwood Formation. The host rock and the upgradient water quality does not have the capacity to bùffer the acid mine drainage conditions at this site.

3.0 SAMPLING RESULTS

3.1 Lower Bog Mine

The Lower Bog Mine portal is located at an approximate elevation of 8,520 feet AMSL and consists of a single bedrock opening, tailings dump and miscellaneous spoil piles. Discharge

from the portal was gaged at approximately 44.5 gpm on July 8, 1992. The water was clear with "yellow boy" or hydrous iron oxide precipitate in the vicinity of the discharge. On that day, field pH was measured at 5.1 and the field analysis of total dissolved solids was 80 ppm. Based on the considerable amount of iron precipitate at the mouth of the portal discharge, these results were somewhat surprising. A single water sample was collected at the site. The laboratory results (Figure 3) suggest that the discharging waters were not in equilibrium at the time of sampling. A laboratory pH value of 3.9 suggested a greater change in pH (from field to lab) than would be anticipated. The laboratory cation-anion balance was 21%. Typically acceptable laboratory balance is less than 5%. The 1992 sample results are similar to the 1988 (Merritt, 1988) sampling effort eliminating laboratory error as the sole problem.

To evaluate the impact of the Lower Bog portal discharge on the North Fork of the American Fork, field parameters were measured at various points within the hydrologic system. The portal discharge enters the main stream at two points (Figure 3): (1) as surface flow adjacent to a tailings dump and, (2) as seepage through the tailings dump. At the surface flow location, the pH had increased from the upstream value of 5.1 to 6.4. At the seepage location the pH had increased from 5.1 to 7.0, suggesting the neutralization of waters in transit from the mouth of the portal to its confluence with the main stream. On the date of sampling (7/8/92), measurements of field parameters upstream and downstream of the point of confluence were made to determine if there was any impact to the waters of the American Fork. Upstream of the portal discharge, a pH of 7.95 and total dissolved solids content (TDS) of 110 ppm were measured. Downstream of the portal discharge a pH of 7.52 and a TDS of 100 ppm were measured, suggesting that dilution is the principal mechanism for the mitigation of adverse impacts. Discharge of the receiving waters on July 8, 1992 was estimated at 3.31 cfs (from basin area reduction of measured channel discharges along the North Fork and Mary Ellen Gulch). The portal discharge was measured at 44.5 gpm or 0.1 cfs reflecting a dilution of 33:1.

The 1992 water quality analysis of the Lower Bog Mine portal indicate that excessive concentrations of trace metals (iron, cadmium, zinc, copper and lead) are associated with the portal discharge. Similar studies at the adjacent mines (Pacific and Mary Ellen Gulch) indicate that copper and iron concentrations are not problematical since these parameters are strictly pH and Eh dependent. Cadmium, zinc and lead behave in a slightly different geochemical manner. Sampling completed by Mangum, 1988 indicated that upstream concentrations of zinc averaged approximately 20 ug/l during a July and September sampling period. Downstream of the Lower Bog discharge, zinc concentration increased to 77 (in July) to 190 ug/l in September. Sampling of macroinvertebrates at two stations (Mangum, 1988) indicated that the effects of the portal discharge resulted in "stress conditions along the lower reach".

3.2 Pacific Mine

The Pacific Mine is located at an elevation of 7800 feet AMSL and consists of two discharging portals, a tailings dump, miscellaneous mine-related structures and spoil piles. An upper or northwest portal was not investigated as part of this study. Previous studies (Merritt, 1988) had indicated that additional dissolution of trace metals occurred where the discharge from the south portal commingled with an abandoned tailings dump. Kastning-Culp, et. al., 1992 investigated the biological uptake of trace metals by an adjacent wetlands/beaver pond north of




FIGURE 3. LOWER BOG SAMPLING PROGRAM

the tailings dump. The 1992 sampling program was designed to investigate the impacts of the portal discharge on the receiving waters (North Fork of the American Fork), the influence of the interaction of the tailings with the portal discharge, and the positive, if any, influence of the beaver pond on the discharging water quality. Figure 4 presents the sampling program conducted at the Pacific Mine site.

Field parameters were measured at the five sampling sites on consecutive days and were found to be repeatable during the sampling period. Drainage from the main portal (AF#6) is characterized by a near neutral (6.54) pH, iron precipitate and high concentrations of trace metals, primarily lead, zinc, copper and cadmium. Flow at the mouth of the main portal was gaged at 144 gpm or 0.32 cfs. At the base of the first bench and approximately 110 feet from the mouth of the main portal, the portal flow splits at a spoils dump and load out structure. The main flow is diverted to the north towards a beaver pond. A secondary flow is diverted to the south, commingling with a tailings dump. Much of the flow along this channel appears to be subsurface flow and may exit the site as seepage. Sample AF#1, which was collected from the tailings surface flow (measured at 6.5 gpm) is characterized by an increase in pH relative to the upstream sampling site (AF#6). Trace metals concentrations at this site either remained the same as AF#6 or decreased as a function of the increase in pH and Eh. The lead concentration, however increased significantly (approximately 10 times). This increase appears to be primarily tailings related. Previous sampling by Merritt, 1988 bore out this relationship though at a significantly greater magnitude (160 time increase in lead concentration). Dr. Merritt's sampling took place during a "rain storm" which may have influenced the magnitude of the trace metal concentrations.

A sample (AF#4) was collected at the mouth of the beaver pond prior to commingling with the waters of the North Fork drainage. Sampling data from this point (Figure 4) suggest that the beaver pond is efficiently removing most trace metals from solution. Most of the iron and copper were precipitated out of the waters prior to entrance into the beaver pond. Cadmium and zinc which exhibit similar geochemical behavior were reduced in concentration by approximately 50%. Lead concentrations were below detection limits at the mouth of the beaver pond.

Samples AF#3 and AF#2 were collected from the main stream at sites upstream and downstream of the Pacific Mine disturbance. The waters upstream of the mine disturbance meet all Class 3A standards for aquatic wildlife. Downstream of the mine (AF#2), the waters exceed state criteria for lead. This sample exhibits an impact of the mine discharge in its four-fold increase in zinc. Zinc levels approach the aquatic standard. Studies by Mangum, 1988 indicated that "the number of organisms (macroinvertebrates) had decreased approximately 70% from an upstream to a downstream station in the vicinity of the Pacific Mine."

Figure 5 characterizes the changes in water quality character (major anices and cations) at the Pacific Mine. Trilinear diagrams typically are used to present the relative chemical characteristics of waters collected from different locations. Qualitatively, if two samples or data points plot in the same field on a trilinear diagram a common source of ions is indicated. It is no surprise that the five samples plot within the same field and can be classified as calcium-magnesium bicarbonate waters. Both ground water and surface water sources at this site are strongly influenced by site geology. The portal discharge is more sulfate- rich than the receiving




FIGURE 4. PACIFIC MINE SAMPLING PROGRAM




FIGURE 5. TRILINEAR DIAGRAM CHARACTERIZING
THE WATER QUALITY AT PACIFIC MINE

△ AF #1
□ AF #2
○ AF #3
* AF #4
● AF #6

waters, yet the relative dilution of the portal discharge by the main streams waters (30:1) result in the "commonality of ions" portrayed on Figure 5. The portal discharge was measured at 0.32 cfs. The North Fork of the American Fork was gaged at 9.2 cfs.

3.3 Mary Ellen Gulch Mines

The Mary Ellen Gulch Mines are located along a south east-flowing tributary drainage to the North Fork of the American Fork at an average elevation of 9,100 feet AMSL. The site consists of a number of mine portals, abandoned structures, sedimentation ponds and detention structures, tailings and waste rock piles and spoil dumps. At the time of the field visit, active mining was ongoing at an adjacent and upstream mine. There was recent evidence of attempts to control the north portal drainage at the Mary Ellen Gulch Mine. Field parameters were collected from two discharging portals: the south portal (pH= 7.2; EC= 205 umhos/cm) and the north portal (pH= 5.95; EC= 180). Since the most significant discharge (70 gpm vs. 2.5 gpm) originates from the north portal, only that portal was sampled (Figure 6). The sampling program at the Mary Ellen Gulch Mines was developed to ascertain the impacts of the AMR portal discharge on the receiving waters, Mary Ellen Gulch. Prior to the initiation of this project it was understood that other abandoned mines and dumps were present in the upper basin, but that the Mary Ellen Gulch north portal may have had the most significant impact on the drainage and the fishery.

On the day the Mary Ellen Gulch Mines were sampled, the Globe Mine, immediately upstream of the AMR site was discharging "milky sediment-laden water". The discharge ceased at approximately 5:30 PM that day. In an attempt to collect the most representative downstream sample, AF#9, was collected at 7:15 PM. Fine sediment, a reflection of the Globe Mine discharge, was present on the stream gravels throughout the downstream reach.

The discharge from the main north portal (AF#7) was acidic (pH=5.95) with "yellow boy" and iron oxide precipitates near the mouth of the portal. The sample data from the 1992 sampling program indicated that the trace metal concentrations of this portal were not very high with only zinc, and iron exceeding aquatic standards. Previous sampling efforts (Merritt, 1988) found that elevated levels of copper, lead and cadmium originated from this portal. A sample collected upstream of the AMR disturbance and along Mary Ellen Gulch, AF#8, is characterized by good water quality. Class 3A aquatic standards were achieved for all parameters. The downstream sample, AF#9, may have been influenced by the discharges from the active underground mine above the AF#7 sampling location. Despite any such influence the 1992 sample analysis was very similar to the previous sample analysis by Merritt, 1988 which exhibited elevated concentrations of zinc, iron, copper and lead. Copper and lead appear to originate from some source other than the mine portal and may be related to the upstream Globe Mine or possibly to adjacent spoils and tailings dumps within the Mary Ellen Gulch basin.

A trilinear diagram (Figure 7) characterizes the transitional change in water quality character (major anions and cations) at the Mary Ellen Gulch Mines. The waters discharging from the portal (AF#7) are calcium-magnesium sulfate waters. The waters of Mary Ellen Gulch prior to "mixing" (AF#8) are calcium-magnesium bicarbonate type waters. Once these waters are mixed (AF#9) at the dilution ratio naturally occurring on-site (10:1) the waters change




FIGURE 6. MARY ELLEN GULCH SAMPLING PROGRAM




FIGURE 7. TRILINEAR DIAGRAM CHARACTERIZING
THE WATER QUALITY AT MARY ELLEN GULCH

□ AF #7
LEGEND ○ AF #8
△ AF #9

chemistry to a calcium magnesium sulfate-bicarbonate- type water. The portal discharge was measured at 70 gpm. The main stem of Mary Ellen Gulch was gaged at 1.50 cfs.

4.0 Water Quality Impacts to the North Fork of the American Fork

4.1 Site Geochemistry

It is important to understand the geochemical changes, which occur as the mine discharge water exits the mine portals and before it enters the main stream. In general the water quality exiting the mine portals (Figure 8) is a calcium-magnesium sulfate-type water. The Pacific Mine drainage is predominantly calcium-magnesium bicarbonate water. The drainage from these portals are typically high in cadmium, copper, lead, iron and zinc. The anomalous concentrations of trace metals in the waters exiting these mine portals are directly related to the trace element geochemistry of the ore zones (Chapter 2.1). Copper and iron concentrations in water are strongly Eh and pH dependent. In the case of the mine portal discharge the majority of the iron precipitates out of solution as the waters become oxidized and the pH increases to neutral. The copper coprecipitates as a copper carbonate and is removed from the solution as Eh increases.

The trace metals zinc, cadmium and lead are somewhat more problematical since they are mobile under a wider range of Eh and pH conditions. Lead is the least mobile of these latter three elements and its solubility under oxidizing conditions is controlled by the presence of the carbonate ion and to a lesser degree, the sulfate ion. Under reducing conditions, lead will precipitate as a sulfide. Lead concentrations in the waters at the American Fork mines do not appear to be directly related to discharge from the mine portals but rather to contact with an outside source, either the tailings at the Pacific Mine or an adjacent upstream mine source, such as the Globe Mine within Mary Ellen Gulch.

Cadmium and zinc have similar geochemical behavior and are mobile under oxidizing conditions and nearly all pH conditions present at the American Fork sites. Cadmium levels are relatively low at the source and appear to rapidly decrease with dilution and to a certain degree by plant uptake. Chelation and/or adsorption of cadmium by organic matter in the beaver pond at the Pacific Mine appears to have a positive impact on trace metal concentration. Further discussion of these processes can be found in Kastning-Culp, et.al. 1992. concentrations of zinc are the most serious trace metal water quality problem in the American Fork Canyon. Zinc concentrations remain elevated at all stations sampled. Dilution of the portal discharge by the main channel flow appears to be the most significant mechanism for the reduction of zinc concentrations. Plant uptake of zinc, adsorption of zinc on hydrous manganese and iron oxides, adsorption and chelation of zinc by organic matter in the beaver pond at the Pacific Mine currently reduce concentrations of zinc in the effluent waters. Over time reducing conditions will develop within the beaver pond, accelerating the process of zinc removal as zinc sulfide precipitate. The limiting factor for sulfide precipitation at all American Fork sites is the degree of sulfate present in the water. With the exception of the Lower Bog site, nearly all project "receiving waters" are carbonate-rich.




FIGURE 8. TRILINEAR DIAGRAM CHARACTERIZING
THE WATER QUALITY AT MINE PORTALS

△ AF #5
LEGEND □ AF #7
○ AF #6

it was felt to be most applicable to this project occause past sampled water quality reflects a long term average condition.

Table 2 presents a summary of the American Fork water quality sampling program in relationship to the four-day average aquatic standard. On a site by site basis, samples AF#2 and AF#9 reflect the water quality at locations downstream of the disturbance and within the receiving waters, the American Fork and Mary Ellen Creek. No downstream sample was collected below the Lower Bog Mine. Previous sampling efforts by Mangum, 1988 document the zinc concentrations above and below the Lower Bog Mine.

Sample AF#2 (Table 2 and Figure 4) was collected approximately 800 feet below the Pacific Mine and exceeds aquatic fisheries standards for lead by a factor of four (4). Zinc concentrations at the downstream sample are slightly below the Aquatic Class 3A standards, yet are significantly elevated (four times) above background or upstream water quality. It is anticipated that zinc concentrations downstream of the Pacific Mine will exceed Class 3A water quality during certain periods of the year. The principal source of the elevated lead concentration at the Pacific Mine is the tailings dump adjacent to the North Fork of the American Fork. The principal source of the elevated zinc concentration is the water discharging from the south portal of the Pacific Mine. Based on the impacts of the Pacific Mine on the receiving water quality, remedial action at this site is recommended.

Sample AF#9 (Table 2 and Figure 6) characterizes the downstream water quality of Mary Ellen Gulch below the Mary Ellen Gulch Mine. This sample exceeds Class 3A water quality standards for cadmium, copper, iron, lead and zinc. Of these parameters, copper, lead and zinc are of primary concern. Copper exceeds standards by a factor of 4.5; lead exceeds standards by a factor of 13.2; zinc exceeds standards by a factor of 3.6. All parameters are significantly elevated above the upstream water quality sample AF#8. An insufficient number of samples were collected at this site to fully characterize the source of the trace metal contamination of Mary Ellen Gulch. The upstream sample, AF#8, eliminates the abandoned Yankee Mines (Figure 1) as a source of the metal contamination. Sample AF#7 was collected from the

Table 2. Water Quality Samples Which Exceed Class 3A Aquatic Fisheries Standards (4-Day Average).

Aquat	ic Standard*				Sample Concentration						
	4-Day	AF#1	AF#2	AF#3	AF#4	AF#5	AF#6	AF#7	AF#8	AF#9	
pН	6.5-9.0 su	-	-			3.9		6.0			
As	190 μg/l										
Cd	1.3 μg/l	14			5	14	12		-	2	
Cu	13.3 μg/l		-			30	47			60	
Fe	1000 μg/l					9100	4500	7800		1100	
Pb	3.8 µg/l	130	15			10	15			50	
Se	5 μg/l	-									
Zn	119 μg/1	1700	_	-	810	660	1800	800		430	

^{*} Hardness dependent criteria (pertaining to Cd, Cr, Cu, Pb, and Zn) assumes 115 mg/l total hardness

discharge waters of the Mary Ellen Gulch North Portal. Although cadmium and zinc were elevated at this source, only zinc exceeded Class 3A standards. It appears that an adjacent source must contribute toxic levels of trace metals, in particular lead and copper. That source could be the upstream and active Globe Mine or possibly runoff from the Mary Ellen Gulch tailings or the abandoned Mary Ellen Gulch South Portal. Before any mine reclamation can proceed at this site, additional water and soil sampling is necessary to clearly define the source of the contamination and maximize the positive effects of the reclamation.

No 1992 downstream sample was collected at the Lower Bog Mine site. Field parameters (Table 1) collected upstream and downstream of the mine discharge and along the North Fork of the American Fork suggest that there is minimal impact to the receiving waters (pH and TDS). Sampling of the discharging waters from the Lower Bog portal suggest that the waters exiting the mine portal reflect poor water quality, exceeding Class 3A standards (Table 2) for pH, cadmium,, iron, copper, lead and zinc. With the exception of pH and iron, the metals concentration of the Lower Bog Mine portal (AF#5) is less than the Pacific Mine portal (AF#6). When comparing the dilution ratio (receiving water flow to the portal discharge) it is apparent that there is greater dilution at the Lower Bog Mine than at the Pacific Mine. Assuming similar geochemical conditions, one can predict that the impact of the Lower Bog Mine discharge on the American Fork River will be less than the impact of the Pacific Mine discharge. The principal contaminants of interest will be zinc and possibly lead. Sampling conducted in 1988 (Mangum, 1988) indicated that zinc concentration will exceed Class 3A standards during the low water period of the year by a factor of 1.6. Because of the site's inaccessibility and the limited magnitude of the problem, no action is recommended at the Lower Bog Mine site.

4.3 Proposed Mine Reclamation

The 1992 water quality investigations quantified the environmental impacts of the AMR disturbances on the North Fork of the American Fork. Additional study is recommended at the Mary Ellen Gulch sites. No further action is recommended at the Lower Bog Mine. Sufficient water quality data are available at the Pacific Mine to document the nature and magnitude of the environmental problem at this site. AMR and/or USFS action is recommended at this site to mitigate the adverse impacts of past mining activities.

Available funding, land and mineral owner consent and final land use may restrict the degree of mine reclamation and ultimately its success in the mitigation of adverse impacts. On this basis a phased approach is recommended. Two interrelated sources of contamination will have to be addressed at the Pacific Mine: (1) portal discharge and (2) the tailings pond adjacent to the creek.

The primary source of contamination, the tailings dump is responsible for the elevated lead levels in the American Fork at sample site AF#2. Lead concentrations are transported to the creek via mine portal discharge as surface and subsurface flow, overland flow in response to rainfall and snowmelt events and bank erosion and channel migration of the American Fork against the tailings embankment. This study did not quantify the relative metals loading of each mechanism of transport.

The tailings dump should be isolated from the American Fork through a combination of cut and fill, rerouting of the portal discharge drainage and revetment of the existing American Fork channel banks. All portal discharges should be routed in a permanent diversion channel directly to the beaver pond. Because of the steep gradient of this channel, riprap protection will be required. The riprap will serve a multiple purpose of protecting the permanent diversion from erosion, oxidizing the discharging portal waters, raising the pH of the waters and coprecipitating the iron from solution, as well as serving as a permanent and maintenance-free barrier to ATV traffic attempting to access the tailings dump site. On site limestone or dolomitic rock can be utilized as riprap. Screening and sorting of this rock will be required to ensure a well graded riprap blanket.

The east slope of the tailings dump should be excavated from the vicinity of the North Fork of the American Fork channel. These materials should be transported to the top of the tailings dump and regraded to a "domed", yet relatively flat (less than 3% grade) surface. Ponded areas on the existing tailings dump should be eliminated. The outslope (east) of the regraded tailings dump should be graded to no steeper than a 4:1. The regraded surface of the tailings should be "deep ripped and limed" to elevate the pH of the tailings above 6.5. Topsoil can be borrowed from adjacent sites and placed on the regraded and limed surface. A minimum of 12 to 15 inches of topsoil should be placed on site. Care should be taken to separate A and B horizon material at the borrow site to ensure that an organic rich layer of A-horizon material is available for final cover. This same material will serve as a natural seed source and will reduce revegetation costs. The site should be broadcast seeded and harrowed. A riprap bank apron or at a minimum, toe slope riprap protection should be placed along the outslope, adjacent to the creek. Depending on the characteristics of available rock, this riprap may have to be imported to the site. Wooden cribs or similar biotechnical slope protection may be substituted for riprap. However longevity of the design should be addressed.


Additional treatment of the discharging portal waters can be accomplished through the construction of a wetland on the upper terrace immediately above the beaver pond. The purpose of this wetland is to accomplish primary treatment of zinc and cadmium, prior to the water's entrance into the beaver pond. The beaver pond would behave as a secondary treatment facility. The wetland would be excavated into the surface adjacent to the "loadout" area. Approximately 4,000 square feet of surface is available for wetland construction. An impermeable liner and coarse limestone gravels would be placed at the bottom of the excavation. Organic matter (humus, manure, soils borrowed from the beaver pond area) would be backfilled above the gravel layer. The site would be topsoiled and planted with the appropriate locally available vegetation. Kastning-Culp, 1992 documents the chelation properties and plant uptake of zinc by local vegetative species. Soil and moss berms would be constructed within the wetland to prevent short-circuiting of the influent waters. The wetland would discharge directly to a ditch, which would flow to the beaver pond and ultimately to the North Fork of the American Fork.


Under a phased approach, the initial reclamation should entail a channel diversion of all portal discharges to the beaver pond. "Follow-up" water quality sampling should take place to evaluate the beaver pond's ability to treat the additional waters. Later phases should include the limited cut and fill and regrading of the tailings dump, channel stabilization of the North Fork in the vicinity of the tailings dump and the construction of the wetland.

A conceptual design of the Pacific Mine proposed reclamation is presented on Figure 9. Surveying and mapping should be completed prior to the finalization of the designs. The design process should include an evaluation of design hydrology, channel hydraulics, soils and vegetation requirements, final earthwork, preparation of final plans and specifications.

FIGURE 9. CONCEPTUAL DESIGN OF RECLAMATION AT THE PACIFIC MINE

APPENDIX A WATER QUALITY DATA





Pacific Mine seepage above confluence w/ American Fork

pH = 7.75temp. = 18.3EC = 325color = clear

UTAH STATE HEALTH DEPARTMENT DIVISION OF LABORATORY SERVICES Environmental Chemistry Analysis Report

Description: AMERICAN FORK #1

Site ID: Source: 00 Date of Review and QA Validation Cost Code: 350B Inorganic Review: 92/07/29

Organic Review: Lab Number: 9204265 Type: 04 Sample Date: 92/07/08 Time: 12:45 Radiochemistry Review:

Tot. Cations: 68 Microbiology Review: Tot. Anions: 140 mg/1 4.1 me/1 Cations:

Grand Total: 208 mg/1 4.2 me/1 Anions:

Laboratory Analyses

			*** F	
L-pH *	7.9	D-Calcium	43 mg/	1
D-Magnesum	23 mg/1	D-Potassum	<1 mg/	1
Bicarbnate	206 mg/1	Carbonate	0 mg/	1
Chloride	<1 mg/1	Sulfate	38.045 mg/	1
Tot. Alk.	169 mg/1	TDS @ 180C	220 mg/	1
H+Arsenic	5.0 ug/1	H+Barium	0.11 mg/	1
H+Cadmium	14 ug/1	, H+Chromium	<5.0 ug/	1
H+Copper	<20.0 ug/1	H+Iron	0.3 mg/	1
H+Lead '	130.0 ug/1	H+Mangan	92.0 ug/	1
H+Selenium	<5.0 ug/1	H+Zinc	1700.0 ug/	1

pH should be performed as a field test.

JUL 3 1 1992

North Fork of American Fork below Pacific Mine

> pH = 8.05temp = 1333 EC = 170color = clear

UTAH STATE HEALTH DEPARTMENT DIVISION OF LABORATORY SERVICES Environmental Chemistry Analysis Report

Description: AMERICAN FORK #2

Site ID:

Source: 00

Date of Review and QA Validation

Cost Code: 350B

92/07/29

Lab Number: 9204266 Type: 04

Sample Date: 92/07/08 Time: 13:05

Radiochemistry Review:

Tot. Cations: 42

Cations: 2.5 me/l

Microbiology Review:

Tot. Anions: Grand Total:

80 mg/1 122 mg/1

Anions:

2.5 me/1.

Inorganic Review:

Organic Review:

Laboratory Analyses

L-pH *	8.0 :	D-Calcium	28 mg/1	
D-Magnesum	12 mg/1	D-Potassum	<1 mg/1	
Bicarbnate	128 mg/1	Carbonate	0 mg/l	
Chloride	<1 mg/1	Sulfate	15.889 mg/1	
Tot. Alk.	105 mg/1	TDS @ 180C	132 mg/1	
H+Arsenic	(5.0 ug/1	H+Barium	0.053 mg/1	
H+Cadmium	c1 ug/1	H+Chromium	<5.0 ug/1	
H+Copper	<20.0 ug/1	H+Iron	0.23 mg/l	
H+Lead	15.0 ug/1	H+Mangan	21.0 ug/1	
H+Selenium	<5.0 ug/1	H+Zinc	99.0 ug/1	

pH should be performed as a field test.

JUL 3 1 1992

N. Fork of American Fork above Pacific Mine

> pH = 8.42temp = 11.3EC = 150Color-= clear

UTAH STATE HEALTH DEPARTMENT DIVISION OF LABORATORY SERVICES Environmental Chemistry Analysis Report

Description: AMERICAN FORK #3

Source: 00 Site ID:

Date of Review and QA Validation

Inorganic Review: 92/07/29

Cost Code: 350B

Lab Number: 9204267 Type: 04 Sample Date: 92/07/08 Time: 12:30

Organic Review:

Radiochemistry Review:

Tot. Cations: 39

77 mg/l Cations: Microbiology Review: 2.3 me/1

Tot. Anions: Grand Total:

116 mg/1

Anions:

2.3 me/1

Laboratory Analyses

L-pH *	7.9 .	D-Calcium	26 mg/1
D-Magnesum	11 mg/1	D-Potassum	<1 mg/1
Bicarbnate	119 mg/1	Carbonate	0 mg/1
Chloride	c1 mg/1	Sulfate	17.572 mg/1
Tot. Alk.	97 mg/1	TDS @ 180C	138 mg/1
H+Arsenic	<5.0 ug/1	H+Barium	0.043 mg/1
H+Cadmium	<1 ug/1	H+Chromium	<5.0 ug/1
H+Copper	<20.0 ug/1	H+Iron	0.14 mg/1
H+Lead	<5.0 ug/1	H+Mangan	16.0 ug/1
H+Selenium	<5.0 ug/1 ·	H+Zinc	23.0 ug/1

pH should be performed as a field test.

JUL 3 1 1992

Seepage Discharge from Beaver Pond above confluence of American Fork pH. = 7.98temp. = 11.7 EC = 280 color = clear

UTAH STATE HEALTH DEPARTMENT DIVISION OF LABORATORY SERVICES Environmental Chemistry Analysis Report

Description: AMERICAN FORK #4

Site ID:

Source: 00

Date of Review and QA Validation

Cost Code: 350B

Inorganic Review:

92/07/29

Lab Number: 9204268 Type: 04

Organic Review:

Sample Date: 92/07/08 Time: 12:15 Tot. Cations:

Radiochemistry Review: Microbiology Review:

Tot. Anions: Grand Total:

136 mg/1

4.1 me/1

Anions: 204 mg/1

Cations:

4.1 me/1

Laboratory Analyses

L-pH *	7.7		D-Calcium 42	mg/1
D-Magnesum	23	mg/1	D-Potassum <1	mg/1
Bicarbnate	202	mg/1	Carbonate 0	mg/1
Chloride	<1	mg/1	Sulfate 35.646	mg/1
Tot. Alk.	165	mg/1	TDS @ 180C 218	mg/1
H+Arsenic	<5.0	ug/1	H+Barium 0.086	mg/1
H+Cadmium	5	ug/1	H+Chromium <5.0	ug/1
H+Copper	<20.0	ug/1	H+Iron 0.39	mg/1
H+Lead	(5.0	ug/1	H+Mangan 18.0	ug/1
H+Selenium	<5.0	ug/1	H+Zinc 810.0	ug/1

pH should be performed as a field test.

DECEIVE

Seepage from Pacific Mine @ Portal pH. = 6.54temp. = 7.8 EC = 230 color = sl. Cloudy, red, Fe ppt

92/07/29

UTAH STATE HEALTH DEPARTMENT DIVISION OF LABORATORY SERVICES Environmental Chemistry Analysis Report

Description: AMERICAN FORK #6

Site ID: Source: 00

Date of Review and QA Validation Cost Code: 3508 Inorganic Review: Organic Review: Lab Number: 9204270 Type: 04

Sample Date: 92/07/08 Time: 15:30 Radiochemistry Review: Tot. Cations: 65 Microbiology Review:

135 mg/1 Cations: 3.9 me/1 Tot. Anions: 200 mg/1 4.0 me/1 Grand Total: Anions:

Laboratory Analyses

L-pH *	6.9		D-Calcium 40 m	ng/1	
D-Magnesum	22	mg/1	D-Potassum <1 m	ng/1	
Bicarbnate	191	mg/1	Carbonate 0 m	ng/1	
Chloride	1.4	mg/1	Sulfate 39.473 m	ng/1	
Tot. Alk.	156	mg/1	TDS @ 180C 208 m	ng/1	
H+Arsenic	20.0	ug/1	H+Barium 0.084 m	ng/1	
H+Cadmium	12	ug/l	H+Chromium <5.0	ug/1	
H+Copper	47.0	ug/1	H+Iron 4.5 m	ng/1	
H+Lead	15.0	ug/l	H+Mangan 15.0 u	ig/1	
H+Selenium	(5.0	ug/1	H+Zinc 1800.0 t	19/1	

pH should be performed as a field test.



North Portal Mary Ellen Guich

pH. = 5.95 temp. = 8.0 EC = 180 Color = clear, Feppt

UTAH STATE HEALTH DEPARTMENT DIVISION OF LABORATORY SERVICES Environmental Chemistry Analysis Report

Description:	AMERICAN	FORK #	7	
Site ID:		Source	e: 00	Date of Review and OA Validation
Cost Code:	350B			Inorganic Review: 92/08/06
Lab Number:	9204271	Type:	04	Organic Review:
Sample Date:	92/07/08	Time:	17:05	Radiochemistry Review:
Tot. Cations:	44			Microbiology Review:
Tot. Anions:	118	mg/1	Cations:	2.6 me/l
Grand Total:	162	mg/1	Anions:	2.7 me/l

Laboratory Analyses

L-pH *	6.0		D-Calcium	20	mq/1
r-bu	0.0		D-Calcium	30	mg/ I
D-Magnesum	12 mg/	/1	D-Potassum	1.2	mg/l
Bicarbnate	30 mg/	/1	Carbonate	0	mg/l
Chloride	<1 mg/	/1	Sulfate	102.13	mg/l
Tot. Alk.	25 mg/	/1	TDS @ 180C	184	mg/l
H+Arsenic	70.0 ug/	/1	H+Barium	0.014	mg/1
H+Cadmium	1 ug/	1	H+Chromium	<5.0	ug/l
H+Copper	<20.0 ug/	1	H+Iron	7.8	mg/l
H+Lead	(5.0 ug/	1	H+Mangan	210.0	ug/1
H+Selenium	(5.0 ug/	1	H+Zinc	800.0	ug/l

pH should be performed as a field test.

RECEIVED

AUG : 1 0 1992

DIV. OIL, GAS, MIN

Mary Ellen Gulch above AML disturbance

pH. = 8.1temp. = 9.1EC = 140color = clear

UTAH STATE HEALTH DEPARTMENT DIVISION OF LABORATORY SERVICES Environmental Chemistry Analysis Report

Description: AMERICAN FORK #8

Site ID: Source: 00

Cost Code: 350B

Lab Number: 9204272 Type: 04 Sample Date: 92/07/08 Time: 17:50

Tot. Cations: 37

Tot. Anions:

73 mg/1 110 mg/1

Cations: Anions:

Date of Review and QA Validation Inorganic Review: 92/07/29 Organic Review:

Radiochemistry Review:

Microbiology Review:

2.1 me/1 2.2 me/1

Laboratory Analyses

Grand Total:

L-pH	* 8.0		D-Calcium 25	mg/l	
D-Hagnesum	10	mg/1	D-Potassum <1	mg/1	
Bicarbnate	106	mg/1	Carbonate 0	mg/1	
Chloride	<1	mg/1	Sulfate 19.91	mg/1	
Tot. Alk.	87	mg/1	TDS @ 180C 124	mg/1	
H+Arsenic	<5.0	ug/1	H+Barium 0.044	mg/l	
H+Cadmium	<1	ug/1	H+Chromium <5.0	ug/1	
H+Copper	<20.0	ug/1	H+Iron 0.08	mg/1	
H+Lead	<5.0	ug/1	H+Mangan <5.0	ug/1	
H+Selenium	<5.0	ug/1	H+Zinc <20.0	ug/1	
	D-Hagnesum Bicarbnate Chloride Tot. Alk. H+Arsenic H+Cadmium H+Copper H+Lead	D-Magnesum 10 Bicarbnate 106 Chloride <1 Tot. Alk. 87 H+Arsenic <5.0 H+Cadmium <1 H+Copper <20.0 H+Lead <5.0	D-Magnesum 10 mg/1 Bicarbnate 106 mg/1 Chloride <1 mg/1 Tot. Alk. 87 mg/1 H+Arsenic <5.0 ug/1 H+Codmium <1 ug/1 H+Copper <20.0 ug/1 H+Lead <5.0 ug/1	D-Magnesum 10 mg/l D-Potassum <1 Bicarbnate 106 mg/l Carbonate 0 Chloride <1 mg/l	D-Magnesum 10 mg/1 D-Potassum <1 mg/1 Bicarbnate 106 mg/1 Carbonate 0 mg/1 Chloride <1 mg/1 Sulfate 19.91 mg/1 Tot. Alk. 87 mg/1 TDS @ 180C 124 mg/1 H+Arsenic <5.0 ug/1 H+Cadmium <1 ug/1 H+Chromium <5.0 ug/1 H+Cropper <20.0 ug/1 H+Iron 0.08 mg/1 H+Lead <5.0 ug/1 H+Mangan <5.0 ug/1

pH should be performed as a field test.

JUL 3 1 1992

Mary Ellen Gulch below AML disturbno pH. = 7.95temp. = 10.4 EC = 170 Color = milky

UTAH STATE HEALTH DEPARTMENT DIVISION OF LABORATORY SERVICES Environmental Chemistry Analysis Report

Description: AMERICAN FORK #9

Site ID: Cost Code:

Source: 00

04

Date of Review and QA Validation

3508

Inorganic Review: 92/07/29

Lab Number: 9204273 Type:

Organic Review: Radiochemistry Review:

Sample Date: 92/07/08 Time: 19:15 Tot. Cations: 43

Microbiology Review: Cations: 2.5 me/1

Tot. Anions:

97 mg/1

Anions:

Grand Total:

140 mg/1

2.6 me/1


Laboratory Analyses

L-pH *	7.9	D-Calcium	29 mg/1
D-Magnesum	12 mg/1	D-Potassum	<1 mg/1
Bicarbnate	94 mg/1	Carbonate	0 mg/1
Chloride	<1 mg/1	Sulfate	49.504 mg/1
Tot. Alk.	77 mg/1	TDS @ 180C	148 mg/1
H+Arsenic	10.0 ug/1	H+Barium	0.034 mg/1
H+Cadmi um	2 ug/1	H+Chromium	<5.0 ug/1
H+Copper	60.0 ug/1	H+Iron	1.1 mg/1
H+Lead	50.0 ug/1	H+Mangan	60.0 ug/1
H+Selenium	<5.0 ug/1	H+Zinc	430.0 ug/1

pH should be performed as a field test.



JUL 3 1 1992



Description: AMERICAN FORK #5

Site ID: Source: 00

Cost Code:

350B

Lab Number: 9204269 Type:

66 mg/1

83 mg/1

Sample Date: 92/07/08 Time: 14:25

Tot. Cations: 17

Tot. Anions:

Grand Total:

04

Cations: Anions:

Date of Review and QA Validation Inorganic Review: 92/07/29

Organic Review: Radiochemistry Review: Microbiology Review:

0.9 me/1 1.4 me/1

Laboratory Analyses

L-pH *	3.9 :	D-Calcium	11 mg/1
D-Magnesum	3.6 mg/1	D-Potassum	1.1 mg/1
Bicarbnate	0 mg/1	Carbonate	0 mg/1
Chloride	1.4 mg/1	Sulfate	64.368 mg/1
Tot. Alk.	0 mg/1	TDS @ 180C	120 mg/1
H+Arsenic	<5.0 ug/1	H+Barium	0.035 mg/l
H+Cadmium	14 ug/1	H+Chromium	<5.0 ug/1
H+Copper	30.0 ug/1	H+Iron	9.1 mg/1
H+Lead	10.0 ug/1	H+Mangan	290.0 ug/1
H+Selenium	<5.0 ug/1	H+Zinc	660.0 ug/1

Lower Bog portal discharge
pH. = 5.11
TDS = 80 ppm
temp = 10.1
Color = clear, Fe ppt


UTAH STATE HEALTH DEPARTMENT DIVISION OF LABORATORY SERVICES Environmental Chemistry Analysis Report

Description: AMERICAN FORK #5 Site ID: Source: 00 Date of Review and QA Validation Cost Code: 350B Lab Number: 9204269 Type: 04 Inorganic Review: 92/07/29 Organic Review: Sample Date: 92/07/08 Time: 14:25 Radiochemistry Review: Tot. Cations: 17 Microbiology Review: Tot. Anions: 66 mg/1 Cations: 0.9 me/1 Grand Total: 83 mg/1 Anions: 1.4 me/l

Laboratory Analyses

L-pH *	3.9	D-Calcium	11 mg/1
D-Magnesum	3.6 mg/1	D-Potassum	1.1 mg/l
Bicarbnate	0 mg/1	Carbonate	0 mg/1
Chloride	1.4 mg/1	Sulfate	64.368 mg/1
Tot. Alk.	0 mg/1	TDS @ 180C	120 mg/1
H+Arsenic	<5.0 ug/1	H+Barium	0.035 mg/1
H+Cadmi um	14 ug/1	H+Chromium	<5.0 ug/1
H+Copper	30.0 ug/1	H+Iron	9.1 mg/1
H+Lead	10.0 ug/1	H+Mangan	290.0 ug/1
H+Selenium	<5.0 ug/1	H+Zinc	660.0 ug/1

PH pH should be performed as a field test.



APPENDIX B
GAGING DATA

								PAGE	OF
			DISCH	ARGE ME	ASUREMEN	T NOTES			7, 9-1-10
						arge (800	feet)		
TE_ 7/	8/92	PAR	TY_Lids	tone/Me	sch				
JUIPMEN	T Bygm	у		And the sale of the latest with the		elocity	_WEATHER	R_cloudy	overcast
CROSS SE	CTION_	GS#2			_ FLOW_1	rapid	Olympia.	n	Walker Carrier
					the second secon	GAGE READ			
						o; high fl			
						cobbles, w			
	graver	T Dec	i, some	SIIL UE	posic. i	Tamilings i	1=.035 D	eu, 110	bar
Distance From Initial Point (ft)			oservation Depth	Revolutions	In		city /s)		scharge (cfs)
From Poir	Width (ft)	Depth (ft)	Opse 0.09	. Revo	Time Ir Seconds	At Point	Mean	Area (ft ²)	Disc (c
LB no	0.0	0.0			(Value)	CAR EL			MALE IN
flow			amp: 1					Name and the	
LB EOW	3.2	0.0				0			
vertica	1						0.76	0.17	0.13
11	3.5	0.58	5-11-1	91	60	1.52			Mar Charle
			40			The state of the s	1.59	0.41	0.65
1 23 2 3	4.2	0.58	407	101	60	1.66		1000000	30000
		1000		* - 1			1.47	0.32	0.47 :
	4.8	0.50		77	60	1.28		42.5	12/12/11/20
			11 E 34		I STATE		1.32	0.12	0.16
10.00	5.1	0.33		81	60	1.35			
					The contract of		0.67	0.07	0.05
RB EOW	5.5	0				0		1	
9-16-16	Marie Edward		N			11-15-76-16			1.46 cfs



Lidstone & Anderson
Water Resources and Environmental Consultant

1.50

from seepage along LB

PAGE	OF	

DISCHARGE MEASUREMENT NOTES


DATE 7/8/	92	PARTY	Lidst	one/Mes	sch	RELIGIO		
PMENT	and the second	CX (n. Skr).	1	METHOD_	Name:	5 7.74	WEATHER	cloudy/overcast
CROSS SECT	TION_GS	3 #1			_FLOW_	clear	rapid	
NITIAL GA	GE REA	DING/TIM	E 10:05	Att	FINAL	GAGE	READING/TIME	10:40 AM
OMMENTS	LB look	ing DS	= 0.00.	Bed ma	aterial	2-4"	cobbles, som	e gravels -

Mannings "n" over channel length 0.045 - 0.050; OB = 0.065

Distance From Initial Point (ft)			Observation © Depth	Revolutions	In		ocity		Discharge (cfs)
From Poir	Width (ft)	Depth (ft)	opse 0.6	Revo	Time In Seconds	At Point	Mean	Area (ft ²)	Disc (c
LB EOW	0.0	0.0	0.0			17.26 1 36			
	4 - 14.					1.172-11-25	0.79	0.36	0.28
	2.2	0.33	407	47	30	1.57		可去区	
		. T.	12				2.05	0.36	074
	3.0	0.58	-4.0	76	30	2.52		ROME II	
			- 455				2.73	0.67	1.83
	4.0	0.75	7.160	44	15	2.93		Sale Co.	
4.55			100	*P #51			2.55	0.73	1.86
	5.0	0.71		32	15	2.16	E WY TO A		
							2.46	0.73	1.80
	6.0	0.75	5-18-1-1	41	15	2.76			
				-1, -4		45,670	2.19	0.71	1.55
	7.0	0.67	17 657	24	15	1.62			
		A King					1.31	0.59	0.77
	8.0	0.50		30	30	0.99	KEON	宣言共 [2]	
	To the state of		Vista III	4-51	Fig.	**************************************	0.86	0.31	0.27
	8.7	0.42	,	22	30	0.72			fire elem
			15 100 100	45- 14	AMILE.	Dr. 19	0.36	0.27	0.10
	10.0	0.0	-135	0	0	0.00			
		VIVIEW	0.5					The Lotter	
					The second				9.2 cfs
	V.C.				16.78	TOTAL ASS			



Lidstone & Anderson Water Resources and Environmental Consultant



AQUATIC TECOSYSTEM TNVENTORY

Macroinvertebrate Analysis

Annual Progress Report

NORTH FORK AMERICAN FORK RIVER AND

MARY ELLEN GULCH CREEK

UINTA NATIONAL FOREST

1988



Forest Service Intermountain Region

89-12

AQUATIC ECOSYSTEM INVENTORY Macroinvertebrate Analysis

NORTH FORK AMERICAN FORK RIVER AND MARY ELLEM GULCH CREEK UINTA NATIONAL FOREST 1988

Fred A. Mangum

Regional Aquatic Ecologist

USDA - Forest Service

Intermountain Region

Aquatic Ecosystem Analysis Laboratory

10. Page School

Brigham Young University

Provo, Utah 84602

(801) 378-4928

89-12 - MAR

TABLE OF CONTENTS

INTR	ODUCTI	ON .														1	Pag.
NORTI	ODUCTI	AWERT					*	•	•								1
MARV	H FORK	ANEKI	CAN	FOR	KRI	VER		٠		•							3
THE LAND	ELLEN	GULCH	CRE	EK	٠.												17

AQUATIC ECOSYSTEM ANALYSIS

FOR NORTH FORK AMERICAN FORK RIVER AND MARY ELLEN GULCH CREEK ON THE

UINTA NATIONAL FOREST

1988

BACKGROUND AND METHODS

In recent years land managers on many of our forests and BLM districts in the west have improved the stability and reliability of land management plans and decisions by sampling aquatic organisms which act as natural monitors of management activities within the drainages on public lands.

During short-term exposure to water of poor quality or adverse changes in habitat, organisms that cannot tolerate the stress are destroyed and the aquatic macroinvertebrate community structure changes. Since aquatic organisms respond to their total environment, they can become an effective tool for detection of environmental changes.

Our analysis of aquatic ecosystems is based upon multiple factors including:

- 1. Various macroinvertebrate data Community dry-weight biomass/sample expressed in gm/m²; number of individuals per taxa (resident populations?); DAT Diversity Index, which combines a measure of dominance and number of taxa; habit, habitat and feeding preferences of individual taxa or species; specific tolerances of taxa; community composition; and BCI (Biotic Condition Index), which indicates as a percentage how close an aquatic ecosystem is to its own potential.
 - 2. Physical parameter data and
 - 3. Water chemistry data

Effective use of the Biotic Condition Index (BCI) depends upon the availability of data on stream gradient, natural capability of instream substrate (may not be the composition present if man-influence? sedimentation is found at the sample station), total alkalinity, and sulfate in mg/1.

Because of the way that macroinvertebrates occupy space within a stream, it generally takes at least three samples to represent the community accurately at a given station. One sample per station costs less but has little value for aquatic habitat assessment, one never knows if such single samples represent the best, the worst or an average of possible conditions at the sampling site. Also as a side benefit, three samples per station provides a basis for various statistical analyses, if random samples are all taken from a rubble substrate in as similar habitat as possible, taking into account mainly the velocity of flow and depth in the stream. Biologists have found that compared to other sampling devices, the Winget-modified surber net yields the highest coefficient of correlation (similarity of samples).

A stream's natural potential for productivity, habitat quality and water quality can be compared to the "actual" by taking quantitative samples of aquatic macroinvertebrates. Careful analysis of macroinvertebrate communities can reveal condition and trends in aquatic ecosystems. Sampling and analysis is conducted in accordance with procedures outlined in FSH R-4 2609.23, March 1985, Fisheries Habitat Surveys Handbook.

This report is based upon 27 macroinvertebrate samples from 6 stations on North Fork American Fork River and 3 stations on Mary Ellen Gulch Creek along with physical and water chemistry data provided by your aquatic specialist. Samples were taken above and below old mine sites to determine possible effects of mine drainage upon the streams water quality and aquatic life. Zinc concentrations on the North Fork American Fork River ranged from less than 0.02 mg/l at Stations 1, 3A and 8 to 0.190 mg/l at Station 3 below the lower bog mine. Zinc concentrations on Mary Ellen Gulch Creek ranged from 0.022 mg/l at Station 12 above the mine drainage to 0.092 mg/l at Station 14 below the mine effluents.

NORTH FORK AMERICAN FORK RIVER

The Upper Station (1) was above the lower bog mine. Macroinvertebrate samples were also taken at Station 3 just below the lower bog mine to evaluate the effects of mine drainage from the old mine site. Of the heavy metals tested, zinc appeared to be the most potentially damaging heavy metal in the effluents from the mines along the North Fork. At Station 1 above the lower bog mine the zinc concentration was less than 0.02 mg/1, which was a non-limiting concentration. The aduatic macroinvertebrate community had fairly good diversity among clean water taxa at this station which included clean water mayflies Bhithrogens and Enhancella dodds; with fair population numbers. The clean water stoneflies Zanada and members of the family Leuctridae were present in good resident population numbers. These species indicated relatively good substrate and good water quality at the upper Station (1). The zinc concentration was slightly higher when sampled in July at this station.

At Station 3 below the lower bog mine the zinc concentration was 0.190 mg/1, which was higher than found in July when it was 0.077 mg/1. The zinc concentration in September exceeded them threshold value of 0.1 mg/1 for sensitive aquatic invertebrates. A comparison of the communities at the upper Station (1) and the station below the bog mine (3) indicates that there were stress conditions at the lower reach withe number of organisms/m² was reduced 78%, from nearly 9,000 to less than 2,000. The macroinvertebrate standing crop was reduced by nearly 90%, from 0.9 to 0.1 g/m² and the BCI value was reduced from 93 to 79.

None of the taxa at Station 3 had good resident population numbers and many were limited to about 3/m2. Those with the highest numbers were the most tolerant species in the community and apparently. Rhyscaphils which had been fairly comfortable with the sizinc concentration in July was affected by the slightly higher concentration in September since with was no longer a dominant species in the community. The Planaria, which were doing so well in July were missing from the community in September. As found in July samples, the mayflies seem to be the most sensitive to the zinc and some of the shredder stoneflies appeared to be fairly resistent to the effects of this heavy metal.

In contrast, the macroinvertebrate community at Station, 3A, located just above the Pacific mine area, hadran excellent, macroinvertebrate community with good diversity and high population numbers for the clean water species, which indicated good water quality and good instream substrate in that stream reach. There were warning numbers of those taxa tolerant to sedimentation, but good diversity and good resident population numbers for most of the taxa in the community indicated that there was good stability in this stream reach.

Also, the observed number of shredders in the community is generally found where riparian habitat is in good to excellent condition. The zinc concentration at this station was less than 0.02 mg/l. Clean water species present included mayflies ERRORUS, Rhithrogens and Ephemerella dodder, stoneflies Skwala and Zapada and members of the family Leuctridae, and caddisflies Parapaychs and Apatania. These species were also found in the July samples at this station.

When sampled in September, the zinc concentration at Station 8 was pless than 0.02 mg/1. This Station was located below the Pacific mine drainage area. The zinc concentration was much lower than the 0.081 mg/1 found in July. However, this community was limited. Many of the species did not have resident population numbers and the number of organisms was reduced about 70%, to less than 4,000/m², compared to over 13,000/m² at Station 3A. The DAT diversity index value of 15.2 was much better than found in July at this station when it was 0.7. Conditions at this station appear to be somewhat better but were still limiting to the macroinvertebrate community in September.

At the Dutchman Flat Station (9) the zinc concentration was 0.037 mg/l, which was less than the 0.043 mg/l found in July. The DAT diversity index of 18 was much better than the 0.8 found in July, but the numbers of organisms in the community was just about the same and was close to that found at the Upper Station (1). Clean water species at this station had fairly good population numbers and included the mayflies Engages. Rhithcagens and Enhancedla dodds, stoneflies Zanada and members of the family Leuctridae, and caddisflies Arctonagene and Paransuche. There were indications of at least moderate amounts of sedimentation at this station. The observed number of shredders in the community is generally found where riparian habitat is in fairly good condition.

At the lowest station on the North Fork the zinc concentration of 0.099 mg/l was higher than found in July when

it was 0.04 mg/ll and was approaching the 0.1 mg/l threshold value for zincl. However, the macroinvertebrate community there included clean water aspecies, which indicated good water quality and good instream substrate at this station. Clean water mayflies included Enggrue, Rhithrogens and Enhancella doddsi, each found in July samples, and stoneflies Zanada, Amphinemura and members of the family Leuctridae, caddisflies ALSIGNESCHE and Paranageche, most with good population numbers. Good resident population numbers for most of the taxa in the community indicated good stability in this stream reach. The observed number of shredders in the community is generally found where riparian habitat is in good condition.

The potential for a resident fishery on this stream appeared to be good at Stations 3A and 11, where the macroinvertebrate biomass was sufficient to provide nutrients for a good fishery. It appeared that the water chemistry was limiting to biotic life at Stations 3 and 8 and could be limiting to the success of a fishery in those stream reaches. The most sensitive stages in the trout life cycle would probably be limited by the adverse water chemistry. The macroinvertebrate biomass of 0.1 g/m^2 at Station 3 and 0.4 and 0.6 g/m^2 at Stations 8 and 9 respectively, would be limiting to the success of a fishery.

The BCI values at most of the stations indicate there is good potential in this stream. The BCI value of 79 at Station 3 indicated just fair conditions there. There were impacts in that stream reach. It appears there may be some opportunity for management to improve the instream habitat quality and water quality in this aquatic ecosystem.

The effects of the mine drainage from the lower bog mine, appeared to have more limiting effects in September than was observed in July and the effluents from the Pacific mine did not appear to be as limiting in September as they were in July. It appeared that even though there were indications of adverse effects in the mine effluents in the reaches mof stream sampled, mparticularly below the mine sites, the effects on the biotic community were not severe but were limiting. Some of the zinc concentrations were mainly below the mine sites to threshold, levels for sensitive aquatic especies. The zinc concentrations were rather quickly diluted as they entered the main stream from the mine sites and as the water flowed down stream further accretion flows appeared to decrease the effects of the adverse water chemistry.

USFS - INTERMOUNTAIN REGION - ANNUAL PROGRESS REPORT HACROINVERTEBRATE ANALYSIS

Aquatic Ecosystem Analysis Laboratory 105 Page School Brigham Young University Provo, Utah 84602

	٨.	Investig	ator	Paul Skabel	und		
		Forest/D	istrict_	Uinta N.F.			
		Stream		NORTH FORK	AMERICAN FORK RIVER		
		State/Co	unty	Utah,	Utah County		
		Forest S	ervice Ca	t. No			
	В.					Biotic	
				Diversity	Standing	Condition	
Organisms				Index	Çrop	Index	
/m ²	Sta	tion Di	ate(s)	DAT (mean)	r/m² (mean)	BCI 50	#Taxa
981	1 (Abv bog M)	9-21-88	10.2	0.9	93	21
922		Bel bog M)		11.5	0.1	79	21
13,091		Abv Pac M)		19.2	1.4	100	32
3,888		Bel Pac M)		15.2	0.4	100	31
7,819		Dutch Fl)	9-22-88	18.2	0.6	98	32
9,555	11 (Bel MEG)	9-22-88	16.4	2.1	100	_ 25
7,866	1		7-20-88	1.0	1,0	100	_ 20
5,193	3		7-20-88	0.6	0.6	82	25
13,891	3A		7-20-88	1.8	1.8	91	25
2,582	8		7-20-88	0,7	0.7	98	25
8,730	9		7-20-88	0.8	0.8	88	23
18,163	11		7-20-88	1.3	1.3	85	_ 22
		<u>.</u>	١.				
	Scal	THE RESERVE OF THE PARTY OF THE		DAT	Standing crop	BCI	
		ccellent		8 - 26	4.0 - 12.0	above 90	
		ood		1 - 17	1.6 - 4.0	80 - 90	
		ir		6 - 10	0.6 - 1.5	72 - 79	
	Po	or		0 - 5	0.0 - 0.5	below 72	



STATION: 1

NORTH FORK AMERICAN FORK RIVER (ABV BOG MINE), UINTA NF

DATE: 09 21 88

REPL	TOTAL NO. SPECIES	MEAN /SQM	CONFIDENCE (8Ø PERC LL		STANDARD DEVIATION	PERCENT SE OF MEAN	COEFF. OF VARIATION	DBAR	R	CTQA	стар
· NUMB	ERS DATA										
3	21	8981.	4947.	13015.	3704.52	23.81	41.25	2.6345	0.4006	53.	54.

SPECIES TOLERANCE CODES

Clean water species

Moderately tolerant species

Shredders - Depend upon deciduous vegetation from riparian areas)

Sediment tolerant

Organic enrichment tolerant

Resistant to adverse chemistry

Large stonefly species



STATION	: 1	NORTH FORK A	MERICAN FORK RI	VER (ABV BOG M)	NE),	JINTA NF		DATE: 09 21	88	
CLASS	ORDER	FAMILY	GENUS	SPECIES		MEAN NO/SQM	LOG10 NO/SQM	TOLERANCE QUOTIENT	LOGIØ X TQ	MEAN WT
INSECTA	EPHEMEROPTERA EPHEMEROPTERA EPHEMEROPTERA EPHEMEROPTERA	EPHEMERELLIDAE SIPHLONURIDAE	CINYGMULA RHITHROGENA EPHEMERELLA EPHEMERELLA AMELETUS BAETIS TAENIONEMA ZAPADA RHYACOPHILA DICRANOTA HEXATOMA PERICOMA PLANARIA	INERMIS DODDSI ACROPEDES	1/4/2011/01/02/10000000	997.09 107.60 35.87 43.04 7.17 4598.11 43.04 14.35 638.43 487.79 408.88 279.76 7.17 14.35 538.00 14.35 315.63 286.93 78.91 57.39	2.999 2.032 1.654 0.856 3.663 1.637 2.805 2.612 2.447 0.856 1.157 2.731 1.157 2.731 1.157 2.459 2.458 1.897 1.759 0.856	30. 21. 48. 2. 48. 72. 24. 48. 32. 16. 18. 18. 18. 108. 108. 108. 108. 108.	90. 43. 75. 3. 41. 264. 39. 56. 90. 43. 47. 44. 21. 42. 295. 42. 270. 285. 205. 172. 92.	
				TOTALS		8981.01	3.953			0.40

TOTAL SAMPLE STATISTICS

STATION: 3 NORTH FORK AMERICAN FORK RIVER (BEL LOWER BOG MINE), UINTA NF DATE: 69 21 88

REPL	TOTAL NO. SPECIES	MEAN /SQM	CONFIDENCE (80 PERC		STANDARD DEVIATION	PERCENT SE OF MEAN	COEFF. OF VARIATION	DBAR	R	СТДА	CTQD
• NUMB	ERS DATA										
3	21	1922.	923.	2922.	917.59	27.56	47.73	2.8333	0.3588	61.	63.



STATIO	N: 3	NORTH FORK	AMERICAN FORK RI	VER (BEL LOWER	BOG MIN	E), UINT	A NF	DATE: 09 21	88	
CLASS	ORDER	FAMILY	GENUS	SPECIES		MEAN NO/SQM	LOG10 NO/SQM	TOLERANCE QUOTIENT	LOGIØ X TQ	MEAN WT GM/SQM
INSECTA INSECT	EPHEMEROPTERA EPHEMEROPTERA PLECOPTERA PLECOPTERA PLECOPTERA TRICHOPTERA TRICHOPTERA TRICHOPTERA OIPTERA DIPTERA DIPTERA DIPTERA DIPTERA DIPTERA DIPTERA DIPTERA DIPTERA OIPTERA OOFRACODA		EPEORUS RHITHROGENA EPHEMERELLA BAETIS ZAPADA RHYACOPHILA GLOSSOSOMA DICRANOTA PERICOMA HEMERODROMIA	INERMIS	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	3.59 7.17 107.60 21.52 32.28 347.91 46.63 3.59 3.59 3.59 3.59 3.59 3.59 3.59 3.5	0.555 0.856 2.0333 1.509 2.208 1.5509 2.541 1.669 0.5555	21. 21. 48. 72. 24. 32. 16. 18. 24. 108. 24. 108. 95. 36. 95. 108. 95. 108. 95.	12. 18. 98. 96. 36. 71. 24. 46. 30. 13. 60. 60. 13. 53. 60. 53. 25. 20. 53. 25. 26.	
				TOTALS		1022 45	3 294			0.10

TOTAL SAMPLE STATISTICS

STATION: 3A NORTH FORK AMERICAN FORK PIVER (ABV PACIFIC MINE), UINTA NF DATE: 89 22 88

REPL	TOTAL NO. SPECIES	MEAN /SQM	CONFIDENCE (80 PER		STANDARD DEVIATION	PERCENT SE OF MEAN	COEFF. OF VARIATION	DBAR	R	CTQA	стар
· NUMB	ERS DATA										
3	32	13091.	8605.	17578.	4128.41	18.17	31.47	4.0198	0.1962	46.	46.

SPECIES ANALYSES

CLASS	ORDER	FAMILY	GENUS	SPECIES		MEAN NO/SQM	LOG10 NO/SQM	TOLERANCE	LOGIØ X TQ	MEAN WT
INSECTA	EPHEMEROPTERA	HEPTAGENIIDAE	EPEORUS		_	114.77	2.060	21.	43.	
INSECTA	EPHEMEROPTERA	HEPTAGENIIDAE	CINYGMULA		-	1291.20	3.111	30.	93.	
INSECTA	EPHEMEROPTERA		RHITHROGENA		_	1040.13	3.017	21.	63.	
INSECTA			EPHEMERELLA	COLORADENSI		14.35	1.157	18.	21.	
INSECTA		EPHEMERELLIDAE	EPHEMERELLA	INERMIS	5	1707.25	3.232	48.	155.	
INSECTA		EPHEMERELLIDAE	EPHEMERELLA	DODDSI		1032.96	3.014	2.	6.	
INSECTA			BAETIS		5,0	1721.60	3.236	72.	233.	
INSECTA	PLECOPTERA	CHLOROPERLIDAE			-/-	301.28	2.479	24.	59.	
INSECTA	PLECOPTERA	PERLODIDAE	SKWALA	PARALLELA	_	28.69	1.458	18.	26.	
INSECTA	PLECOPTERA	PERLODIDAE	MEGARCYS		-1	43.04	1.634	24.	39.	
INSECTA	PLECOPTERA -	TAENIOPTERYGIDAE	TAENIONEMA		0	315,63	2.499	48.	120.	
INSECTA	PLECOPTERA	CAPNIIDAE			-1	631.25	2.800	32.	90.	
INSECTA	PLECOPTERA	NEMOURIDAE	ZAPADA		-	329.97	2.518	16.	40.	
INSECTA	PLECOPTERA	NEMOURIDAE	MALENKA		5	129.12	2.111	36.	76.	
INSECTA	PLECOPTERA	LEUCTRIDAE			_	272.59	2.436	. 18.	44.	
INSECTA	TRICHOPTERA	HYDROPSYCHIDAE	CHEUMATOPSYCHE		5	344.32	2.537	108.	274.	
INSECTA	TRICHOPTERA	HYDROPSYCHIDAE	PARAPSYCHE		_	559.52	2.748	6.	16.	
INSECTA	TRICHOPTERA	LIMNEPHILIDAE	OLIGOPHLEBODES		_	14.35	1.157	24.	28.	
INSECTA	TRICHOPTERA	LIMNEPHILIDAE	APATANIA		-,	28.69	1.458	18.	26.	
INSECTA	TRICHOPTERA	RHYACOPHILIDAE	RHYACOPHILA		-1, ch	157.81	2.198	18.	40.	
INSECTA	TRICHOPTERA	GLOSSOSOMATIDAE	GLOSSOSOMA		-1	172.16	2.236	24.	54.	
INSECTA	TRICHOPTERA	LEPIDOSTOMATIDAE			-	43.04	1.634	18.	29.	
INSECTA	DIPTERA	TIPULIDAE	DICRANOTA		_	86.08	1.935	24.	46.	
INSECTA	DIPTERA	TIPULIDAE	HEXATOMA		5	14.35	1.157	36.	42.	
INSECTA	DIPTERA	SIMULTIDAE			0	28.69	1.458	108.	157.	
INSECTA	DIPTERA	CHIRONOMIDAE			5,0	1692.91	3.229	108.	349.	
INSECTA	DIPTERA	EMPIDIDAE			5.	71.73	1.856	95.	176.	
INSECTA	DIPTERA	CERATOPOGONIDAE			S. ch	28.69	1.458	108.	157.	
INSECTA	DIPTERA	PSYCHODIDAE	PERICOMA		s, cu	200.85	2.303	36.	83.	
CRUSTACEA	OSTRACODA				5	329.97	2.518	108.	272.	
OLIGOCHAETA					0,3	28.69	1.458	108.	157.	
ARACHNIDA	HYDRACARINA				5,0	315.63	2.499	98.	245.	
				TOTALS		13091.34	4.117			1.48

1

TOTAL SAMPLE STATISTICS

STATION: 8 NORTH FORK AMERICAN FORK RIVER (BEL PACIFIC MINE), UINTA NF DATE: 89 22 88

2501	TOTAL NO.	MEAN	CONFIDENCE (80 PERC		STANDARD	PERCENT SE	COEFF. OF				
REPL	SPECIES	/SQM		OL.	DEVIATION	OF MEAN	VARIATION	DBAR	R	CTQA	CTQD
* NUMB	ERS DATA	and the second							warmen and		
3	31	3888.	2627.	5148.	1157.60	17.19	29.77	3.4025	0.3144	52.	50.

SPECIES ANALYSES

STATIO	N: 8	NORTH FORK	AMERICAN FORK RIVE	ER (BEL PACIFI	C MINE)	, UINTA	NF	DATE: 09 22	88	
CLASS	ORDER	FAMILY	GENUS ·	SPECIES		MEAN NO/SQM	LOG10 NO/SQM	TOLERANCE QUOTIENT	LOCIØ X TQ	MEAN WT GM/SQM
INSECTA	EPHEMEROPTERA EPHEMEROPTERA EPHEMEROPTERA EPHEMEROPTERA EPHEMEROPTERA	HEPTAGENIIDAE HEPTAGENIIDAE HEPTAGENIIDAE EPHEMERELLIDAE EPHEMERELLIDAE BAETIDAE CHLOROPERLIDAE PERLODIDAE TAENIOPTERYGIDAE NEMOURIDAE NEMOURIDAE	EPEDRUS CINYGMULA RHITHROGENA EPHEMERELLA EPHEMERELLA BAETIS MEGARCYS TAENIONEMA ZAPADA MALENKA		111101101111	39.45 39.45 114.77 7.17 86.08 75.32 1477.71 25.11 17.93 89.67 68.15 129.12	1.596 1.596 2.060 0.856 1.935 1.877 3.170 1.400 1.254 1.953 1.833 2.111	21. 30. 21. 18. 48. 2. 72. 24. 24. 24. 32. 16. 36.	34. 48. 43. 15. 93. 4. 228. 34. 36. 94. 59. 37.	
INSECTA INSECT	PLECOPTERA TRICHOPTERA TRICHOPTERA TRICHOPTERA TRICHOPTERA TRICHOPTERA TRICHOPTERA DIPTERA	NEMUNIDAE LEUCTRIDAE HYDROPSYCHIDAE HYDROPSYCHIDAE HYDROPSYCHIDAE RHYACOPHILIDAE GLOSSOSOMATIDAE ELMIDAE TIPULIDAE TIPULIDAE TIPULIDAE TIPULIDAE CHIRONOMIDAE EMPIDIOAE CERATOPOGONIDAE PSYCHODIDAE	CHEUMATOPSYCHE ARCTOPSYCHE ARCTOPSYCHE PARAPSYCHE RHYACOPHILA .GLOSSOSOMA DICRANOTA HEXATOMA PERICOMA	ch, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5,	15/1775 7 0 355	17.93 32.28 161.40 64.56 132.71 261.83 10.76 10.76 7.17 7.17 347.91 469.85 3.59 17.93 10.76 10.76	1.254 1.579 2.748 1.810 2.123 2.418 1.032 1.032 0.056 0.858 2.541 2.675 1.254 1.032 2.134 0.555	18. 18. 18. 6. 18. 24. 104. 108. 24. 36. 108. 95. 108. 108. 108.	37. 23. 163. 40. 11. 38. 58. 107. 111. 21. 31. 274. 254. 60. 45. 111. 111. 209. 60.	
				TOTALS		3887.95	3.590			0.40

14

TOTAL SAMPLE STATISTICS

STATION: 9

NORTH FORK AMERICAN FORK RIVER (DUTCHMAN FLAT), UINTA NF

DATE: 09 22 88

REPL	TOTAL NO. SPECIES	MEAN /SQM	CONFIDENCE (80 PERC		STANDARD DEVIATION	PERCENT SE OF MEAN	COEFF. OF VARIATION	DBAR	R	CTQA	CTQD
• NUMP	ERS DATA										
3	32	7819.	4711.	10926.	2853.86	21.07	36.50	3.5363	0.2929	50.	51.

SPECIES ANALYSES

STATION: 9 NORTH FORK AMERICAN FORK RIVER (DUTCHMAN FLAT), UINTA NF DATE: 09 22 88 MEAN LOG1Ø TOLERANCE LOGIØ X MEAN WT FAMILY CLASS ORDER **GENUS** SPECIES NO/SQM NO/SQM QUOTIENT TQ GM/SQM INSECTA EPHEMEROPTERA HEPTAGENIIDAE **EPEORUS** 114.77 2.060 21. 43. EPHEMEROPTERA HEPTAGENIIDAE CINYGMULA INSECTA 161.48 2.208 30. 66. INSECTA EPHEMEROPTERA HEPTAGENIIDAE RHITHROGENA 667.12 2.824 21. 59 EPHEMEROPTERA EPHEMERELLIDAE EPHEMERELLA INERMIS INSECTA 86.08 1.935 48. 93. INSECTA EPHEMEROPTERA EPHEMERELLIDAE **EPHEMERELLA** DODDSI 186.51 2.271 2. 5 INSECTA EPHEMEROPTERA SIPHLONURIDAE **AMELETUS** 28.69 1.458 48. 70. 5,0 INSECTA EPHEMEROPTERA BAETIDAE BAETIS 2403.07 3.381 72. 243. INSECTA PLECOPTERA 14.35 1.157 48. 56. CHLOROPERLIDAE PLECOPTERA INSECTA 269.00 2.430 24. 58. PLECOPTERA INSECTA PERLODIDAE MEGARCYS 21.52 1.333 24. 32. TAENIOPTERYGIDAE TAENIONEMA 0 INSECTA PLECOPTERA 964.81 2.984 48. 143. INSECTA CAPNIIDAE PLECOPTERA 418.05 2.619 32. 84 NEMOURIDAE ZAPADA INSECTA PLECOPTERA 254.65 2.496 16. 38. INSECTA PLECOPTERA LEUCTRIDAE 53.80 1.731 18 31 INSECTA **PLECOPTERA** PERLODIDAE **ISOGENDIDES** 17.93 1.254 24. 30. INSECTA TRICHOPTERA HYDROPSYCHIDAE CHEUMATOPSYCHE 39.45 1.596 108. 172. HYDROPSYCHIDAE INSECTA TRICHOPTERA ARCTOPSYCHE 57.39 1.759 18. 32. INSECTA TRICHOPTERA HYDROPSYCHIDAE PARAPSYCHE 39.45 1.596 6. 10 INSECTA TRICHOPTERA RHYACOPHILIDAE RHYACOPHILA 28.69 1.458 18. 26. INSECTA TRICHOPTERA GLOSSOSOMATIDAE GLOSSOSOMA 60.97 1.785 24. 43. TRICHOPTERA INSECTA RHYACOPHILIDAE RHYACOPHILA ACROPEDES C 14.35 1.157 18. 21. INSECTA COLEOPTERA ELMIDAE 57.39 1.759 104 183 INSECTA DIPTERA 154.23 2.188 108. 236. INSECTA DIPTERA TIPULIDAE DICRANOTA 86.08 1.935 24. 46 INSECTA DIPTERA TIPULIDAE HEXATOMA 10.76 1.032 36. 37. 0 SIMULIIDAE INSECTA DIPTERA 93.25 1.970 108. 213. 5,0 1169.25 INSECTA DIPTERA CHIRONOMIDAE 3 068 108. 331. INSECTA EMPIDIDAE DIPTERA 50.21 1.701 95. 162. =. ch INSECTA DIPTERA CERATOPOGONIDAE 10.76 1.032 108. 111. INSECTA DIPTERA PSYCHODIDAE PERICOMA s, ch 17.93 1.254 36. 45. CRUSTACEA OSTRACODA 68.15 1.833 108. 5 198. ARACHNIDA HYDRACARINA 200.85 2.303 98. 226. 5,0

TOTALS

7818.93

3.893

0.60

-

MARY ELLEN GULCH CREEK

At the Upper Station (12) above the mine, clean water species included mayflies Bhithrogens and Enemeralla dodds with fairly good population numbers, and the stonefly Zanada with excellent population numbers, which indicated good water quality and some good instream substrate in that stream reach. The observed number of shredders in the community is generally found where riparian habitat is in good condition. The zinc concentration there was just 0.022 mg/l, which was below the threshold value. Good diversity and resident population numbers for many of the taxa in the community at Station 12 indicated good stability in that stream reach.

Of the stations sampled, Station 14 below Mary Ellen Gulch Creek Mine showed the most severe impacts. All of the analysis elements indicated that there were severe impacts at this station. At the control station above the mine effluents the DAT was 11.7; at Station 14 below the mine effluents it was 1.9. The standing crop had decreased 83% from 2.3 to 0.4 g/m², the BCI value from 89 to 78, and the number of taxa from 20 to 15. The only species with a resident population number at Station 14 was the adverse water chemistry and sediment tolerant Chiconomids, which numbered far less than was found in July. Most of the other taxa in the community had extremely limited numbers in their populations and were not living successfully in the reach sampled. The zinc concentration of 0.092 mg/1 was close to the threshold value for sensitive aquatic species. There were 6,500 organisms/m², but 5,200 were Chiconomids.

At the mouth of Mary Ellen Creek (Station 10) near its confluence with North Fork of the American Fork River the macroinvertebrate community showed that the ecosystem had recovered considerably by the time it reached that point. Clean water species present at that station included mayflies Epachus, Bhithrogena and Ephemerella doddai, stoneflies Zanada and members of the family Leuctridae, and caddisflies Accionates and Parapsyche, which indicated relatively good water quality in that stream reach and that the toxic effects of the zinc and other possible heavy metals that may synergistically have been operating at Station 14, no longer had a serious detrimental effect in the aquatic ecosystem. The zinc concentration at Station 10 was 0.041 mg/l which was below the threshold value and was lower than found in July. The same clean water species were present at tation 10 as were found at

the Control Station (12) above the mine. These clean water taxa were not present at the station directly below the mine.

The potential for a resident fishery on this stream appeared to be fairly good in the reaches of stream sampled above the mine and at the mouth. However, it appeared the fishery would be extremely limited at Station 14 below the mine. Scarcity of clean water taxa in the community and abundance of sediment tolerant taxa, particularly the Chironomids, indicated there would be a very limited amount of suitable spawning substrate. Also, the water chemistry would probably not be compatible with the most sensitive life stages in the life cycle of a trout. The macroinvertebrate biomass at Stations 10 and 12 would be sufficient to provide nutrients for a fishery, but 0.4 g/m² at Station 14 would be limiting to a fishery.

A BCI value of 100 at Station 10 indicated that this reach of stream was close to its potential; the BCI value of 89 at Station 12 indicated good conditions at the upper station, and a BCI of 78 at Station 14 indicated just fair conditions in that stream reach and that was based on a community where most species did not have resident population numbers.

September samples were expected to show more severe, effects from the mine drainages on these streams than were observed in July. This was true at Stations 3 and 14, however, in general it appeared that distant, down-stream effects were less than those found in July. It appears that the sediment and heavy metals, particularly the zinc, entering these aquatic ecosystems is detrimental to the aquatic life in the vicinity of the effluents from the old mines. As accretion flows dilute the heavy metals down-stream reaches appear to become more and more capable of supporting aquatic life. It appears there would be opportunities for management to improve the instream habitate quality and water quality below each of the mine sites monitored in 1988 along the North Fork American Fork River and Mary Ellen Creek ecosystems.

USFS - INTERMOUNTAIN REGION - ANNUAL PROGRESS REPORT HACROINVERTEBRATE ANALYSIS

Aquatic Ecosystem Analysis Laboratory 105 Page School Brigham Young University Provo, Utah 84602

A. Investigator_____Paul Skabelund

	Forest	/District	Uinta N.F.			
	Stream	100	MARY ELLEN	CREEK		
	State/0	County	Utah,	Utah County		
	Forest	Service (Cat. No			
rganisms	B. Station	Date(s)	Diversity Index DAT (mean)	Standing Crop g/m ² (mean)	Biotic Condition Index BCI 50	
8,013 6,685 6,528	10 (Mouth) 12 (Ab M) 14 (Bel M)		15.2 11.7 1.9	1.4 2.3 0.4	100 89 78	#1
13,884 12,424 30,110	10 12 14	7-20-88 7-21-88 7-21-88	7.8 11.6 1.2	2.1 2.1 2.0		
	=======================================	•	DAT	Shanding and	BCI	
	Excellent Good Fair Poor		DAT 18 - 26 11 - 17 6 - 10 0 - 5	Standing grop 4.0 - 12.0 1.6 - 4.0 0.6 - 1.5 0.0 - 0.5	above 90 80 - 90 72 - 79 below 72	

TOTAL SAMPLE STATISTICS

STATION: 11

AMERICAN FORK RIVER, UINTA NF

DATE: 09 22 88

REPL	TOTAL NO. SPECIES	MEAN /SQM	CONFIDENCE (80 PERC		STANDARD DEVIATION	PERCENT SE OF MEAN	COEFF. OF VARIATION	DBAR	R	CTQA	стар
* NUMB	ERS DATA										
3	25	9555.	7238.	11871.	2127.33	12.85	22.26	3.7368	0.1959	49.	47.

SPECIES ANALYSES

STATION: 11

AMERICAN FORK RIVER, UINTA NF

DATE: 09 22 88

CLASS	ORDER	FAMILY	GENUS	SPE	CIES	MEAN NO/SQM	LOG10 NO/SQM	TOLERANCE QUOTIENT	LOGIØ X TQ	MEAN WT
INSECTA INSECT	EPHEMEROPTERA EPHEMEROPTERA EPHEMEROPTERA EPHEMEROPTERA PLECOPTERA DIPTERA	HEPTAGENIIDAE HEPTAGENIIDAE HEPTAGENIIDAE HEPTAGENIIDAE BAETIDAE BAETIDAE CHLOROPERLIDAE PERLODIDAE TAENIOPTERYGIDAE CAPNIIDAE NEMOURIDAE NEMOURIDAE HYDROPSYCHIDAE HYDROPSYCHIDAE HYDROPSYCHIDAE TIPULIDAE TIPULIDAE CHIRONOMIDAE EMPIDIDAE BLEPHARICERIDAE	EPEORUS CINYGMULA RHITHROGENA EPHEMERELLA BAETIS MEGARCYS TAENIONEMA ZAPADA AMPHINEMURA ARCTOPSYCHE PARAPSYCHE RHYACOPHILA DICRANOTA PLANARIA	DODDSI	17110 1107111111 TOUNING	243.89 286.93 1004.27 157.81 2395.89 387.36 14.35 789.07 731.88 157.81 14.35 88.08 487.79 301.28 57.39 57.39 67.39 632.11 387.36 172.16 122.8.69 28.69 28.69 100.43 14.35	2.387 2.458 3.0098 3.379 2.6887 2.8864 2.198 2.198 1.1535 2.688 2.4759 1.7520 2.588 2.2588 2.191 1.458 2.459 2.111 1.458 1.458 2.197	21. 30. 21. 2. 72. 24. 48. 32. 16. 6. 18. 18. 18. 24. 198. 24. 198. 24. 198. 24. 198. 24. 198. 24. 198. 24. 198. 24. 24. 26. 27. 28. 28. 28. 28. 28. 28. 28. 28	50. 74. 63. 4. 243. 62. 28. 139. 92. 45. 13. 211. 35. 16. 45. 190. 42. 316. 280. 212. 46. 157. 167. 196.	
				тот	ALS	9554.88	3.980			2.10

N

TOTAL SAMPLE STATISTICS

STATION: 10

MARY ELLEN CREEK (MOUTH), UINTA NF

DATE: 09 22 88

REPL	TOTAL NO.	MEAN	CONFIDENC (80 PER		STANDARD	PERCENT SE	COEFF. OF				
MELL	SPECIES	/SQM	LL	UL	DEVIATION	OF MEAN	VARIATION	DBAR	R	CTQA	CTQD
• NUMB	ERS DATA										
3	25	8013.	4399.	11626.	3318.29	23.91	41.41	3.6554	0.2135	45.	43.

SPECIES ANALYSES

STATION: 10 MARY ELLEN CREEK (MOUTH), UINTA NF

DATE: 09 22 88

CLASS	ORDER	FAMILY	GENUS	SPECI	ES	MEAN NO/SQM	LOGIØ NO/SQM	TOLERANCE QUOTIENT	LOG1Ø X TQ	MEAN WT GM/SQM	
INSECTA INSECT	EPHEMEROPTERA EPHEMEROPTERA EPHEMEROPTERA EPHEMEROPTERA EPHEMEROPTERA PLECOPTERA PLECOPTERA PLECOPTERA PLECOPTERA PLECOPTERA PLECOPTERA TICHOPTERA TRICHOPTERA TRICHOPTERA TRICHOPTERA TRICHOPTERA TRICHOPTERA TRICHOPTERA DIPTERA DIPTERA DIPTERA DIPTERA DIPTERA TRICLADIDA HYDRACARINA	HEPTAGENIIDAE HEPTAGENIIDAE HEPTAGENIIDAE HEPTAGENIIDAE EPHEMERELLIDAE EPHEMERELLIDAE BAETIDAE CHLOROPERLIDAE PERLODIDAE TAENIOPTERYGIDAE CAPNIIDAE NEMOURIDAE LEUCTRIDAE HYDROPSYCHIDAE HYDROPSYCHIDAE HYDROPSYCHIDAE RHYACOPHILIDAE GLOSSOSOMATIDAE TIPULIDAE CHIRONOMIDAE EMPTOIDAE PELECORHYNCHIDAE PLANARIIDAE	EPEORUS CINYGMULA RHITHROGENA EPHEMERELLA EPHEMERELLA BAETIS MEGARCYS TAENIONEMA ZAPADA ARCTOPSYCHE PARAPSYCHE RHYACOPHILA GLOSSOSOMA DICRANOTA GLUTOPS PLANARIA	INERMIS DODDSI	11/10/10/11/11/11/00 00 N	168.57 1104.69 642.01 89.67 243.89 1979.84 43.04 885.91 25.11 204.44 613.32 545.17 344.32 14.35 157.81 154.23 143.47 1076 68.15 218.79 60.97 14.35 200.85 53.80 25.11	2.227 3.043 2.808 1.953 2.387 3.297 1.634 2.731 2.737 2.537 1.157 2.188 2.167 1.032 1.785 1.157 2.303 1.785 1.167	21. 30. 21. 48. 2. 72. 48. 24. 24. 24. 16. 18. 72. 18. 6. 18. 24. 24. 24. 29. 18. 6. 18. 95. 95. 96. 198. 97. 98. 98. 99. 99. 99. 99. 99. 99	47. 91. 59. 94. 5. 237. 78. 71. 34. 111. 89. 44. 46. 83. 40. 13. 39. 25. 44. 253. 170. 35. 249. 170. 180. 190. 1		
				TOTAL		8012.61	3.904			1.40	

2

TOTAL SAMPLE STATISTICS

STATION: 12

MARY ELLEN CREEK, UINTA NF

DATE: 09 22 88

	TOTAL NO.	MEAN	CONFIDENCE (8Ø PERC		STANDARD	PERCENT SE	COEFF. OF				
REPL	SPECIES	/SQM	LL	UL	DEVIATION	OF MEAN	VARIATION	DBAR	R	CTQA	CTQD
* NUMB	ERS DATA										
3	28	26685.	13503.	39866.	12105.49	26.19	45.36	3.0992	0.2833	57.	56.



STATION: 12

MARY ELLEN CREEK, UINTA NF

DATE: 09 22 88

CLASS	ORDER	FAMILY	GENUS	SPEC	IES	MEAN NO/SQM	LOG10 NO/SQM	TOLERANCE QUOTIENT	LOGIØ X	MEAN WT
INSECTA INSECT	EPHEMEROPTERA EPHEMEROPTERA EPHEMEROPTERA PLECOPTERA PLECOPTERA PLECOPTERA PLECOPTERA PLECOPTERA TRICHOPTERA TRICHOPTERA DIPTERA DIPTERA DIPTERA DIPTERA OIPTERA OIPTERA OSTRACODA	HEPTAGENIIDAE EPHEMERELLIDAE EPHEMERELLIDAE	CINYGMULA RHITHROGENA EPHEMERELLA EPHEMERELLA AMELETUS BAETIS MEGARCYS TAENIONEMA ZAPADA CHEUMATOPSYCHE RHYACOPHILA DICRANOTA	INERMIS	[14110] 140] 140 10 10 10 10 10 10 10 10 10 10 10 10 10	5910.83 114.77 2266.77 86.08 315.63 946.86 86.08 200.85 28.69 1291.20 2410.24 315.63 1778.99 172.16 5652.59 28.69 4791.79 28.69 229.56	3.772 2.060 3.355 2.499 2.976 2.303 1.458 3.111 3.382 2.499 3.250 3.752 1.458 3.680 1.458 2.361	30. 21. 48. 2. 48. 72. 24. 24. 16. 108. 18. 95. 108. 108. 108. 95. 108.	113. 43. 161. 4. 120. 214. 46. 55. 70. 100. 54. 270. 59. 54. 405. 138. 157. 397. 167. 231.	2.30

TOTAL SAMPLE STATISTICS

STATION: 14

MARY ELLEN CREEK, UINTA NF

DATE: 69 22 8

REPL	TOTAL NO. SPECIES .	MEAN /SQM	CONFIDENCE (80 PERC LL		STANDARD DEVIATION	PERCENT SE OF MEAN	COEFF. OF VARIATION	DBAR	R	CTQA	стар
* NUMB	ERS DATA										
3	15	6528.	4880.	8176.	1513.35	13.38	23.18	1.3199	0.6639	62.	64.



STATION: 14 MARY ELLEN CREEK, UINTA NF

CLASS	ORDER	FAMILY	GENUS	SPECIES	MEAN NO/SQM	LOG1Ø NO/SQM	TOLERANCE QUOTIENT	LOGIØ X	MEAN WT GM/SQM
INSECTA	EPHEMEROPTERA EPHEMEROPTERA PLECOPTERA PLECOPTERA PLECOPTERA TRICHOPTERA DIPTERA DIPTERA DIPTERA DIPTERA DIPTERA TRICHOPTERA DIPTERA DIPTERA TRICHOPTERA DIPTERA DIPTERA DIPTERA TRICHODIA	HEPTAGENIIDAE SIPHLONURIDAE BAETIDAE CHLOROPERLIDAE CAPNIIDAE NEMOURIDAE RHYACOPHILIDAE TIPULIDAE SIMULIIDAE CHIRONOMIDAE EMPIDIDAE CERATOPOGONIDAE PLANARIIDAE	EPEORUS AMELETUS BAETIS ZAPADA RHYACOPHILA DICRANOTA	ACROPEDES TICK Ch, 5, 5, 0	14.35 14.35 14.35 100.43 100.43 100.43 401.71 57.39 28.69 5236.53 14.35 258.24 28.69 57.39	1.157 1.157 2.157 1.157 2.002 1.769 2.002 2.604 1.759 1.458 3.719 1.157 2.412 1.458 1.759	21. 48. 72. 48. 24. 32. 16. 18. 24. 108. 108. 108. 108. 108.	24. 56. 155. 56. 48. 56. 32. 47. 42. 157. 402. 118. 268. 157. 172.	
				TOTALS	6527.73	3.815			0.40

PRELIMINARY SURVEY OF WATER QUALITY

IN MINE DRAINAGE IN SHEEPROCK MOUNTAINS AND

NORTH FORK OF THE AMERICAN FORK RIVER

FOR

UINTA NATIONAL FOREST

BY

AVERE B. MERRITT, Ph.D., P.E. Environmental Engineer

> Provo, Utah July 1988

me should

o to one that ever as

Introduction

As part of an abandoned/inactive mine survey, several mines in the Sheeprock Mountains and in the American Fork River drainage were visited and water samples taken from mine drainage waters and nearby natural drainage streams on May 12 and May 18, 1988 respectively.

This survey was intended to help identify the locations and water quality parameters that would need more intensive sampling and evaluation later.

Quality Comparison Basis

Although heavy metal standards for quality for a cold water sports fishery would need to be somewhat more stringent than for drinking water, in this survey phase drinking water standards are used for comparison. Macroinvertebrate samples will then be used in selected areas to indicate the nature of the ecosystem stresses and then heavy metal conclusions drawn from those and additional water sampling results.

(a) The maximum contaminant level for nitrate is applicable to both community water systems and non-community water systems. The levels for the other inorganic chemicals apply only to community water systems. Compliance with naximum contaminant levels for inorganic chemicals is calculated pursuant to 6 141,23.

(b) The following are the maximum contaminant levels for inorganic chemier than fluoride:

Contaminant	Level, milligrams per liter
Arsenic Barium Cadmium Chromium Lead Mercury Nitrate (as N) Selenium	 1. 0.010 0.05 0.05 0.002 10. 0.01

. (c) When the annual average of the maximum daily air temperatures for the location in which the community water system is situated is the following, the maximum contaminant levels for fluoride

Temperalme Degrees Fahrenheit	Degrees Celsius	Level, 4 milliersms per liter
52.9 to 54-2	17.0 and below 12.1 to 14.6 14.7 to 17.6 17.7 to 21.4 21.5 to 26.2 26.3 to 32.5	2.4 2.2 2.0 1.8 1.6 1.4

§ 141.11 Maximum contaminant levels EPA Regulations

§ 141.12 Maximum contaminant levels for organic chemicals.

The following are the maximum contaminant levels for organic chemicals. They apply, only to community water systems. Compliance with maximum contaminant levels for organic chemicals is calculated pursuant to § 141.24.

Level milligrams per liter (a) Chiorinated hydrocarbons:
Endrin (1,2,3,4,10, 10-hexachioro8,7-epoxy-1,4, 4s,5,6,7,8,8-octahydro-1,4-endo, endo-5,6 - dimethano naphthalene).

Lindane (1.2.3.4.5.6-herachloro- 0.004 cyclohexane, gamma isomer). (1,1,1-Trichloro- 0.1 Methoxychlor 2, 2 - bis [p-methoxyphenyl]

ethane). (C, H, Cl, Technical 0.005 Toxaphene chlorinated camphene, 67-09 percent chiorine).

Secondary Maximum Contaminant Levels for public water systems are:

Contaminant	SMCL
Chloride	250 mg/L
Color	
Copper	1 mg/L
Corrosivity	Noncorrosive
Foaming agents	0.5 mg/L
Iron	0.3 mg/L
Manganese	0.05 mg/L
Odor	3(TON)
pH	6.5-8.5
Sulfate	250 mg/L
Total dissolved solids	500 mg/L
Zinc	5 mg/L

These levels represent reasonable goals for drinking jater quality. The states may establish higher or lower levels which may be appropriate dependent upon local conditions such as unavailability of alternate source waters or other compelling factors, provided that public health and welfare are not adversely affected.

I. Sheeprock Mountains

A. <u>HARKER MINE</u> in the Harker Creek Drainage of the Sheeprock Mountains near Vernon, Utah.

East Portal (probably evaporatory shaft) was flowing about 0.1 cfs of clear water. The test results indicate the water to be of good quality with no heavy metal concentrations of concern, with only As (Arsenic) (2.5 ug/l) and Ba (Barium) (27 ug/l) above detection limits but both far below maximum allowed levels.

South Portal was flowing about 0.2 cfs of water, some signs of chemical instability in orangish precipitates and/or algae in pooled water at portal. Detectable levels of Cd (Cadmium), Pb (Lead), Ba (Barium), Fe (Iron), Mn (Manganese), and Zn (Zinc) were present. Lead at 585 ug/l and Zinc at 2700 ug/l are of some concern along with a pH of

Conclusions These mine drainage waters are normally the main part if not the total flow of small Harker Creek this high in the drainage. During the late summer the stream is probably dry in spots down the stream below the mines. The aquatic habitat in these upper waters is naturally stressed (sediments, high temperatures, no flow) and not capable of supporting a balanced aquatic ecosystem including fish. When these mine drainage waters do flow into the lower reaches of the canyon drainage, mineral precipitation and dilution would make the relatively small amount of Pb and Zn of little concern.

Recommendation No action to be taken with the possible exception of piping the South Portal flow down past the spoils pile, a distance of perhaps 150 to 200 feet--a low-priority project in my opinion.


B. NORTH OAK BRUSH MINE in the North Oak Brush drainage of the Sheeprock Mountains near Vernon, Utah.

<u>East Portal</u> was the only portal observed to have portal drainage waters. The flow was approximately 0.1 cfs of clear water. As, Cd, Cu (Copper), Pb, Ba, Fe, Mn, and Zn were above detection limits but all rather low except Pb at 115 ug/l, Cd at 8 ug/l and Zn at 1200 ug/l which are still moderate.

The Creek was sampled about one-fourth mile below the mine. It was flowing about 0.3 cfs at this point. All detected metals in the mine drainage were at considerably lower levels at this point and none higher than drinking water standards.

<u>Conclusions</u> This mine drainage makes up a large part of the Creek flow this high in the drainage. The flow downstream is likely intermittent seasonally. The aquatic ecosystem is naturally stressed and not capable of supporting fish. The metals from the mine drainage are rather small quantities.

Recommendations - no action.



II. North Fork of American Fork River above American Fork, Utah.

A. LOWER BOG MINE approximately 2 miles upstream of the Pacific Mine on east side of creek.

The portal drainage flow was about 0.1 cfs and the portal is covered by rubble. This is an Acid drainage of pH <4.5. Yellow precipitates are present. The flow percolates into the soil over some 200 feet and is not a surface flow into the creek (but is likely commingling with other percolating waters and seeping into the creek). Detectable levels of As, Cd, Ba, Fe, Mn and Zn are present but at fairly low levels with only Cd at 12 ug/l above DW standards.

<u>Conclusions</u> It was surprising that this acid drainage didn't contain higher levels of heavy metals. The fact that it percolates on into the stream undoubtedly mitigates its impact.


<u>Recommendation</u> - The relatively small percolating flow suggests no action on this mine drainage. However additional water quality samples and macroinvertebrate in the streamflow above and below the area during low stream in summer is desirable.

B. PACIFIC MINE

At Portal A drainage flow of about 0.2 cfs is not acidic. Detectable levels of most heavy metals were present but only As, Cu, and Pb are at significant levels at about one half of DW standards. As the flow continues on toward the stream, generally across spoil material, about 1/4 mile away it picks up metals and at the stream considerably higher levels are found with particular concern focusing on Pb at 4000 ug/l about 100 times DW standards of 50 ug/l. On the sampling day, a drizzling rain was causing a small runoff from the spoils/tailings; runoff flow of about 0.2 cfs was sampled at the bottom of the old spoils lagoon area near the stream. This sample gave by far the highest levels of heavy metals and As, Cd, & Pb were above DW standards with Pb by far the highest at 20,000 ug/l about 400 times the standard.

<u>Upper Portal</u> (NW Portal) A small mine drainage flow is piped from the portal and discharged a short distance downhill. The water is of high quality except Pb at 60 ug/l which is just above DW standards of 50 ug/l.

American Fork River A water sample from the stream (American Fork River) about 1 mile downstream contained some surface runoff and eroded sediment (light) on the day sampled. Quality was very good overall with only Pb at 60 ug/l of concern. This indicates that he upstream mine drainage was having some effect on the stream but the net result on the water quality was moderate to nie. Since considerable amounts of heavy metals are likely precipitating in the stream, macroinvertebrate samples are needed to assess the impact.



<u>Conclusions</u>. The mine drainage waters are not a serious concern at the portals although they do carry slightly high levels of some heavy metals. The real problem is the spoils/tailings. These need to be treated/stabilized and drainage waters routed around the tailings. I am in full agreement with Ben Albrechtsen in his July 1985 file report.

Recommendations. Additional water quality samples should be taken in the stream above and below the site. This should be complemented by macroinvertebrate samples and habitat surveys.

[M, | ler Tunnel]

Lower Pacific Mine just across the stream from the junction in the road (Baker Junction) and about 1/2 mile downstream from Pacific Mine. The drainage water of about 0.1 cfs is of very good quality and shows essentially no heavy metals.

Recommendations. No action except that local runoff from the spoils piles go directly into the stream and the stream is undercutting the toe of the pile. This does not affect the mine drainage water.

C. MARY ELLEN MINE AREA

Mary Ellen Mine drainage flow of about 0.3 cfs at portal contained detectable levels of As, Cd, Cu, Pb, Ba, Fe, Mn and Zn, but only As at 100 ug/l was above the 50 ug/l DW standard. The flow had a pH of _______ which is slightly acidic. The "yellow boy" precipitates in the flow is in concert with the low pH. Some other surface waters in the area give indication of low pH--yellow precipitates. The sample on Mary Ellen Creek about 1/4 mile below the mine had detectable levels of most of the same metals but none exceeded DW standards although Pb was 4 times higher at 40 ug/l, likely indicating the impact of surface drainage leaching from the spoils areas upstream.

<u>Conclusions</u>. Given the rather large areas of spoils/tailings the effect on the stream water quality was less than expected although the rain ceased about 2 hours earlier and surface wash had diminished compared to the Pacific Mine area samples.

<u>Recommendations</u>. Additional water quality and macroinvertebrate samples should be taken during summer lower flow conditions.

Recommended July 88 Sampling Site Water Quality, Lower Bog Water Quality, As. Cd. Cu. Pb. Hg. Ag. Ba. Cr. Fe, Mn, Se, Zn. Pacific Mary Ellen

Schematic Site Map Lower Bog 100-400 yds below Lower Bag V, River Pacific NW Pac Portal DX Pac Portal 5 100-400 gds below tailings pond Dutchman
Flat x 6
At culvert
'early's sil (early site) F. at Mary Ellen M.E. Portal & @ 100-400 yds D 100-400 gds below M.E. tacking 5 Buth River Confluence
Am FK R. gds below confluence

APPENDIX

Water Sample Testing Results

HARKER MINE EAST PORTAL UINTA NAT. FOREST ATN. PAUL P.O.BOX 829 PROVO UT

. 377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: HARKER MINE EAST PORTAL Source: 00 Site ID:

Cost Code: 350B

Lab Number: 8802698 Type: 04

Sample Date: 88/05/12 Fime: 10:10

Tot. Cations:

Anions: nd lotal:

50 me/l Cations:

50 me/l Anions:

Date of Review and QA Validation Inorganic Review: 88/06/09

Organic Review:

Radiochemistry Review: 1.7 Microbiology Review:

fot. Alk.	84	mg/l	rps @ 180C	150	mg/1
1-Arsenic	2.5	ug/l	1-Barium	0.027	mg/l
I'-Cadmium	<1	ug/1	l'-Chromium	(5.0	ug/l
1-Copper	<20.0	ug/l	1-lron	<0.02	mg/1
I'-Lead	(5.0	ug/l	l'-Manganes	(5.0	ug/1
Mercury	<0.2	ug/l	T-Selenium	<0.5	ug/l
I-Silver	(2.0	ug/l	l-Zinc	<20.0	ug/1

Description: HARKER MINE SOUTH PORTAL Site 10: Source: 00

Site 1D: Source: 00

Cost Code: 3508

Lab Number: 8802697 Type: 04

Sample Date: 88/05/12 Fime: 10:30

Cations:
Anions: 17 me/l Cations:
rand lotal: 17 me/l Anions:

Date of Review and QA Validation Inorganic Review: 88/06/09 Organic Review: Radiochemistry Review:

0.6 Microbiology Review:

l'ot. Alk.	28 mg/l	FDS @ 180C	100 mg/l
1-Arsenic	<1.0 ug/l	T-Barium	0.028 mg/l
I'-Cadmium	#16 ug/1:	I-Chromium	<5.0 ug/1
1-Copper	<20.0 ug/1	T-Iron	0.76 mg/l
I'-Lead	585:0 ug/1"	I'-Manganes	420.0 ug/l
Mercury	(0.2 ug/l	T-Selenium	<0.5 ug/l
I-Silver	<2.0 ug/1	ſ-∠inc	2700.0 ug/1

HARKER NAT. CREEK PARALLEL TO SOUTH MINE UINTA NAT. FOREST ATN. PAUL P.O.BOX 829 PROVO

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: HARKER NAT. CREEK PARALLEL TO SOUTH MINE

Site 1D:

Cost Code: 3508

Lab Number: 8802699 Type: 04

Source: 00

Date of Review and QA Validation Inorganic Review: 88/06/09

Sample Date: 88/05/12 lime: lot. Cations:

and Total:

28 me/l Cations:

28 me/l Anions:

Organic Review: Radiochemistry Review: 0.9 Microbiology Review:

Laboratory Analyses

Tot. Alk.

47 mg/l

LD2 @ 180C

82 mg/l

NORTH OAK BRUSH EAST PORTAL UINTA NAT. FOREST ATN. PAUL P.O.BOX 829 PROVO UT

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: NORTH OAK BRUSH EAST PORTAL

Site 10:

Source: 00

Cost Code: 3508 Lab Number: 8802696 Type: 04

Date of Review and QA Validation

Sample Date: 88/05/12 Time: 13:30

Inorganic Review: 88/06/09

lot. Cations:

Organic Review:

Tot. Anions: and lotal:

31 me/1 Cations:

Radiochemistry Review:

31 me/l Anions:

1.0 Microbiology Review:

Fot. Alk.	52 mg/1	FDS @ 180C	124	mg/l
1-Arsenic	8.5 ug/l	T-Barium	0.011	7 24
I'-Cadmium	1:8 ug/1	l'-Chromium	(5.0	ug/l
T-Copper	(41:0: ug/l)	T-Iron	5.5	mg/l
T-Lead	115.0 ug/1.	l'-Manganes	83.0	ug/l
Mercury	<0.2 ug/1	T-Selenium	(0.5	ug/l
T-Silver	<2.0 ug/1	ľ-Zinc	¥1200.0	ug/1-

NORTH OAK BRUSH STREAM 1/4 MILE BL MINE UINTA NAT. FOREST ATN. PAUL P.O.BOX 829 PROVO UT

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: NORTH OAK BRUSH STREAM 1/4 MILE BL MINE

Site 1D:

Source: 00

Cost Code: 350B Lab Number: 8802695 Type: 04 Sample Date: 88/05/12 Time: 14:45

Date of Review and QA Validation Inorganic Review: 88/06/09

lot. Cations:

t. Anions: nd lotal:

41 me/l Cations: 41 me/l Anions:

Organic Review: Radiochemistry Review: 1.4 Microbiology Review:

Tot. Alk.	68	mg/1	TDS @ 180C	118	mg/l
T-Arsenic	<1.0	ug/l	T-Barium	0.017	mg/l
I-Cadmium	1	ug/1	I-Chromium	(5.0	ug/l
1-Copper	(20.0	ug/l	T-lron	1.2	mg/1
I-Lead	40.0	ug/1	l'-Manganes	160.0	ug/l
Mercury	<0.2	ug/l	1-Selenium	(0.5	ug/l
I-Silver		ug/1	ľ-∠inc	80.0	ug/1

PORTAL LOWER BOG MINE UINTA NATIONAL FOREST 88 W 100 N U1 84603 PROVO

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PORTAL LOWER BOG MINE Site 10: Source: 00

Cost Code: 3508 Lab Number: 8802857 Type: 04

Sample Date: 88/05/18 Fime: 12:30

lot. Cations:

t. Anions: and lotal: me/1 Cations: me/l Anions: Date of Keview and QA Validation

Inorganic Review: Organic Keview:

Radiochemistry Review: Microbiology Keview:

0 mg/1	rps @ 1800	90 mg/1
1.5 ug/l	1-Barium	0.037 mg/1
112 ug/1'.	I-Chromium	(5.0 ug/1
(20:0 ug/1	'I-lron	7.9 mg/1
(5.0 ug/1	l'-Manganes	270.0 ug/1
(0.2 ug/1	7-Selenium	(0.5 ug/1
<2.0 ug/1	ľ-Zinc	510.0 ug/1
	1.5 ug/1 112 ug/1 . <20.0 ug/1 <5.0 ug/1 <0.2 ug/1	1.5 ug/l 1-Barium 12 ug/l 1-Chromium (20.0 ug/l 1-lron (5.0 ug/l 1-Manganes (0.2 ug/l 1-Selenium

OUTLET PACIFIC MINE MAIN PORTAL AT AULT UINTA NATIONAL FOREST 88 W 100 N PROVO UI 84603

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PACIFIC MINE MAIN PORTAL AT AUET

Site 1D: Cost Code: 3508 Source: 00

Lab Number: 8802854 Type: 04 Date of Keview and QA Validation

Sample Date: 88/05/18 Time: 10:00 Inorganic Review: lot. Cations: Organic Review:

Anions: Radiochemistry Review: me/l Cations:

me/l Anions: id lotal: Microbiology Keview:

lot. Alk.	163 mg/1	TDS @ 180C	202 mg/1
1-Arsenic	22.0 ug/1	1-Barium	0.069 mg/1
I-Cadmium	是6部LOCKLean	I-Chromium	<5.0 ug/1
1-Copper	134:0 ug/1	1-lron	4.0 mg/1
I'-Lead	25_0-ug/1	l'-Manganes	11.0 ug/1
Mercury	0.2 ug/1	1-Selenium	<0.5 ug/l
I-Silver	(2.0 ug/1	ĭ-∠inc	\$ 800 0 2ug/1

PACIFIC MINE PORTAL FLOW 200 YDS. BELOW PORTA UINTA NATIONAL FOREST 88 W 100 N UT 84603 377-5780 PROUD

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PACIFIC MINE PORTAL FLOW 200 YDS. BELOW PORTA

Site ID:

Source: 00

Cost Code: 3508 Lab Number: 88028 Lab Number: 8802862 Type: 04 Sample Date: 88/05/18 Fime:

Date of Review and QA Validation Inorganic Review: 88/06/22

Tot. Cations:

Tot. Anions: and lotal:

91 me/l Cations:

91 me/l Anions:

Organic Review: Radiochemistry Review: 3.0 Microbiology Review:

Tot. Alk.	152 mg/1	FDS @ 180C	202 mg/1
1-Arsenic	24.0 ug/l	T-Barium	0.11 mg/l
ſ-Cadmium	\$90ug71	Γ-Chromium	(5.0 ug/1
1-Copper	62.0 ug/1	T-lron	6.6 mg/l
I-Lead	180.0:ug/1	T-Manganes	23.0 ug/l
Mercury	(0.2 ug/l	T-Selenium	<0.5 ug/l
ſ-Silver	<2.0 ug/l	Γ-Zinc	1300.0-ug/11

PACIFIC PORTAL AT CREEK (MARKINGS WIPED OFF UINTA NATIONAL FOREST 88 W 100 N PROVO U1 84603 377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PACIFIC PORTAL AT CREEK (MARKINGS WIPED OFF

Site 10:

Source: 00

Cost Code:

3508

Lab Number: 8802859 Type: 04 Sample Date: 5/18/88

l'ime:

Date of Review and QA Validation

Inorganic Review: Organic Keview:

lot. Cations:

t. Anions: and lotal:

me/1 Cations: me/1 Anions: Radiochemistry Review: Microbiology Keview:

Tot. Alk.	164 mg/1	rps @ 180C	200	mg/1
1-Arsenic	22.5 ug/l	1-Barium	0.28	mg/1
f-Cadmium	33-1%0g716-	I-Chromium	(5.0	ug/1
1-Copper	60 0 ug/1	1-1ron	5.3	mg/l
I'-Lead	4000;0 ug/1]	l'-Manganes	23.0	ug/l
Mercury	0.63 ug/1/	1-Selenium	(0.5	ug/l
T-Silver	15:0 ug/1	ľ-Zinc	1600.0	ug/1',

PAC.MINE NW PORTAL PIPED OUT OF MINE UINTA NATIONAL FOREST 88 W 100 N PROVO U1 84603

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PAC.MINE NW PORTAL PIPED OUT OF MINE

Site 10:

Source: 00

Cost Code:

3508

Lab Number: 8802856 Type: 04

Sample Date: 88/05/18 | Time: 10:10

Date of Keview and QA Validation

Inorganic Review: Organic Keview:

lot. Cations: t. Anions: ind lotal:

me/l Cations: me/l Anions:

Radiochemistry Review: Microbiology Review:

lot. Alk.	198 mg/1	TDS @ 180C	208 mg/1
1-Arsenic	1.0 ug/1	1-Barium	0.15 mg/1
I-Cadmium	<1 ug/1 ·	I-Chromium	(5.0 ug/1
1-Copper	<20.0 ug/1	1-1ron	0.091 mg/1
I-Lead	\$60 O 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	l'-Manganes	19.0 ug/1
Mercury	<0.2 ug/1	1-Selenium	(0.5 ug/1
I-Silver	<2.0 ug/1	l'-∠inc	78.0 ug/1

PACIFIC N TAILING UINTA NATIONAL FOREST 88 W 100 N U1 84603 PROVO

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: PACIFIC N FAILING

Site 10:

Cost Code: 3508 Lab Number: 8802860 Type: 04 Sample Date: 88/05/18 Time:

Source: 00

Date of Keview and QA Validation

Inorganic Review: Organic Review:

lot. Cations:

Anions:

me/l Cations: me/l Anions: Radiochemistry Review:

Microbiology Keview:

Tot. Alk. 1-Arsenic 1-Cadmium 1-Copper 1-Lead Mercury	21 mg/1 90.0 ug/1 	IDS @ 180C 1-Barium 1-Chromium 1-Iron 1-Manganes 1-Selenium	140 mg/1 0.15 mg/1 <5.0 ug/1 13.0 mg/1 48.0 ug/1 1.0 ug/1
I-Silver	45,0 ug/1	ſ-∠inc	7/700:0 ug/1

LOWER PAC.MINE PORTAL ACROSS STREAM FROM BAKE UINTA NATIONAL FOREST

88 W 100 N

PROVO

UT 84603

377-5780

UTAH STATE HEALTH LABORATORY

Environmental Chemistry Analysis Report

Description: LOWER PAC.MINE PORTAL ACROSS STREAM FROM BAKE! (117:21)

Site ID: Source: 00

lot. Cations:

Site ID: Source: 00
Cost Code: 3508
Lab Number: 8802863 Type: 04
Sample Date: 88/05/18 Fime: 10:45

Anions: 109 me/1 Cations: d Total: 109 me/1 Anions:

Date of Review and QA Validation Inorganic Review: 88/06/22

Organic Review:

Radiochemistry Review: 3.6 Microbiology Review:

fot. Alk.	183	mg/1	TDS @ 180C	204	mg/1
1-Arsenic	<1.0	ug/l	1-Barium	0.036	mg/l
I-Cadmium	<1	ug/l	I-Chromium	(5.0	ug/1
T-Copper	<20.0	ug/l	T-lron	0.048	mg/1
[-Lead	(5.0	ug/l	ſ-Manganes	6.0	ug/1
Mercury	<0.2	ug/l	1-Selenium	(0.5	ug/1
T-Silver	(2.0	ug/1	ſ-/inc	(20.0	ug/1

NORTH FORK AMERICAN RIVER AT DUTCHMAN FLAT UINTA NATIONAL FOREST 88 W 100 N

PROVO

U1 84603

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: NORTH FORK AMERICAN RIVER AT DUTCHMAN FLAT

Sice 10:

Source: 00

Sice 10: Source: 00
Cost Code: 3508
Lab Number: 8802855 Type: 04
Sample Date: 88/05/18 Time: 16:10

<u>Date of Review and QA Validation</u> Inorganic Review:

Organic Keview:

lot. Cations: Anions:

me/l Cations: me/l Anions:

Radiochemistry Review: Microbiology Keview:

fot. Alk.	83 mg/l	TDS @ 180C	102 mg/1
1-Arsenic	2.5 ug/1	1-Barium	0.056 mg/1
I-Cadmium	<1 ug/1 -	I-Chromium	<5.0 ug/1
1-Copper	<20.0 ug/1 ~	1-lron	0.45 mg/1
I-Lead	60.0 ug/11 A	l'-Manganes	31.0 ug/1
Mercury	(0.2 ug/1	1-Selenium	<0.5 ug/1
I-Silver	<2.0 ug/1	l'−∠inc	77.0 ug/l

MARY ELLEN PORTAL UINTA NATIONAL FOREST 88 W 100 N PROVO UI 84603

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: MARY ELLEN PORTAL

Site 1D: Source: 00
Cost Code: 3508
Lab Number: 8802858 Type: 04
Sample Date: 88/05/18 Time: 15:00

lot. Cations:

Anions: me/l Cations: id Total: me/l Anions:

Date of Review and QA Validation

Inorganic Review: Organic Keview:

Radiochemistry Review: Microbiology Keview:

lot. Alk.	36 mg/l	TDS @ 180C	206	mg/1
1-Arsenic	100.0 ug/1	1-Barium	0.019	mg/I
I'-Cadmium	4-ug/11 -	I'-Chromium	(5.0	ug/1
1-Copper	40.0 ug/1	1-Iron	9.9	mg/1
I'-Lead	10.0 ug/1	l'-Manganes	140.0	ug/1
Mercury	<0.2 ug/1	1-Selenium	(0.5	ug/1
1-Silver	<2.0 ug/1	l'-∠inc	1200.0	ug/1

MARY ELLEN CREEK 1/4 MILE BELOW MINE AREA UINTA NATIONAL FOREST 88 W 100 N

PROVO

UT 84603

377-5780

UTAH STATE HEALTH LABORATORY Environmental Chemistry Analysis Report

Description: MARY ELLEN CREEK 1/4 MILE BELOW MINE AREA

Site 1D:

Source: 00

Cost Code:

350B

Lab Number: 8802861 Type: 04 Sample Date: 88/05/18 Fime:

Date of Review and QA Validation

Inorganic Review: 88/06/22 Organic Review:

lot. Cations: Tat Anions:

i lotal:

55 me/l Cations:

Radiochemistry Review:

55 me/l Anions: 1.8 Microbiology Review:

fot. Alk.	92 mg/1	TDS @ 180C	132 mg/l
T-Arsenic	<1.0 ug/l 2	1-Barium	0.039 mg/1
T-Cadmium	2 ug/1 4 - 1	f-Chromium	<5.0 ug/1
T-Copper	42.0 ug/1 (T-Iron	1.1 mg/l
T-Lead	40.0 ug/1	l'-Manganes	46.0 ug/l
Mercury	<0.2 ug/l	1-Selenium	<0.5 ug/1
T-Silver	<2.0 ug/l	ſ-∠inc	\$310.0 ug/1



Department of Plant, Soil and Insect Sciences

College of Agriculture

P.O. Box 3354 Laramic, WY 82071-3354 Phone: (307) 766-3103 Fax: (307) 766-3379

24 July, 1992

Mr. Paul Skablund Hydrologist Uinta National Forest 88 West 100 North Provo, UT 84601 JUL 29 1992

Dear Mr. Skablund:

We would like to inform you of a research project that will likely be of significant interest to your organization. As you well know, mitigation of water pollution in the western United States is a matter of increasing concern. In particular, the mitigation of heavy-metal contaminated mineland effluent is an area of intense environmental interest. Heavy-metal effluent from hard rock mines at high elevations presents unique challenges in terms of cost, accessibility, and ecological damage. However, in the last two years we have succeeded in the first phase of a three phase program to develop a wetland system capable of effectively treating effluents which contain a range of heavy metals.

We have discovered a number of plants that accumulate heavy metals, including arsenic, lead, copper, zinc, and cadmium). Our reference metal, copper, is accumulated at a concentration 1,000 to 4,000 times that found in the water; one remarkable sedge that grows in the Rocky Mountains accumulates over 25,000 times the background concentration of this metal. All of these plants have been successfully cultivated in the greenhouse, where we are continuing to study the ecological parameters which optimize accumulation rates. In addition, we have found two fungi that are capable of removing 30-40% of the heavy metal from a liquid medium contaminated with 100 ppm copper in 9 days. The primary mechanism of removal appears to be metabolic chelating of the metal, which has particularly promising applications for *in situ* mitigation.

The next phase of our work will be small-scale experimental wetlands on contaminated mine sites. These "micro"-wetlands will be used to systematically vary ecological conditions and plant/fungi associations to determine the optimal system for heavy metal mitigation at a particular site. The final phase will involve the full-scale development of a wetland or series of wetlands capable of providing effective treatment of contaminated effluent. At this time, we are seeking funding for the initiation of the last two phases of this program. It appears that most, high altitude habitats would be amenable to this treatment program, and we are highly optimistic that mitigative wetlands will provide an extremely cost-effective management tool.

Enclosed you will find a more detailed presentation of the results to date and our future plans. If you are interested in further information or discussing the possibility of full or partial funding, please write to us or call (307) 766-3103 and ask for any of us.

Sincerely,

Stephen E. Williams Professor, Soil Science Nancy K. Culb Research Associate, Plant Science

Jeffrey A. Lockwood Associate Professor,

Entomology



Title Development of High Mountain Plant Communities as Wetland Mitigation Systems for Heavy Metal Mine Effluent.

Present Duration January 1991 - February 1993

Background

Heavy metal pollution from mine effluent is a serious and widespread problem in the western states. Although considerable work has documented the impact of heavy metals on aquatic and riparian flora, prior to our research there were no published, comprehensive field studies of the impacts of metals on high mountain stream plant communities. As a consequence the potential for manipulating high-elevation wetland plants and fungi for mitigation of mine effluent was unknown, although plant communities had proven to be effective filters of heavy metals in other ecosystems (Cairns 1980, Brooks et al. 1985).

It is well known that wetland plants can act as pollutant filters, collecting and holding nutrients, sediment, silt and other natural and anthropogenic pollutants, including heavy metals. There are several reasons why heavy metals are trapped in wetlands. Some metals will precipitate in the anaerobic zone, commonly present in wetlands. The high levels of decaying organic matter will further chelate many of these same metals. Many plants and microbiotic organisms that tolerate heavy metals also accumulate them in their tissues. At low elevations, the value of wetlands in sequestering iron and manganese from mine drainages has already been recognized (Holbrook and Maynard 1985, Gerber et al. 1985), and efforts have been made to construct wetlands for the purpose of trapping heavy metals (Gerber et al. 1985).

An effort to use wetlands to mitigate heavy metal mine discharge in high mountain streams is unique in that: 1) most wetland projects on abandoned mine lands have had the restoration of ' wildlife habitat, not mitigation of effluent, as the primary goal (Taub 1969, Brooks et al. 1985, Cairns 1987), 2) no wetland restoration projects have been developed at high elevations, and 3) wetland projects have been used to mitigate impacts of acids, iron and manganese, but the management of copper, zinc, lead and arsenic has not been attempted. Effective mitigation of mine effluent with wetlands generally includes integration of physical, chemical, and biological parameters (Brooks et al. 1985). In this context, it seems unlikely that manipulation of wetlands alone will completely reduce the heavy metal content of mine effluent flows. However, strategic management of plant and fungal communities is likely to play a significant role in a comprehensive ecological effort (e.g., settling ponds, impoundment, precipitation, etc.). Following physical, chemical, and biological treatments, wetland systems may be expected to function effectively in a management program.

Objectives

This wetland mitigation project has been in progress for 1.5 years. To date, the objective of this study has been to discover wetland plants and fungi that could tolerate and sequester heavy metals in their tissues. Both the plants and the environments in which they reside were examined. Reproductive and growth requirements were investigated.

Site Selection

During the summer of 1992, sites were assessed for their potential use in this study. Elevation, presence of mine effluent flowing across fairly horizontal gradients that contained wetland plants and accessibility were the main criteria used for selecting the areas. The sites chosen for study in summer 1992 were as follows, 1) Hughesville-Barker Block P Mine and Mill tailings in Lewis and Clark National Forest, Montana, 2) The Ontario Mine in Helena National Forest, 3) The Independence and McClaren mines in Gallatin National Forest, Montana, 4) Kirwin Mine west of Meeteetse in Wyoming, 5) Ferris-Haggarty Mine in the Sierra Madres, southern Wyoming, 6) Pacific and Mary Ellen Mine in the Wasatch Mountains, Utah. Additional sites that were assessed but rejected for the purposes of this study were the Mike Horse Mine in Helena National Forest, Montana and the Lower Bog Mine in the Wasatch Mountains, Utah.

Methods

Community and habitat analysis were undertaken on the chosen study sites. The Daubenmire quadrat method was used to identify the dominant and subordinant species. Diversity was quantified by counting number of plant species per site. The site was mapped on a 7.5 minute quad. Slope and aspect were determined with a clinometer and compass. The topographic position of the site was determined (crest, upper slope, mid-slope, lower slope, valley bottom, bench or terrace, saddle or gap). It was determined whether lighting was open, partial, filtered, or Soil moisture was described as inundated shaded on the site. (hydric), saturated (wet-mesic), moist (mesic), dry-mesic, or dry (xeric). Elevation was determined using a topographic map. The environment of the site was described (sand or gravel bar; wet meadow dominated by grasses; marsh dominated by sedges or rushes; swamp dominated by shrubs or trees; bog mire [mosses in acidic, wet peat soil]; fen mire with vascular plants in alkaline, wet peat soil; swale with moist surface soil; seep; terrace within three vertical feet or 100 feet of running surface water; snow catchment area; floating or quaking vegetation mat). Soil was collected to quantify pH, N, P, K and heavy metal composition. It was also collected for mycological sampling. The pH of the water on each site was tested, and samples were collected for heavy metal analysis in the laboratory.

Voucher specimens were collected of all the different species on each site. Vigor was described for each species. The reproductive fitness of each species in the most heavily impacted area was ascertained. Evidence of hybridity, disease, and symbiotic or parasitic relationships was noted. Plants were collected for heavy metal analysis, copper toxicity and reproductive studies in the greenhouse.

Vascular plants were identified using the Rocky Mountain Herbarium. Mosses were sent to the Clinton Herbarium in Buffalo, New York, to be identified. The pH, N, P, and K of the soil was determined by using a LaMotte soil testing kit. Plant available heavy metals were extracted from the soils using the ABDTPA method (Soltanpour and Schwab, 1977). Heavy metals were extracted from the plants via nitric acid digests (Havlin and Maynard, 1985). The University of Wyoming soil-testing lab analyzed duplicate samples of the water, soil and plants for Cu, Zn, Cd, Pb, Hg and As via inductively coupled plasma spectrometry.

Soil fungi from mines with soils containing more than 100 ppm of copper were plated on an agarose medium containing 100 ppm of copper and analyzed for frequency and dominance. Dominant and subdominant species were identified. Dominant species were grown in a liquid medium containing between 80 and 100 ppm of copper. The medium was analyzed for copper before innoculation and after a period of nine days. The fungi that grew in the culture was also analyzed after nine days.

Summary of Project Work Done to Date

To date, this study has shown that wetlands containing both plant and fungal species can be used very effectively to prevent release of heavy metals into stream systems. Both abiotic and biotic factors act to prevent the movement of heavy metals. Soils, especially those high in organic matter, and of fine particle size, chelate heavy metals and hold them in place. Notable in our study was the fact that the soils found directly under the plants contained higher levels of heavy metals, than those soils which were bare (ranging from 2:1 at the Ontario Mine to 24:1 at Ferris-Haggarty). Plants also uptake these metals, preventing their escape.

In this study, many plants were shown to substantially accumulate various heavy metals. All species accumulated significant amounts of heavy metals relative to the amounts found in water. For copper, the highest accumulators in context of background levels in water and soil, were Pohlia annotina (a moss) accumulating 3,000 times the amount in water, and 48 times the amount in soil, Deschampsia cespitosa (a grass) accumulating 2,000 times the level in water, and 31 times the soil, Pohlia wahlenbergii (a moss) accumulating 14,800 times the level of copper in the water, and 7 times the background soil level, and Senecio fremontii (a forb) accumulating 31 times the level in water and 5 times the soil level.

For zinc, the highest accumulators were Pohlia wahlenbergii (a moss) accumulating 3,800 times the level in water and 10 times the soil level, Equisetum arvense (a horsetail) accumulating 1,100 times the amount of zinc in the water and 8 times the amount in the soil, Poa interior (a grass) accumulating 2,100 times the level in the water and 6 times the soil background level, and Agrostis exarata (a grass) accumulating 260 times the zinc level in the water, and 4 times the soil level.

The highest lead accumulators were Carex microptera (a sedge) accumulating 6,000 times the level in the water, and 3,000 times the soil, and Poa interior (a grass) accumulating 5,300 times the amount in the water, and 2,700 times the soil level.

Arsenic was most effectively accumulated in Pohlia wahlenbergii (a moss). It accumulated 3,200 times the level of arsenic in the water, and 1,100 times the soil level, Bryum lisae (a moss) contained 6,400 times the amount in the water, and 1,100 times the level in the soil, Epilobium glaberrimum (a forb) accumulated 5,800 times the amount in the water, and 1,000 times the soil level, and Carex scopulorum (a sedge) accumulated 26,400 times the amount in the water and 113 times the soil level.

Cadmium was accumulated best by Pohlia wahlenbergii (a moss) at 6,400 times the level in the water, and 2 times the level in the soil.

All of these plants were capable of accumulating multiple metals. The highest overall accumulator was Pohlia wahlenbergii. Plants were able to accumulate arsenic most effectively, followed by lead, copper, zinc and finally cadmium. Different species varied in their ability to accumulate different heavy metals, so a mix of different species would be best for introduction into man-made wetlands, where a spectrum of contaminants is present.

Deschampsia cespitosa was common on all of the copper sites.

Carex aquatilis, and Pohlia nutans tolerated both zinc and copper sites. Carex microptera was common on sites which were high in zinc, and Carex rostrata appeared on zinc, lead and arsenic contaminated sites.

All of the plants that were collected in the field have been established in the greenhouse. The seeds of sedges that were collected germinated fully within 6 days, after 2 months of cold stratification. Germination was poor and occurred over a period of 6 weeks to 3 months without stratification. The seeds of the Epilobiums germinated in 3-6 days without stratification. Mosses have survived well within misting benches or where humidity is kept at high levels. Laboratory toxicity studies have shown that all of the species will survive well in the most copper-polluted sites found during our study. Although we have yet to find the extreme level of copper fatal to the plants, this is clearly far above that present in the field.

All of the fungal species analyzed accumulated copper. These included Tolypocladium inflatum, Trichocladium sp., and Penicillium sp. nov. "A". Tolypocladium inflatum from the Ferris-Haggarty Mine accumulated an average of 2,800 ppm of copper. Tolypocladium inflatum from the Kirwin Mine accumulated 3,400 ppm of copper. Trichocladium sp. from the McClaren's Mine at Cooke City, MT accumulated an average of 900 ppm and Penicillium sp. nov. "A" from the Pacific Mine in Utah accumulated 1,600 ppm of copper. These amounts were accumulated out of 80-95 ppm copper-amended medium. The accumulation of the copper in the fungi did not account for all or most of the copper removed from the medium by Tolypocladium inflatum and Trichocladium sp. In the case of Tolypocladium inflatum, an average of 41% of the copper had been removed from the solution by the fungus at the end of the 9 day experimental period. Of the copper that was removed, 77% was removed by a mechanism other than accumulation by the fungi. Trichocladium sp., removed an average of 31% of the copper from the solution. Of this, only 3% was found in the fungus; the remaining 97% was taken out of solution by another mechanism. Based on the scientific literature, we hypothesized that both species are producing a metabolite that is chelating the copper and removing it from the As such, the potential for fungi to play a solution. significant role in copper mine effluent mitigation appears reasonably high.

Studies to be completed before February 1993

Field work for the summer of 1993 is in progress. The accumulation of heavy metals in the dominant species of plants on three study sites, over space and time is being examined at the Ontario Mine in Montana and the the Kirwin and Ferris-Haggarty mines in Wyoming. The shoots and roots of plants collected are being separated for heavy metal analysis, to examine possible impacts on wildlife. Seeds of candidate plants for wetland mitigation studies will be collected from the Pacific Mine in Utah. Growth and copper toxicity studies are being completed in the greenhouse. Studies of environmental parameters (pH and temperature) that may effect uptake of heavy metals by plants will be attempted in the fall. Analysis of the fate of copper after senescence of the plant will also be undertaken at this time. In vitro fungal studies to determine the mechanism of copper removal are continuing. A two year report of the project will be submitted to the Department of Environmental Quality, Abandoned Minelands by December 1st, 1992.

Proposed work after February 1993

The present research is designed to produce the technology necessary for wetland mitigation of copper mine effluent. main reason we are working on copper pollution is because of the serious environmental situation that exists at the Ferris-Haggarty mine in the Sierra Madres, Wyoming. However, in our study we have discovered that there were few mines in the west where the main contaminant in the effluent was copper. Many abandoned mines have effluent flows containing higher levels of zinc, lead or arsenic with copper as a secondary contaminant. All of the mines studied contained heavy metals in combination. To design wetland mitigation systems for these mines' affluents, it would be advisable to initiate plant and fungal studies of the other metals present on these mine sites, similar to what has been done with copper. The effects of these metals in combinations similar to what is present in the field should also be examined.

It would be useful to establish control sites that are similar in every other way to the heavy metal polluted sites so that variations between polluted and unpolluted sites could be analyzed. Study sites already established can be analyzed for ecological parameters over several summers.

Small-scale field studies should be initiated at a site before a full-blown wetland mitigation project is undertaken. Problems encountered in the small scale in situ studies can be solved before large monetary commitments are made. These field studies will be in the form of micro-wetlands created by shunting a part of a mine effluent flow into a level, dyked area. Each micro-wetland will have parameters varied using a statistically valid experimental design to optimize the information gained (plant assemblage, innoculation by fungi, soil amendments). Data will be gathered throughout the field season (see attached diagram of possible study design). The Ferris-Haggarty, Kirwin, Ontario, Mary Ellen, and/or Pacific mines would be good sites to set up these microwetlands, if owner permission can be obtained.

The possibility of using tolerant species of fungi for treatment of heavy metal mine effluent should be explored. The prospect of both mitigating heavy metal contaminated mine effluent and recovering heavy metals from some of the fungal species that we have discovered during this study is exciting. This primary treatment system could prove to be relatively low in initial cost and long-term maintenance.

Literature Cited

- Brooks, R. P., D. E. Samuel and J. B. Hill. 1985. Wetlands and water management on mined lands. Proceedings of a conference at the Pennsylvania State University. University Park, Pennsylvania.
- Cairns, J. 1980. The recover process in damaged ecosystems.
 Ann Arbor Science Publishers, Ann Arbor, Michigan.
- Cairns, J. 1987. Disturbed ecosystems as opportunities for research in restoration ecology. pp. 307-320 in W. R. Jordan M.E. Gilpin and J.D. Aber (eds.). Restoration ecology: A synthetic approach to ecological research. Cambridge University Press, New York.
- Gerber, D. W., J. E. Burris and R. W. Stone. 1985. Removal of dissolved iron and manganese ions by a Sphagnum moss system. Proceedings of a conference at the Pennsylvania State University. University Park, Pennsylvania, pp. 365-373.
- Havlin, J. L. & P. L. Soltanpour. 1980. A nitric acid plant tissue digest method for use with inductively coupled plasma spectrometry. Comm. in Soil Science and Plant Analysis 11: 969-980
- Holbrook, J. A. and B. R. Maynard. 1985. Needed wetland research: a federal perspective. Proceedings of a conference at the Pennsylvania State University. University Park, Pennsylvania, pp. 195-198.
- Soltanpour, P. N. and A. P. Schwab. 1977. A new soil test for simultaneous extraction of macro and micro nutrients in alkaline soils. Comm. Soil Sci. and Plant Anal. 8: 195-297
- Taub, F. 1969. Gnotobiotic models of freshwater communities. Vehr. Internat. Verein. Limnol. 17: 485-496



UINTA NATIONAL FOREST

JUN 0 1 1992

epartment of Plant, Soil and Insect Sciences College of Agriculture P.O. Box 3354 Phone: (307) 766-3103 Fax: (307) 766-3379

30 May 1992

Uinta National Forest Supervisor's Office Paul Skablund Forest Hydrologist 88 West 100 North Provo UT 84601

Dear Mr. Skablund:

Here is your part of the year-end report on the mitigation of heavy metal mine effluent by wetlands, that I promised you. I hope you will find the data I gathered at the Pacific and Mary Ellen mines informative. If you need any more information, please call me.

If it is all right with you, I would like to visit the Pacific Mine sometime in late August or September to gather seed heads off of the plants up there. I am going to grow larger amounts of these plants in the greenhouse for use in wetland mock-up studies. I will send you the data I gather from these studies.

I hope you will gain some insight from this report. The beaver pond at the Pacific Mine is doing a great deal to mitigate flow-through of heavy metals into the North Fork of the American Fork River. I am glad to hear that you are trying to keep people off of the mine tailings. They are heavily laced with a variety of heavy metals and probably would not be good for the health of anybody who spent a lengthy period of time on them, especially on a windy day.

Thank-you for your consideration,

Nancy Kastning-Culp Research Associate



Year End Report

On Mitigation Systems for Hard Rock Mine Effluent

In Utah

Investigators:
Nancy Kastning-Culp
Larry DeBrey
Jeff Lockwood
of the
Department of Plant, Soil and Insect Sciences
University of Wyoming

30 May 1992

Abstract/Summary of All Mines Studied

This study shows that wetlands can be used very effectively to prevent release of heavy metals into stream systems. Both abiotic and biotic factors act to prevent the movement of heavy metals. Soils, especially those high in organic matter, and of fine particle size, chelate heavy metals and hold them in place. Notable in our study was the fact that the soils found directly under the plants contained higher levels of heavy metals, than those soils which were bare (a minimum of 2:1 at the Ontario Mine, maximum 24:1 at Ferris-Haggarty). Plants also uptake these metals, preventing their escape.

In this study, many plants were shown to accumulate various heavy metals to a great degree. All species accumulated significant amounts of heavy metals in comparison to the amounts found in water. For copper the best accumulators as compared to background levels in water and available in soil, were Pohlia annotina (a moss) accumulating 3,032 times the amount in water, and 48 times the amount in soil, Deschampsia cespitosa (a grass) accumulating 1,979 times the level in water, and 31 times the soil, Pohlia wahlenbergii (a moss) accumulating 14,813 times the level of copper in the water, and 7.0 times the background soil level, and Senecio fremontii (a forb) accumulating 31 times the level in water and 5 times the soil level.

For zinc, the best accumulators were <u>Pohlia wahlenbergii</u> (a moss) accumulating 3,814 times the level in water and 10 times the soil level, <u>Equisetum arvense</u> (a horsetail) accumulating 1,120 times the amount of zinc in the water and 7.5 times the amount in the soil, <u>Poa interior</u> (a grass) accumulating 2,128 times the level in the water and 5.7 times the soil background level, and <u>Agrostis exarata</u> (a grass) accumulating 261 times the zinc level in the water, and 4.5 times the soil level.

The best lead accumulators were <u>Carex microptera</u> (a sedge) accumulating 5,954 times the level in the water, and 2,977 times the soil, and <u>Poa interior</u> (a grass) accumulating 5,347 times the amount in the water, and 2,674 times the soil level.

Arsenic accumulated best in Pohlia wahlenbergii (a moss). It accumulated 3,221 times the level of arsenic in the water, and 1,073 times the soil level, Bryum lisae (a moss) contained 6,443 times the amount in the water, and 1,073 times the level in the soil, Epilobium glaberrimum (a forb) accumulated 5,814 times the amount in the water, and 969 times the soil level, and Carex scopulorum (a sedge) accumulated 26,432 times the amount in the water and 113 times the soil level. Cadmium was accumulated best by Pohlia wahlenbergii (a moss) at 6,393 times the level in the water, and 1.7 times the level in the soil.

All of these plants were capable of accumulating multiple metals. The best overall accumulator was <u>Pohlia wahlenbergii</u>. Plants were able to uptake arsenic most effectively followed by lead, copper, zinc and then cadmium. Different species vary in their ability to accumulate different heavy metals, so a mix of different species would be best for introduction into man-made wetlands, where a range of contaminants is present.

<u>Deschampsia</u> <u>cespitosa</u> was common on all of the copper sites. <u>Carex</u> <u>aquatilis</u>, and <u>Pohlia</u> <u>nutans</u> tolerated both zinc and copper sites. <u>Carex</u> <u>microptera</u> was common on sites which were high in zinc, and <u>Carex</u> <u>rostrata</u> appeared on zinc, lead and arsenic sites.

Field and Lab Methods

In the field, community and habitat analyses were undertaken. The Daubenmire quadrat method was used to identify the first and second most dominant species. Diversity was quantified by counting number of species per site. The site was mapped on a 7.5 minute quad. Slope and aspect were determined with a clinometer and compass. The topographic position of the site was determined (crest, upper slope, mid-slope, lower slope, valley bottom, bench or terrace, saddle or gap). It was determined whether lighting was open, partial, filtered, or shaded on the site. Soil moisture was described as inundated (hydric), saturated (wet-mesic), moist (mesic), dry-mesic, dry (xeric). Elevation was determined using a topographic map. The environment of the site was described (sand or gravel bar; wet meadow dominated by grasses; marsh dominated by sedges or rushes; swamp dominated by shrubs or trees; bog mire [mosses in acidic, wet peat soil]; fen mire with vascular plants in alkaline, wet peat soil; swale with moist surface soil; seep; terrace within three vertical feet or 100 feet of running surface water; snow catchment area; floating or quaking vegetation mat). We collected soil to quantify pH, N, P, K and heavy metal composition. We also collected soil for mycological sampling (10 samples from the most heavily impacted site). We tested the pH of the water on the site before collecting it for heavy metal analysis in the lab.

Voucher specimens were collected of all the different species on the site. Vigor was described for each species. The reproductive fitness of each species in the most heavily impacted area was ascertained. We looked for evidence of hybridity, disease, and symbiotic or parasitic relationships. We then collected plants for heavy metal analysis and live-plant greenhouse studies.

Vascular plants were identified using the microscopes at the Rocky Mountain Herbarium. Mosses were sent to the Clinton Herbarium in Buffalo, New York, to be identified by Patricia Eckel, a western moss specialist. The pH, N, P, and K of the soil was determined by using a LaMotte soil testing kit. Plant available heavy metals were extracted from the soils using the ABDTPA method (Soltanpour, 1977). Heavy metals were extracted from the plants via nitric acid digests (Havlin, 1980). The University of Wyoming soil-testing lab analyzed duplicate samples of the water, soil and plant samples for Cu, Zn, Cd, Pb, Hg and As using an inductively coupled plasma spectrometer.

Mycological soil samples from Kirwin Mine and Ferris-Haggarty Mine in Wyoming, McClaren's Mine in Montana, and the Pacific Mine in Utah were diluted 1:100 in sterile, deionized water, and 1 ml each of each sample was dispensed onto three plates of 10, 100, and 1000 ppm copper-enriched Martin's Medium. Colonies were counted and hyphal tip picks were made from the 100 ppm copper-enriched Martin's into 100 ppm copper enriched potato-dextrose agar tubes.

Utah

Mary Ellen Mine - Wasatch Mountains - Wasatch County

Glacial of Yough of The Mary Ellen mine is 2.1 miles up a four wheel drive road. It is in a cirque basin surrounded by peaks of the Wasatch mountains. The Mary Ellen gulch area was extensively mined. Mine effluent originates from the side of a hill, flows past tailings piles and into Mary Ellen Creek. Forest service tests in 1981 and 1982 show zinc to be the main contaminant followed by copper. The soil around and under the effluent is stained a bright orange-red. Several plants grow directly in and by the effluent including a moss, Epilobium (a willow-wort), Mimulus (also called monkeyflower), Carex (a sedge), and Juncus (a rush). See Figure 3 for the map of the Mary Ellen microsite.

Microsite Information:

Microsite Code: ME1

UT: Utah County. Wasatch Mountains ca 20 air miles north of Provo, ca 1 air mile east-south-east of East Twin Peak at the Mary Ellen Mine seep (T3S R3E S22). From Provo: Go to American Fork Canyon. Follow road to Dutchmans Flat. Go up the center 4 wheel drive road into Mary Ellen Gulch. You should be on the right side of the creek. Turn left at all forks. The road dead ends into the mine.


Habitat and Community Information:

This site is located at mid-slope by a seep. The elevation is 9,500 ft. The slope is facing south-east. Lighting is full.

Basic soil and water chemistry: pH of Water: 6 Soil pH: 6 N: 5ppm P: 50ppm K: <50ppm

The dominant species is <u>Bryum lisae</u> var. <u>cuspidatum</u> (a moss) and <u>Poa interior</u> (a grass) is subdominant. Nine species were sampled by quadrat on this site.

Figure 3
Map of Mary Ellen Microsite



Individual Species Information:

Date collected: 07/30/91 Chem Code: ME1BL Species: Bryum lisae var. cuspidatum

Vigor: Exceptionally vigorous Cover: 65 % Frequency: 90 %

Chem Code: ME1CC

Species: Corydalis caseana Date collected: 07/30/91
Vigor: vigorous Cover: 0 % Frequency: 0 %
Comments: This species did not appear in thrown quadrats.

Chem Code: ME1CM

Species: Carex microptera

Vigor: vigorous

Date collected: 07/30/91

Cover: 0 % Frequency: 0 % Comments: This species did not appear in thrown quadrats.

Chem Code: ME1EG

Species: Epilobium glaberrimum Date collected: 07/30/91 Vigor: Vigorous Cover: 0 % Frequency: 0 % Comments: This species did not appear in thrown quadrats.

Chem Code: ME1PI

Species: Poa interior Date collected: 07/30/91 Cover: 6 % Frequency: 20 % Vigor: Normal

Heavy Metal Chemistry (mg/kg):

Plants:

Cu	Zn	Code: ME1BL Cd 17.66	Pb	lisae Hg <5	As 644.2
Cu	Zn	Code: ME1CC Cd 2.475	Pb	alis <u>casean</u> Hg <5	a As 87
	Zn	Code: ME1CM Cd 2.975	Pb	Hg	As
Cu	Zn	Code: ME1EG Cd 13.65	Pb	Hg	rimum As 581.35
Soil:	06 (25 (01	C. l. WEIG	D- W-	FILL	
Cu	Zn	Code: ME1S Cd 3.84	Pb	Hg <.2	As

Water:

Date:	06/25/91	Code: ME1W		Mary Eller	mine effluent
Cu	Zn	Cd	Pb	Hg	As
.02	1.55	<.01	<.1	<.1	<.1

These plants accumulated high levels of both zinc and arsenic.

Bryum lisae (a moss) accumulated 806 times the amount of zinc in the water, and 2.9 times the amount of zinc in the soil. It accumulated 6,442 times the amount of arsenic in the water and 1,074 times the amount of arsenic in the soil. It also accumulated copper at a rate 8.3 times the level in the soil, and 12,825 times the level in the water.

Epilobium glaberrimum (a willow-wort) accumulated 675 times the amount of zinc in the water and 2.4 times the amount of zinc in the soil. This plant accumulated 5,814 times the amount of arsenic as in the water, and 969 times the amount of arsenic in the soil. It also contained 6.06 times the amount of copper in the soil and 9,325 times the level in the water.

Pacific Mine - Wasatch Mountains

The Pacific Mine is located on the left fork of the main dirt road originating from Tibble Fork Reservoir. The mine effluent originates from a hillside, pools in a flat area, and flows through a tailings pile before entering into a beaver-caused wetlands complex. This area is a perfect study site in which to test the hypothesis that wetlands systems which include heavy metal tolerant or accumulating species of plants would mitigate heavy metal effluent. In fact, University of Wyoming water quality studies indicate that water quality increases dramatically after running through the wetlands and beaver dam. This site also yielded the species which accumulated the most heavy metals. wahlenbergii var. glaciale (a moss) accumulated 13,004 ppm of zinc. This is 3,813 times the background water level and 10 times the background soil levels. This moss also accumulated 1,185 ppm of copper, which is 1,481 times the background level of water and 7 times the background level of the soil. All of the metals for which we tested were accumulated by this plant. The area impacted by the mine drainage is rich in vegetation, including two species of Carex (a sedge), a species each of Juncus (a rush), Poa (a grass), and Epilobium (a willow-wort), and a species of moss. The main contaminant in the effluent is zinc. See Figure 4 for map of Pacific Mine microsites.

Microsite Information:

Microsite Code: PM1

UT: Utah County. Wasatch Mountains ca 20 air miles north-northeast of Provo, ca 1 air mile west of Miller Hill at the Pacific Mine (T3S R3E S22). From Provo: Go to American Fork Canyon. Follow the road to its main fork above all named flats and go left. The Pacific Mine is on the left approximately 1 mile up the road. This microsite is the seep mouth on the Pacific Mine.

Habitat and Community Information:

This site is located on a lower slope by a seep. The elevation is 7,800 ft. The slope is east-facing. Lighting is Full.

Basic soil and water chemistry: pH of Water: 6 Soil pH: 7 N: 20ppm P: 25ppm K: 150ppm

The dominant plant species is $\underline{Carex\ microptera}$ (a sedge), and $\underline{Juncus\ ensifolius}$ (a rush) is subdominant. Seven species were sampled by quadrat on this site.

The dominant soil microfungi is

Figure 4
Map of Pacific Mine Microsites



Individual Species Information:

Chem Code: PM1CM

Species: Carex microptera
Vigor: Vigorous

Date collected: 07/31/91
Cover: 37 % Frequency: 40

Chem Code: PM1JE

Species: Juncus ensifolius
Vigor: Vigorous

Date collected: 07/31/91
Cover: 26 % Frequency: 30 %

Chem Code: PM1PW Species: Pohlia wahlenbergii Date collected: 07/31/91

var. glaciale

Vigor: exceptionally vigorous Cover: 3.4% Frequency: 20 % Comments: This moss is especially loaded with heavy metals.

Chem Code: PM1PI

Date collected: 07/31/91 Cover: 8 % Frequency: 20 % Species: <u>Poa interior</u> Vigor: Vigorous

Heavy Metal Chemistry (mg/kg):

Plants:

Date: 07/31/91 Cu Zn 556.5 5403	Code: PM1CM Cd 69.97	Carex microptera Pb Hg As 595.35 <5 305.15
Date: 07/31/91 Cu Zn 237 2662.5	Code: PM1JE Cd 28.64	Juncus ensifolius Pb Hg As 242.35 <5 171.85
Date: 07/31/91 Cu Zn 724.5 7259	Code: PM1PI Cd 79.83	Pb. Hg As 534.7 <5 489.9
		Pb Hg As 388.05 <5 644.25
Soil:		
Date: 07/31/91 Cu Zn 166.96 1261.96	Code: PM1S Cd 74.99	Pacific mine seep Pb Hg As <.2 <.2 <.2 .6
Water:		
Date: 06/25/91 Cu Zn .12 4.64	Code: 'PM1W Cd <.01	At effluent mouth Pb Hg As <.1 <.1 <.1
Date: 07/31/91 Cu Zn .08 3.41	Code: PM1W Cd .02	At effluent mouth Pb Hg As .1 <.1 .2
Date: 07/31/91 Cu Zn .08 3.32	Code: PM1WA Cd .01	effluent pool below mouth Pb Hg As .1 .1 .1
		Below tailings before dam Pb Hg As .6 .4 <.1

The Pacific Mine had unusually high amounts of copper, zinc, lead and cadmium. Some arsenic and mercury were also present.

Pohlia wahlenbergii (a moss) accumulated notable levels of copper, zinc and arsenic. It accumulated 14,813 times the background level of copper in the water, and 7.1 times the amount in the soil. It also accumulated 3,814 times the amount of zinc in the water and 10.3 times more than the amount in the soil. Arsenic was accumulated at 3,221 times what was in the water and 1,074 times the amount in the soil.

Carex microptera (a sedge) accumulated notable levels of copper, zinc and lead. It accumulated 6,956 times the amount of copper as was in the water, and 3.33 times what was in the soil. Zinc was accumulated at 1,584 times what was in the water, and 4.28 times what was in the soil. Lead accumulated at 5,954 times the amount in the water and at least 2,977 times what was in the soil.

<u>Juncus ensifolius</u> (a rush) accumulated 781 times the amount of zinc in the water and 2.11 times the amount in the soil. Copper accumulated at 2,962 times the level in the water, and 1.42 times the amount available in the soil.

Poa interior (a grass) accumulated copper, zinc and lead. Copper was accumulated at 9,056 times what was in the water, and 4.34 times the amount in the soil. Zinc was accumulated at 2,128.74 times the amount in the water, and 5.7 times the amount in the soil. Lead was accumulated at 5,347 times the amount in the water, and at least 2,674 times the amount in the soil.

Mycology:

Pacific Mine: Microhabitat samples A-C were predominantly silty to sandy and were orangish in color. Microhabitat samples D-J were high in partially decayed organic matter and were dark brown in color. No growth occurred on the 1000 ppm copper-amended medium. Growth was slow but with many colonies on the 100 ppm copperamended medium. Growth was so profuse on the 10 ppm medium at the 1:100 dilution that separation of colonies for counting and picking for most of the different soil samples was impossible. species were isolated from the 100 ppm P-D-A copper-amended tube slants and were identified. There was an average of 39.3 colonies/plate on the 100 ppm copper-amended Martin's medium. The dominant species is an undescribed *Penicillium*. The description is being developed by Christianson and Tuthill. They are presently characterizing it as Penicillium sp. nov. "A", and have also discovered it in iron rich mine tailings. Its frequency was 78% (Table 1). Another undecribed *Penicillium* (sp. #1) from the raistickii series was present, along with *P. janthinellum*. Quantitative analysis of the fungi showed that it accumulated an average of 1572 ppm of copper. The liquid medium in which the fungi was grown showed no drop in copper in solution during the duration of the experiment. It is felt that this is an artifact resulting from dehydration of the medium. Quantitative analysis for copper is going to be repeated.

Table 1. Average colony counts and frequency of identified and unknown species from Pacific Mine growing on 100 ppm Martin's Medium.

	Penicillium sp. #1	Penicillium janthinellum	Penicillium sp. nov. "A"
Colony counts	4	4	28
Frequency	11%	11%	78%

Table 2. Analysis of fate of copper in potato-dextrose shake cultures amended with copper and innoculated with dominant species of fungi from the Pacific Mine.

Rep #	Sp.	Site	ppm Cu pre	ppm Cu post	Dif.	% dif.	g. fun.	ppm Cu fun.	pH med aft
10	dic.	PM1	84	83.5	0.5	1%	.517	1655	3.4
11			82	82	0.0	0%	.513	1541	3.3
12	15	46.00	82	84.5	-2.5	-3%	.507	1519	3.2

Rep # = Replicate #; Sp. = Species; Site = Site where species was collected; ppm Cu pre = original parts per million of copper in solution before innoculating with the fungus; ppm Cu post = parts per million of copper after nine days of fungal growth in the shake culture; Dif. = ppm Cu pre - ppm Cu post; % dif. = (dif./ppm Cu pre)100; g. fun. = grams dry weight of fungus used to analyze ppm Cu in the fungus; ppm Cu fun. = parts per million of copper in the fungus; pH med aft = pH of the medium after 9 days of fungal growth; Con. = control with no fungus added.

Microsite Code: PM2

UT: Utah County. Pacific Mine swamp (T3S R3E S22). See PM1 for directions. This microsite is the beaver pond where the seep feeds in.

Habitat and Community Information:

This site is located in a valley bottom in a marsh. The elevation is 7,800 ft. The slope is east-facing. Lighting is full.

Basic soil and water chemistry: pH of Water: 7 Soil pH: 7 N: 5ppm P: 50ppm K: 60ppm

The dominant species is <u>Carex rostrata</u> (a sedge), and <u>Equisetum arvense</u> (a horsetail) is subdominant. Two species were sampled by quadrat on this site.

Individual Species Information:

Chem Code: PM2CR

Species: Carex rostrata Date collected: 07/31/91 Vigor: exceptionally vigorous Cover: 91 % Frequency: 100%

Chem Code: PM2EA

Species: Equisetum arvense
Vigor: vigorous

Date collected: 07/31/91
Cover: 9 % Frequency: 20 %

Heavy Metal Chemistry (mg/kg):

Plants:

Date: Cu 102	Zn	Code: PM2CR Cd 10.71	Pb Hg 149.35 <5	As 47.3
Date: Cu 179	07/31/91 Zn 4079	Code: PM2EA Cd 49.7	Equisetum arve	ense As 93.7
Soil:				
Date: Cu 28.16	Zn	Code: PM2S Cd 23.79	Pacific mine Pb Hg 29.24 <.2	As .5
Water:				
Date: Cu .14	06/25/91 Zn 3.64	Code: PM2W Cd .04	In beaver pond Pb Hg <.1 <.1	1. As <.1

Carex rostrata (a sedge) and Equisetum arvense (a horsetail) both accumulated zinc from the beaver pond. Carex rostrata accumulated 360 times the amount of zinc in the water, and 2.4 times the amount in the soil. Equisetum arvense accumulated 1,121 times the amount of zinc in the water and 7.5 times the amount of zinc in the soil. It also accumulated 6.4 times the amount of copper in the soil and 1,279 times the amount of copper in the water.

Bibliography

Havlin, J. L. & P. N. Soltanpour. 1980. A nitric acid plant tissue digest method for use with inductive coupled plasma spectrometry. Comm. Soil Sci. and Plant Anal. 11: 969-980

Skabelund, P. 1989. Uinta National Forest water quality data. Unpubl. 5 pp

Soltanpour, P. N. & A. P. Schwab. 1977. A new soil test for simultaneous extraction of macro and micro nutrients in alkaline soils. Comm. Soil Sci. and Plant Anal. 8: 195-297

Mangum, Fred, Aquatic Ecosystem Analysis Laboratory, 1988.



"Preliminary Survey of Water Quality in Mine Drainage In Sheeprock Mountains and North Fork of the American Fork River", Merritt, James B., Provo, Utah, July 1988.

Appendix D

"Year End Report On Mitigation Systems For Hard Rock Mine Effluent In Utah", Culp, Nancy, Et Al, University of Wyoming, May 1992.

BIBLIOGRAPHY

Culp, Nancy, Et Al, University of Wyoming, May 1992, "Year End Report On Mitigation Systems For Hard Rock Mine Effluent In Utah".

Keetch, Fred, Date Unknown, American Fork Canyon Mining History.

Lidstone and Anderson, Inc. 1993, "American Fork Hydrology and Water Quality Study", Lidstone and Anderson, Inc.

Mangum, Fred A., 1988, "Aquatic Ecosystem Inventory", Macroinvertebrate Analysis, North Fork American Fork River and Mary Ellen Gulch Creek, Uinta National Forest, Aquatic Ecosystem Analysis Laboratory.

Merritt, Lavere B., July 1988, "Preliminary Survey of Water Quality in Mine Drainage In Sheeprock Mountains and North Fork of the American Fork River".

Skabelund, Paul, 1989, Uinta National Forest Water Quality, Unpubl. 5 pp.